



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 14, 2026 – 01:56 PM JST

PDB ID : 24IS / pdb_000024is
Title : alpha-1,2-glucosidase from Arthrobacter humicola A8F5, glucose complex
Authors : Yasukochi, R.; Fushinobu, S.
Deposited on : 2026-03-05
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	662	6% 74% 15% • 9%
1	B	662	6% 77% 13% • 7%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 9478 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called alpha-1,2-glucosidase from *Arthrobacter humicola* A8F5.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	605	Total	C	N	O	S	0	0	0
			4532	2861	822	836	13			
1	B	614	Total	C	N	O	S	0	0	0
			4598	2900	836	849	13			

- Molecule 2 is beta-D-glucopyranose (CCD ID: BGC) (formula: $C_6H_{12}O_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	159	Total O 159 159	0	0
4	B	141	Total O 141 141	0	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: alpha-1,2-glucosidase from *Arthrobacter humicola* A8F5



- Molecule 1: alpha-1,2-glucosidase from *Arthrobacter humicola* A8F5





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	104.46Å 107.09Å 110.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.07 – 1.90 49.07 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.07-1.90) 100.0 (49.07-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.63 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.215 , 0.263 0.225 , 0.268	Depositor DCC
R_{free} test set	4623 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å ²)	25.8	Xtriage
Anisotropy	0.113	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 35.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9478	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.45 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.9023e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	0/4648	1.27	25/6343 (0.4%)
1	B	0.76	0/4716	1.28	21/6435 (0.3%)
All	All	0.76	0/9364	1.27	46/12778 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	10
1	B	0	8
All	All	0	18

There are no bond length outliers.

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	611	GLU	CG-CD-OE1	-8.33	99.23	118.40
1	B	611	GLU	CG-CD-OE1	-7.91	100.20	118.40
1	B	221	THR	CA-CB-OG1	-7.76	97.96	109.60
1	B	427	ASP	CA-CB-CG	7.02	119.62	112.60
1	A	299	ASP	CA-CB-CG	6.94	119.54	112.60
1	A	611	GLU	CG-CD-OE2	6.88	134.22	118.40
1	A	335	HIS	CA-CB-CG	-6.65	107.15	113.80
1	A	427	ASP	CA-CB-CG	6.60	119.20	112.60
1	A	237	PRO	N-CA-CB	-6.48	99.85	103.22
1	B	261	ARG	N-CA-CB	6.40	119.39	109.85
1	A	538	ARG	NE-CZ-NH1	-6.32	115.18	121.50
1	B	611	GLU	CG-CD-OE2	6.29	132.86	118.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	538	ARG	CG-CD-NE	-6.28	98.17	112.00
1	B	75	LEU	N-CA-CB	-6.22	100.15	110.23
1	A	377	PRO	N-CA-CB	-6.20	96.75	103.25
1	A	251	ASP	CA-CB-CG	6.17	118.77	112.60
1	A	195	THR	CA-CB-OG1	-6.11	100.44	109.60
1	A	439	GLN	N-CA-CB	5.99	118.79	110.17
1	B	429	ARG	CB-CA-C	5.95	120.33	110.81
1	B	0	HIS	CA-CB-CG	5.93	119.73	113.80
1	A	494	ASP	CA-CB-CG	5.91	118.51	112.60
1	A	155	MET	CG-SD-CE	-5.86	88.00	100.90
1	B	0	HIS	CB-CA-C	5.86	121.23	110.10
1	B	145	ASP	CA-CB-CG	5.81	118.41	112.60
1	A	208	ARG	CB-CA-C	5.72	118.90	109.70
1	A	412	TYR	CA-CB-CG	5.63	124.04	113.90
1	A	378	ASP	CB-CA-C	-5.49	101.34	109.90
1	A	485	ARG	NE-CZ-NH1	-5.42	116.08	121.50
1	A	261	ARG	CB-CA-C	5.38	118.33	110.26
1	B	538	ARG	NE-CZ-NH1	-5.33	116.17	121.50
1	B	430	ASP	CA-CB-CG	5.32	117.92	112.60
1	B	300	THR	N-CA-C	-5.28	105.70	111.82
1	A	378	ASP	CA-CB-CG	5.27	117.87	112.60
1	B	485	ARG	NE-CZ-NH1	-5.26	116.24	121.50
1	B	458	LEU	N-CA-CB	5.23	117.65	110.07
1	A	590	GLU	CB-CG-CD	5.22	121.48	112.60
1	A	145	ASP	CA-CB-CG	5.22	117.82	112.60
1	B	391	ARG	NE-CZ-NH1	-5.19	116.31	121.50
1	B	23	ASP	CA-CB-CG	5.16	117.75	112.60
1	B	97	ARG	NE-CZ-NH1	-5.14	116.36	121.50
1	A	320	THR	CA-CB-OG1	-5.14	101.89	109.60
1	B	476	ARG	CG-CD-NE	-5.12	100.74	112.00
1	B	469	GLU	N-CA-CB	5.11	115.66	109.74
1	A	538	ARG	CD-NE-CZ	5.04	131.46	124.40
1	A	464	PHE	CA-CB-CG	5.03	118.83	113.80
1	B	505	ASP	CA-CB-CG	5.01	117.61	112.60

There are no chirality outliers.

All (18) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	231	ARG	Sidechain
1	A	312	ARG	Sidechain
1	A	374	ARG	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	429	ARG	Sidechain
1	A	438	ARG	Sidechain
1	A	476	ARG	Sidechain
1	A	487	ARG	Sidechain
1	A	538	ARG	Sidechain
1	A	555	ARG	Sidechain
1	A	9	ARG	Sidechain
1	B	180	ARG	Sidechain
1	B	208	ARG	Sidechain
1	B	312	ARG	Sidechain
1	B	383	ARG	Sidechain
1	B	487	ARG	Sidechain
1	B	538	ARG	Sidechain
1	B	73	ARG	Sidechain
1	B	97	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4532	0	4429	57	0
1	B	4598	0	4494	55	0
2	A	12	0	12	0	0
2	B	12	0	12	0	0
3	A	8	0	12	0	0
3	B	16	0	24	4	0
4	A	159	0	0	1	0
4	B	141	0	0	1	0
All	All	9478	0	8983	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (105) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:236:ARG:H	3:B:703:EDO:H22	1.25	0.96
1:B:239:ARG:NH1	1:B:252:ASN:OD1	2.09	0.86
1:A:465:ASP:OD2	1:A:476:ARG:NH2	2.08	0.86
1:B:465:ASP:OD2	1:B:476:ARG:NH2	2.10	0.84
1:B:63:LEU:HD11	1:B:127:VAL:HG21	1.66	0.76
1:A:2:THR:O	1:A:161:SER:HB3	1.87	0.74
1:A:63:LEU:HD11	1:A:127:VAL:HG21	1.70	0.72
1:B:236:ARG:N	3:B:703:EDO:H22	2.04	0.70
1:B:418:HIS:NE2	1:B:441:GLU:HG2	2.08	0.68
1:B:99:ARG:HH12	1:B:134:ARG:NH2	1.94	0.66
1:A:379:ASP:OD1	1:A:383:ARG:NE	2.29	0.65
1:A:32:SER:CB	1:A:73:ARG:HH22	2.09	0.65
1:B:598:LEU:HD13	1:B:644:ILE:HD11	1.77	0.65
1:A:303:GLU:HG3	1:A:381:VAL:HG22	1.77	0.65
1:A:25:ASP:OD1	1:B:9:ARG:NH2	2.29	0.64
1:A:180:ARG:HH11	1:A:185:SER:HB3	1.63	0.64
1:A:9:ARG:NH2	1:B:25:ASP:OD1	2.31	0.64
1:B:71:LEU:HD13	1:B:186:LEU:HD21	1.84	0.60
1:A:543:GLU:HA	1:A:555:ARG:HG2	1.84	0.59
1:A:598:LEU:HD23	1:A:644:ILE:HD11	1.83	0.59
1:A:628:PRO:HG2	1:B:101:LEU:CD1	2.34	0.58
1:A:71:LEU:HD13	1:A:186:LEU:HD21	1.86	0.57
1:B:596:ASP:OD1	1:B:648:MET:HE1	2.05	0.57
1:A:299:ASP:O	4:A:801:HOH:O	2.18	0.57
1:A:32:SER:HB3	1:A:73:ARG:HH22	1.69	0.56
1:A:628:PRO:HG2	1:B:101:LEU:HD12	1.87	0.56
1:A:208:ARG:HD3	1:B:558:ALA:O	2.06	0.55
1:A:588:ILE:HB	1:A:589:PRO:HD3	1.88	0.55
1:B:178:ARG:HH11	1:B:178:ARG:HG3	1.72	0.54
1:A:433:GLN:OE1	1:A:559:GLY:HA3	2.06	0.54
1:B:14:VAL:HG22	3:B:705:EDO:H11	1.89	0.54
1:A:513:ILE:HD12	1:A:514:PRO:HD2	1.90	0.54
1:B:89:GLU:OE1	1:B:91:GLN:NE2	2.38	0.54
1:B:646:GLN:HE21	1:B:652:LEU:HG	1.73	0.53
1:A:64:LEU:HD22	1:A:155:MET:HE3	1.91	0.53
1:B:421:SER:HB2	1:B:427:ASP:HB2	1.90	0.53
1:A:422:ASN:ND2	1:A:428:SER:O	2.40	0.53
1:A:257:VAL:HG13	1:A:273:ILE:HD12	1.91	0.52
1:B:378:ASP:HB3	1:B:381:VAL:HG23	1.92	0.52
1:A:99:ARG:HH12	1:A:134:ARG:NH2	2.09	0.51
1:A:382:ILE:HG21	1:A:463:LEU:HD11	1.91	0.51
1:B:646:GLN:HG2	1:B:652:LEU:HA	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:32:SER:HB3	1:B:73:ARG:NH1	2.26	0.50
1:A:513:ILE:HD12	1:A:514:PRO:CD	2.42	0.49
1:A:400:GLY:HA3	1:A:478:PHE:HE1	1.77	0.49
1:A:101:LEU:CD1	1:B:628:PRO:HG2	2.42	0.49
1:B:99:ARG:HH12	1:B:134:ARG:HH22	1.59	0.49
1:B:303:GLU:OE1	1:B:378:ASP:OD2	2.30	0.49
1:A:67:ASP:O	1:A:155:MET:HG3	2.12	0.48
1:B:352:PRO:HB2	1:B:353:PRO:CD	2.43	0.48
1:B:433:GLN:OE1	1:B:559:GLY:HA3	2.14	0.48
1:A:295:LEU:HD23	1:A:642:VAL:HG21	1.95	0.48
1:B:549:GLY:HA3	1:B:570:VAL:HG11	1.95	0.48
1:A:63:LEU:HG	1:A:220:LEU:HD21	1.96	0.47
1:A:412:TYR:CE1	1:A:422:ASN:HA	2.50	0.47
1:B:176:ALA:HA	1:B:188:VAL:O	2.15	0.47
1:B:444:ILE:HG23	1:B:503:ALA:HB1	1.97	0.47
1:B:362:THR:N	1:B:363:PRO:HD2	2.30	0.46
1:B:206:LEU:HD23	1:B:206:LEU:N	2.31	0.46
1:B:382:ILE:CG2	1:B:463:LEU:HD11	2.46	0.46
1:A:601:ALA:O	1:A:604:SER:HB2	2.15	0.46
1:A:4:GLN:O	1:A:5:PRO:O	2.33	0.46
1:A:67:ASP:C	1:A:155:MET:HG3	2.41	0.46
1:A:176:ALA:HA	1:A:188:VAL:O	2.17	0.45
1:B:155:MET:O	1:B:159:ARG:HG3	2.16	0.45
1:B:90:THR:HG21	3:B:705:EDO:O2	2.17	0.45
1:B:469:GLU:HB3	1:B:470:PRO:HD2	1.97	0.45
1:A:321:ARG:HA	1:A:330:PRO:HG3	1.99	0.44
1:A:576:ALA:HB2	1:A:640:SER:HB3	1.98	0.44
1:B:533:ARG:O	1:B:533:ARG:HD3	2.18	0.44
1:A:206:LEU:HB3	1:A:212:PHE:HB2	2.00	0.44
1:B:424:GLY:HA3	1:B:446:LEU:CD1	2.47	0.44
1:A:445:ALA:HB3	1:A:504:LEU:HB2	1.99	0.44
1:A:513:ILE:HG12	1:A:557:ALA:HB2	2.00	0.44
1:B:158:ILE:HG23	1:B:163:PHE:CE1	2.53	0.43
1:A:492:VAL:O	1:A:498:PRO:HA	2.17	0.43
1:A:442:GLY:CA	1:A:443:PRO:C	2.91	0.43
1:A:430:ASP:HA	1:A:439:GLN:OE1	2.19	0.43
1:B:28:LEU:HD11	1:B:62:GLY:HA2	2.01	0.43
1:A:596:ASP:OD1	1:A:648:MET:HE2	2.18	0.43
1:A:620:ASP:O	1:B:97:ARG:NH2	2.51	0.43
1:B:178:ARG:HG3	1:B:178:ARG:NH1	2.32	0.43
1:A:4:GLN:C	1:A:5:PRO:O	2.60	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:564:SER:OG	1:A:567:CYS:HB2	2.18	0.42
1:A:382:ILE:CG2	1:A:463:LEU:HD11	2.49	0.42
1:A:190:ALA:HB1	1:A:193:ALA:HB3	2.01	0.42
1:B:334:LEU:HD12	1:B:334:LEU:C	2.45	0.42
1:B:362:THR:N	1:B:363:PRO:CD	2.82	0.42
1:B:588:ILE:HB	1:B:589:PRO:HD3	2.02	0.42
1:A:334:LEU:C	1:A:334:LEU:HD12	2.45	0.41
1:B:402:GLY:HA3	4:B:914:HOH:O	2.20	0.41
1:A:424:GLY:HA3	1:A:446:LEU:CD1	2.51	0.41
1:B:157:GLN:O	1:B:158:ILE:C	2.64	0.41
1:A:400:GLY:O	1:A:408:GLY:HA2	2.20	0.41
1:A:412:TYR:HE2	1:A:446:LEU:HD11	1.85	0.41
1:A:492:VAL:HG12	1:A:493:ASP:N	2.36	0.41
1:B:377:PRO:HG3	1:B:382:ILE:HD11	2.03	0.41
1:B:400:GLY:HA3	1:B:478:PHE:HE2	1.86	0.41
1:B:418:HIS:CD2	1:B:441:GLU:HG2	2.55	0.41
1:B:362:THR:HB	1:B:363:PRO:HD3	2.02	0.40
1:B:555:ARG:HA	1:B:560:PHE:CD1	2.56	0.40
1:A:613:PHE:HA	1:A:627:TYR:H	1.86	0.40
1:B:1:MET:HE3	1:B:1:MET:HB2	1.94	0.40
1:A:555:ARG:HA	1:A:560:PHE:CD1	2.56	0.40
1:B:299:ASP:OD1	1:B:301:GLY:N	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	597/662 (90%)	580 (97%)	15 (2%)	2 (0%)	36	29
1	B	608/662 (92%)	583 (96%)	23 (4%)	2 (0%)	36	29
All	All	1205/1324 (91%)	1163 (96%)	38 (3%)	4 (0%)	36	29

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	5	PRO
1	B	302	MET
1	A	302	MET
1	B	5	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	447/489 (91%)	436 (98%)	11 (2%)	42	37
1	B	453/489 (93%)	440 (97%)	13 (3%)	37	31
All	All	900/978 (92%)	876 (97%)	24 (3%)	39	34

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0	HIS
1	A	1	MET
1	A	3	VAL
1	A	28	LEU
1	A	96	LEU
1	A	145	ASP
1	A	304	THR
1	A	377	PRO
1	A	378	ASP
1	A	381	VAL
1	A	628	PRO
1	B	0	HIS
1	B	3	VAL
1	B	97	ARG
1	B	145	ASP
1	B	180	ARG
1	B	208	ARG
1	B	303	GLU
1	B	377	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	420	LEU
1	B	552	THR
1	B	555	ARG
1	B	648	MET
1	B	652	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	10	GLN
1	B	157	GLN
1	B	269	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	B	705	-	3,3,3	0.22	0	2,2,2	0.34	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	A	703	-	3,3,3	0.23	0	2,2,2	0.13	0
3	EDO	B	702	-	3,3,3	0.20	0	2,2,2	0.16	0
3	EDO	B	703	-	3,3,3	1.18	0	2,2,2	0.25	0
3	EDO	B	704	-	3,3,3	0.32	0	2,2,2	0.68	0
2	BGC	A	701	-	12,12,12	1.78	4 (33%)	17,17,17	1.38	3 (17%)
3	EDO	A	702	-	3,3,3	0.72	0	2,2,2	0.58	0
2	BGC	B	701	-	12,12,12	1.28	2 (16%)	17,17,17	1.10	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	705	-	-	0/1/1/1	-
3	EDO	A	703	-	-	1/1/1/1	-
3	EDO	B	702	-	-	1/1/1/1	-
3	EDO	B	703	-	-	0/1/1/1	-
3	EDO	B	704	-	-	1/1/1/1	-
2	BGC	A	701	-	-	0/2/22/22	0/1/1/1
3	EDO	A	702	-	-	1/1/1/1	-
2	BGC	B	701	-	-	0/2/22/22	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	BGC	C4-C5	3.16	1.59	1.53
2	A	701	BGC	O4-C4	2.88	1.49	1.43
2	A	701	BGC	C4-C3	2.51	1.58	1.52
2	A	701	BGC	O5-C5	2.49	1.50	1.44
2	B	701	BGC	C4-C5	2.41	1.58	1.53
2	B	701	BGC	O4-C4	2.04	1.47	1.43

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	BGC	O3-C3-C2	-2.83	103.80	110.35
2	A	701	BGC	C1-O5-C5	-2.31	109.30	113.66
2	A	701	BGC	O5-C1-C2	-2.26	106.26	110.28
2	A	701	BGC	O2-C2-C3	2.19	115.42	110.35

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	702	EDO	O1-C1-C2-O2
3	B	704	EDO	O1-C1-C2-O2
3	B	702	EDO	O1-C1-C2-O2
3	A	703	EDO	O1-C1-C2-O2


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	705	EDO	2	0
3	B	703	EDO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	605/662 (91%)	0.39	40 (6%) 24 26	19, 30, 64, 98	0
1	B	614/662 (92%)	0.43	39 (6%) 25 27	19, 31, 62, 101	0
All	All	1219/1324 (92%)	0.41	79 (6%) 25 26	19, 31, 63, 101	0

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	420	LEU	7.2
1	A	304	THR	6.4
1	A	3	VAL	5.7
1	B	652	LEU	5.6
1	A	301	GLY	4.8
1	B	301	GLY	4.4
1	B	300	THR	4.4
1	A	300	THR	4.4
1	B	304	THR	4.3
1	A	2	THR	4.2
1	A	105	ILE	4.1
1	A	412	TYR	4.1
1	B	2	THR	4.1
1	A	108	PRO	4.0
1	B	3	VAL	3.9
1	B	302	MET	3.9
1	A	353	PRO	3.8
1	B	353	PRO	3.8
1	B	240	THR	3.8
1	A	352	PRO	3.7
1	A	303	GLU	3.6
1	B	415	ALA	3.5
1	A	106	PRO	3.4
1	A	421	SER	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	176	ALA	3.4
1	A	267	LEU	3.3
1	A	240	THR	3.3
1	A	0	HIS	3.2
1	B	650	VAL	3.2
1	A	338	ARG	3.1
1	A	302	MET	3.1
1	A	104	GLY	3.1
1	A	298	LEU	3.0
1	A	107	GLY	3.0
1	B	242	PRO	3.0
1	B	106	PRO	3.0
1	B	338	ARG	2.9
1	B	651	GLY	2.9
1	A	651	GLY	2.8
1	A	109	THR	2.8
1	A	650	VAL	2.8
1	B	352	PRO	2.8
1	A	5	PRO	2.8
1	B	172	ALA	2.7
1	B	267	LEU	2.7
1	B	298	LEU	2.7
1	A	1	MET	2.7
1	A	32	SER	2.7
1	B	107	GLY	2.6
1	B	170	VAL	2.6
1	B	196	PRO	2.5
1	B	104	GLY	2.5
1	A	269	HIS	2.5
1	B	0	HIS	2.5
1	B	1	MET	2.5
1	A	176	ALA	2.4
1	B	103	ARG	2.4
1	A	324	PRO	2.3
1	A	305	GLY	2.3
1	B	171	SER	2.3
1	B	381	VAL	2.3
1	B	413	ARG	2.3
1	A	242	PRO	2.2
1	B	421	SER	2.2
1	A	429	ARG	2.2
1	A	4	GLN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	5	PRO	2.2
1	A	555	ARG	2.2
1	B	305	GLY	2.2
1	B	555	ARG	2.2
1	B	241	GLY	2.1
1	A	373	TRP	2.1
1	A	325	ALA	2.1
1	A	378	ASP	2.1
1	B	144	ILE	2.1
1	B	268	PRO	2.0
1	A	103	ARG	2.0
1	A	297	PRO	2.0
1	B	210	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BGC	B	701	12/12	0.68	0.21	40,63,74,77	0
2	BGC	A	701	12/12	0.72	0.17	36,49,55,57	0
3	EDO	B	704	4/4	0.72	0.18	41,47,53,53	0
3	EDO	B	703	4/4	0.77	0.15	25,31,32,38	0
3	EDO	B	702	4/4	0.84	0.14	51,53,56,57	0
3	EDO	A	702	4/4	0.85	0.13	27,30,32,33	0
3	EDO	A	703	4/4	0.89	0.13	44,47,50,52	0
3	EDO	B	705	4/4	0.93	0.12	28,42,43,45	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around BGC B 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around BGC A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.