



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 24, 2026 – 02:09 PM JST

PDB ID : 27IQ / pdb_000027iq
Title : Crystal structure of the mature form of human ASPRV1 bound to flaggrin peptide
Authors : Chen, Z.; Feng, X.; Ding, J.
Deposited on : 2026-06-03
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	142	12% 80% 16% ..
1	B	142	% 77% 11% 11%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2110 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Retroviral-like aspartic protease 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	142	Total	C	N	O	S	0	0	0
			1087	696	184	204	3			
1	B	126	Total	C	N	O	S	0	0	0
			968	622	165	178	3			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	190	SER	-	expression tag	UNP Q53RT3
A	212	ASN	ASP	conflict	UNP Q53RT3
A	322	GLY	-	expression tag	UNP Q53RT3
A	323	SER	-	expression tag	UNP Q53RT3
A	324	PHE	-	expression tag	UNP Q53RT3
A	325	LEU	-	expression tag	UNP Q53RT3
A	326	TYR	-	expression tag	UNP Q53RT3
A	327	GLN	-	expression tag	UNP Q53RT3
A	328	VAL	-	expression tag	UNP Q53RT3
A	329	SER	-	expression tag	UNP Q53RT3
A	330	THR	-	expression tag	UNP Q53RT3
A	331	HIS	-	expression tag	UNP Q53RT3
B	190	SER	-	expression tag	UNP Q53RT3
B	212	ASN	ASP	conflict	UNP Q53RT3
B	322	GLY	-	expression tag	UNP Q53RT3
B	323	SER	-	expression tag	UNP Q53RT3
B	324	PHE	-	expression tag	UNP Q53RT3
B	325	LEU	-	expression tag	UNP Q53RT3
B	326	TYR	-	expression tag	UNP Q53RT3
B	327	GLN	-	expression tag	UNP Q53RT3
B	328	VAL	-	expression tag	UNP Q53RT3
B	329	SER	-	expression tag	UNP Q53RT3
B	330	THR	-	expression tag	UNP Q53RT3
B	331	HIS	-	expression tag	UNP Q53RT3

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total 2	Ca 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	27	Total 27	O 27	0	0
3	B	26	Total 26	O 26	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Retroviral-like aspartic protease 1



- Molecule 1: Retroviral-like aspartic protease 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.00Å 64.93Å 76.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	22.74 – 2.09 22.74 – 2.09	Depositor EDS
% Data completeness (in resolution range)	99.0 (22.74-2.09) 98.7 (22.74-2.09)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.09Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, R_{free}	0.262 , 0.292 0.262 , 0.293	Depositor DCC
R_{free} test set	1550 reflections (9.99%)	wwPDB-VP
Wilson B-factor (Å ²)	39.4	Xtriage
Anisotropy	0.190	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 37.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2110	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 99.66 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1202e-13. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/1107	0.75	2/1497 (0.1%)
1	B	0.37	0/985	0.57	0/1332
All	All	0.45	0/2092	0.67	2/2829 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	324	PHE	CA-CB-CG	7.34	121.14	113.80
1	A	321	ASP	N-CA-C	-5.48	105.81	112.88

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	301	ARG	Sidechain
1	B	301	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1087	0	1102	23	0
1	B	968	0	997	14	0
2	A	2	0	0	0	0
3	A	27	0	0	0	0
3	B	26	0	0	0	0
All	All	2110	0	2099	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:325:LEU:HD23	1:B:251:MET:HE1	1.66	0.77
1:A:216:GLN:HB2	1:A:330:THR:HG23	1.67	0.74
1:A:330:THR:O	1:A:331:HIS:C	2.30	0.74
1:A:325:LEU:HD22	1:B:219:VAL:HG21	1.71	0.72
1:A:217:VAL:HG13	1:A:330:THR:HG21	1.72	0.70
1:A:244:LYS:HE3	1:A:331:HIS:CE1	2.29	0.68
1:A:217:VAL:HG13	1:A:330:THR:CG2	2.29	0.63
1:A:266:LEU:HD22	1:A:308:LYS:HG3	1.81	0.62
1:B:266:LEU:HD22	1:B:308:LYS:HG3	1.82	0.61
1:A:325:LEU:HD23	1:B:251:MET:CE	2.36	0.54
1:A:294:ALA:HA	1:A:304:THR:O	2.07	0.52
1:A:244:LYS:HE3	1:A:331:HIS:HE1	1.76	0.50
1:A:243:VAL:HG13	1:A:328:VAL:HG13	1.94	0.48
1:B:294:ALA:HA	1:B:304:THR:O	2.13	0.47
1:A:292:HIS:HB2	1:A:305:LEU:HD12	1.97	0.46
1:A:212:ASN:HD22	1:A:214:GLY:H	1.65	0.44
1:B:212:ASN:HD22	1:B:214:GLY:H	1.65	0.44
1:A:242:VAL:HG13	1:A:250:GLU:HG3	2.00	0.43
1:B:252:LYS:HB2	1:B:252:LYS:HE2	1.81	0.43
1:A:245:VAL:HG23	1:A:247:ASN:OD1	2.18	0.43
1:A:324:PHE:HB2	1:A:326:TYR:CE1	2.52	0.43
1:A:214:GLY:HA3	1:B:212:ASN:ND2	2.34	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:256:VAL:HG12	1:B:273:LEU:HD22	2.01	0.42
1:A:212:ASN:ND2	1:B:214:GLY:HA3	2.34	0.42
1:A:324:PHE:CD1	1:B:216:GLN:HG3	2.54	0.42
1:A:237:GLN:HB2	1:A:256:VAL:HG22	2.00	0.42
1:A:325:LEU:CD2	1:B:219:VAL:HG21	2.46	0.42
1:B:297:ASP:HB3	1:B:302:THR:OG1	2.20	0.42
1:B:212:ASN:ND2	1:B:214:GLY:H	2.19	0.41
1:A:303:CYS:SG	1:A:305:LEU:CD2	3.09	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	140/142 (99%)	134 (96%)	6 (4%)	0	100	100
1	B	124/142 (87%)	121 (98%)	3 (2%)	0	100	100
All	All	264/284 (93%)	255 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/118 (100%)	112 (95%)	6 (5%)	21	21

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	105/118 (89%)	104 (99%)	1 (1%)	68	76
All	All	223/236 (94%)	216 (97%)	7 (3%)	35	39

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	243	VAL
1	A	256	VAL
1	A	265	LYS
1	A	301	ARG
1	A	305	LEU
1	A	324	PHE
1	B	265	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	212	ASN
1	A	327	GLN
1	A	331	HIS
1	B	212	ASN
1	B	241	ASN
1	B	300	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	142/142 (100%)	0.95	17 (11%) 9 9	26, 42, 59, 72	0
1	B	126/142 (88%)	0.32	1 (0%) 82 84	26, 42, 56, 62	0
All	All	268/284 (94%)	0.65	18 (6%) 24 25	26, 42, 58, 72	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	331	HIS	11.0
1	A	316	GLY	8.0
1	A	330	THR	7.2
1	A	324	PHE	6.9
1	A	321	ASP	6.8
1	A	318	SER	6.4
1	A	319	LEU	6.4
1	A	325	LEU	6.1
1	A	326	TYR	5.9
1	A	317	GLY	5.6
1	A	320	GLU	4.8
1	B	299	GLU	4.4
1	A	299	GLU	4.1
1	A	329	SER	4.1
1	A	322	GLY	4.1
1	A	328	VAL	3.6
1	A	327	GLN	2.9
1	A	323	SER	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CA	A	401	1/1	0.92	0.34	30,30,30,30	0
2	CA	A	402	1/1	0.92	0.37	30,30,30,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around CA A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CA A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.