



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2026 – 01:03 pm BST

PDB ID : 29JO / pdb_000029jo
Title : Crystal structure of the unliganded PNGaseABif E236Q mutant
Authors : Hurtado-Guerrero, R.; Taleb, V.
Deposited on : 2026-03-17
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	538	8% 88% 10% ..

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4135 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PNGaseABif E326Q.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	538	Total	C	N	O	S	0	0	0
			4056	2512	692	842	10			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	74	Total	O	0	0
			74	74		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PNGaseABif E326Q



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	85.42Å 77.67Å 80.21Å 90.00° 109.50° 90.00°	Depositor
Resolution (Å)	75.61 – 1.80 75.61 – 1.80	Depositor EDS
% Data completeness (in resolution range)	88.7 (75.61-1.80) 88.7 (75.61-1.80)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.8.0430	Depositor
R, R_{free}	0.201 , 0.240 0.208 , 0.244	Depositor DCC
R_{free} test set	2074 reflections (4.53%)	wwPDB-VP
Wilson B-factor (Å ²)	24.6	Xtriage
Anisotropy	0.468	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 31.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4135	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.67	0/4142	1.12	12/5662 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	543	ASP	CA-CB-CG	6.78	119.38	112.60
1	A	456	ARG	CB-CA-C	-6.57	98.13	110.16
1	A	43	THR	CA-CB-OG1	-6.44	99.94	109.60
1	A	210	ASP	CB-CA-C	6.38	120.25	110.62
1	A	474	ASP	CA-CB-CG	6.16	118.76	112.60
1	A	57	THR	CA-CB-OG1	-6.13	100.40	109.60
1	A	544	ASP	CA-CB-CG	5.97	118.57	112.60
1	A	427	ASP	CA-CB-CG	5.86	118.46	112.60
1	A	210	ASP	CA-CB-CG	5.75	118.36	112.60
1	A	475	PHE	CA-CB-CG	5.73	119.53	113.80
1	A	352	ASP	CA-CB-CG	5.42	118.02	112.60
1	A	81	THR	CA-CB-OG1	-5.26	101.71	109.60

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	218	ARG	Sidechain
1	A	392	ARG	Sidechain
1	A	495	ARG	Sidechain
1	A	50	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4056	0	3868	29	0
2	A	5	0	0	0	0
3	A	74	0	0	0	0
All	All	4135	0	3868	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:328:ARG:HB3	1:A:328:ARG:HH21	1.47	0.80
1:A:125:GLN:HE21	1:A:126:GLU:H	1.43	0.67
1:A:474:ASP:OD2	1:A:476:MET:HE3	1.95	0.66
1:A:548:THR:OG1	1:A:569:SER:HB3	1.98	0.63
1:A:474:ASP:O	1:A:481:TYR:HB2	2.02	0.60
1:A:82:PRO:HG3	1:A:148:PHE:O	2.03	0.59
1:A:474:ASP:HB3	1:A:481:TYR:HA	1.87	0.56
1:A:536:ILE:HG12	1:A:554:GLU:HG2	1.89	0.55
1:A:253:ALA:HB2	1:A:427:ASP:O	2.09	0.52
1:A:423:ILE:HD11	1:A:433:GLN:NE2	2.28	0.49
1:A:253:ALA:HB2	1:A:429:ASN:H	1.79	0.48
1:A:429:ASN:HB3	1:A:474:ASP:HA	1.97	0.46
1:A:250:PHE:O	1:A:251:SER:HB2	2.15	0.45
1:A:257:GLN:O	1:A:258:MET:CB	2.64	0.45
1:A:203:ASN:HB2	1:A:204:PRO:HD2	2.00	0.44
1:A:203:ASN:HB2	1:A:204:PRO:CD	2.47	0.43
1:A:426:SER:O	1:A:427:ASP:CB	2.68	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:297:ARG:HA	1:A:298:PRO:HA	1.85	0.42
1:A:379:GLY:O	1:A:380:ASN:C	2.63	0.42
1:A:425:VAL:O	1:A:426:SER:C	2.62	0.42
1:A:426:SER:O	1:A:427:ASP:HB2	2.20	0.42
1:A:107:ASP:OD1	1:A:107:ASP:N	2.52	0.42
1:A:294:THR:HG23	1:A:521:MET:CE	2.49	0.41
1:A:315:THR:N	1:A:316:PRO:CD	2.83	0.41
1:A:474:ASP:O	1:A:481:TYR:CB	2.68	0.41
1:A:326:LYS:HD2	1:A:328:ARG:HH12	1.86	0.41
1:A:215:MET:HE3	1:A:215:MET:HB3	1.99	0.41
1:A:125:GLN:HE21	1:A:126:GLU:N	2.15	0.40
1:A:253:ALA:CB	1:A:429:ASN:H	2.34	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	536/538 (100%)	508 (95%)	21 (4%)	7 (1%)	9 3

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	251	SER
1	A	258	MET
1	A	427	ASP
1	A	380	ASN
1	A	525	ASN
1	A	477	SER
1	A	291	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	446/446 (100%)	425 (95%)	21 (5%)	23	11

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	54	ARG
1	A	81	THR
1	A	215	MET
1	A	245	ASP
1	A	246	LEU
1	A	254	VAL
1	A	256	THR
1	A	258	MET
1	A	302	LEU
1	A	312	LEU
1	A	318	VAL
1	A	328	ARG
1	A	376	GLN
1	A	423	ILE
1	A	429	ASN
1	A	435	MET
1	A	450	ASP
1	A	474	ASP
1	A	501	THR
1	A	505	LEU
1	A	525	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	63	HIS
1	A	125	GLN
1	A	130	GLN
1	A	222	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	232	ASN
1	A	341	ASN
1	A	360	ASN
1	A	433	GLN
1	A	437	GLN
1	A	459	ASN
1	A	479	ASN
1	A	551	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	601	-	4,4,4	0.46	0	6,6,6	0.11	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	538/538 (100%)	0.50	41 (7%)	20 18	17, 26, 70, 163	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	254	VAL	9.7
1	A	480	GLY	7.5
1	A	253	ALA	6.4
1	A	475	PHE	6.4
1	A	501	THR	5.3
1	A	473	TYR	5.2
1	A	255	SER	5.2
1	A	503	ALA	4.9
1	A	477	SER	4.9
1	A	502	THR	4.9
1	A	479	ASN	4.7
1	A	500	GLY	4.6
1	A	478	ASP	4.4
1	A	476	MET	4.2
1	A	379	GLY	4.1
1	A	425	VAL	4.0
1	A	252	ASP	3.9
1	A	427	ASP	3.9
1	A	426	SER	3.7
1	A	380	ASN	3.7
1	A	524	ALA	3.6
1	A	381	GLY	3.6
1	A	256	THR	3.5
1	A	474	ASP	3.2
1	A	32	ASN	3.0
1	A	382	ASN	3.0
1	A	429	ASN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	528	LEU	2.7
1	A	527	ALA	2.7
1	A	505	LEU	2.7
1	A	526	GLY	2.6
1	A	569	SER	2.6
1	A	164	ASP	2.6
1	A	258	MET	2.5
1	A	525	ASN	2.5
1	A	428	GLY	2.4
1	A	251	SER	2.4
1	A	378	THR	2.3
1	A	257	GLN	2.2
1	A	34	ILE	2.1
1	A	54	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	A	601	5/5	0.89	0.10	54,54,60,64	0

6.5 Other polymers [i](#)

There are no such residues in this entry.