



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2026 – 01:07 pm BST

PDB ID : 29MJ / pdb_000029mj
Title : Akt1 mutant in complex with miransertib (ARQ 092)
Authors : Depta, L.; Niggenaber, J.; Mueller, M.P.; Rauh, D.
Deposited on : 2026-03-23
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	445	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 69%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 21%; height: 10px; background-color: grey;"></div> </div>
1	B	445	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 67%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 23%; height: 10px; background-color: grey;"></div> </div>
1	C	445	<div style="display: flex; align-items: center;"> <div style="width: 10%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 66%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 24%; height: 10px; background-color: grey;"></div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8388 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RAC-alpha serine/threonine-protein kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	353	2789	1803	464	509	13	0	2	0
1	B	341	2758	1783	461	502	12	0	2	0
1	C	336	2601	1682	434	472	13	0	1	0

There are 27 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP P31749
A	114	ALA	GLU	engineered mutation	UNP P31749
A	115	ALA	GLU	engineered mutation	UNP P31749
A	116	ALA	GLU	engineered mutation	UNP P31749
A	205	THR	SER	engineered mutation	UNP P31749
A	262	GLU	ASP	engineered mutation	UNP P31749
A	?	-	GLU	deletion	UNP P31749
A	268	ARG	LYS	engineered mutation	UNP P31749
A	269	ASP	ASN	engineered mutation	UNP P31749
B	1	GLY	-	expression tag	UNP P31749
B	114	ALA	GLU	engineered mutation	UNP P31749
B	115	ALA	GLU	engineered mutation	UNP P31749
B	116	ALA	GLU	engineered mutation	UNP P31749
B	205	THR	SER	engineered mutation	UNP P31749
B	262	GLU	ASP	engineered mutation	UNP P31749
B	?	-	GLU	deletion	UNP P31749
B	268	ARG	LYS	engineered mutation	UNP P31749
B	269	ASP	ASN	engineered mutation	UNP P31749
C	1	GLY	-	expression tag	UNP P31749
C	114	ALA	GLU	engineered mutation	UNP P31749
C	115	ALA	GLU	engineered mutation	UNP P31749
C	116	ALA	GLU	engineered mutation	UNP P31749
C	205	THR	SER	engineered mutation	UNP P31749

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	262	GLU	ASP	engineered mutation	UNP P31749
C	?	-	GLU	deletion	UNP P31749
C	268	ARG	LYS	engineered mutation	UNP P31749
C	269	ASP	ASN	engineered mutation	UNP P31749

- Molecule 2 is 3-[3-[4-(1-azanylcyclobutyl)phenyl]-5-phenyl-imidazo[4,5-b]pyridin-2-yl]pyridin-2-amine (CCD ID: 6S1) (formula: C₂₇H₂₄N₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
2	A	1	Total	C	N	0	0
			33	27	6		
2	B	1	Total	C	N	0	0
			33	27	6		
2	C	1	Total	C	N	0	0
			33	27	6		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	50	Total 50	O 50	0	0
5	B	38	Total 38	O 38	0	0
5	C	37	Total 37	O 37	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RAC-alpha serine/threonine-protein kinase



- Molecule 1: RAC-alpha serine/threonine-protein kinase



- Molecule 1: RAC-alpha serine/threonine-protein kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	44.80Å 164.25Å 188.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.37 – 2.50 47.37 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.37-2.50) 100.0 (47.37-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.64 (at 2.51Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.228 , 0.264 0.228 , 0.264	Depositor DCC
R_{free} test set	2474 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	62.4	Xtrriage
Anisotropy	0.337	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 63.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8388	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.45 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.9969e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, 6S1, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.12	0/2865	0.30	0/3884
1	B	0.11	0/2831	0.30	0/3829
1	C	0.13	0/2668	0.33	0/3622
All	All	0.12	0/8364	0.31	0/11335

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2789	0	2613	27	0
1	B	2758	0	2639	24	0
1	C	2601	0	2406	30	0
2	A	33	0	0	0	0
2	B	33	0	0	0	0
2	C	33	0	0	0	0
3	A	4	0	6	0	0
4	A	6	0	8	1	0
4	B	6	0	8	0	0
5	A	50	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	38	0	0	1	0
5	C	37	0	0	3	0
All	All	8388	0	7680	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:156:LEU:HD21	1:C:166:LEU:HB2	1.74	0.70
1:C:15[B]:ARG:HH11	1:C:20:LYS:HB3	1.60	0.66
1:A:208:PRO:O	1:A:289:LYS:NZ	2.24	0.63
1:C:273:ARG:NH1	5:C:602:HOH:O	2.32	0.62
1:A:150:PHE:O	1:B:428:GLN:NE2	2.33	0.62
1:A:419:LYS:NZ	5:A:601:HOH:O	2.27	0.61
1:B:242:GLU:OE1	1:B:249:ARG:NH1	2.32	0.61
1:B:375:GLU:HG3	1:B:406:ARG:HD2	1.83	0.60
1:A:147:MET:HB3	1:B:436:ARG:HH21	1.66	0.60
1:C:149:GLU:OE1	1:C:149:GLU:N	2.35	0.59
1:C:324:ASN:OD1	5:C:601:HOH:O	2.17	0.59
1:B:339:MET:HA	1:B:342:MET:HE2	1.84	0.58
1:B:174:ARG:NH2	5:B:601:HOH:O	2.38	0.57
1:C:69:ARG:NH1	5:C:604:HOH:O	2.36	0.57
1:C:242:GLU:OE1	1:C:249:ARG:NH1	2.38	0.56
1:A:339:MET:HA	1:A:342:MET:HE2	1.86	0.55
4:A:503:GOL:H11	1:C:163:LYS:HZ3	1.71	0.55
1:A:36[B]:ILE:HD11	1:A:325:ASP:OD1	2.06	0.55
1:B:6:ILE:HD13	1:B:30:LYS:HG2	1.88	0.54
1:B:274:ASP:O	1:B:279:ASN:ND2	2.41	0.54
1:A:408:PHE:HD1	1:A:411:ILE:HD12	1.73	0.54
1:A:149:GLU:N	1:A:149:GLU:OE1	2.42	0.53
1:C:76:ARG:NH2	1:C:85:GLU:OE2	2.41	0.53
1:C:281:MET:HE2	1:C:291:THR:HG22	1.90	0.53
1:B:76:ARG:NH2	1:B:85:GLU:OE2	2.40	0.53
1:A:273:ARG:NH1	5:A:609:HOH:O	2.40	0.52
1:A:164:VAL:HG22	1:A:179:LYS:HG3	1.90	0.52
1:C:339:MET:HA	1:C:342:MET:HE2	1.92	0.51
1:A:251:ARG:HG2	1:A:411:ILE:HD11	1.93	0.51
1:A:62:LEU:HD12	1:A:104:GLN:HG2	1.94	0.50
1:C:234:GLU:HA	1:C:281:MET:HG2	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:230:ALA:HB1	1:C:282:LEU:O	2.12	0.49
1:A:6:ILE:HD13	1:A:30:LYS:HG2	1.93	0.49
1:C:15[B]:ARG:NH1	1:C:20:LYS:HB3	2.27	0.48
1:A:428:GLN:OE1	1:A:436:ARG:NH2	2.39	0.48
1:B:36:ILE:HD11	1:B:325:ASP:OD1	2.14	0.47
1:C:340:TYR:CD1	1:C:348:PRO:HG3	2.49	0.47
1:B:213:LEU:HD11	1:B:225:PHE:HB3	1.97	0.46
1:C:242:GLU:O	1:C:244:VAL:N	2.48	0.46
1:A:242:GLU:O	1:A:244:VAL:N	2.47	0.45
1:B:340:TYR:CE2	1:B:348:PRO:HA	2.51	0.45
1:A:413:TRP:O	1:A:416:VAL:HG22	2.17	0.45
1:B:179:LYS:O	1:B:224:CYS:HA	2.17	0.45
1:A:274:ASP:O	1:A:279:ASN:ND2	2.50	0.45
1:B:416:VAL:HG22	1:B:421:LEU:HD11	1.99	0.45
1:C:348:PRO:HB2	1:C:349:PHE:CD2	2.51	0.45
1:C:179:LYS:HD2	1:C:227:MET:CE	2.46	0.44
1:A:213:LEU:HD11	1:A:225:PHE:HB3	1.99	0.44
1:B:210:LEU:O	1:B:289:LYS:NZ	2.51	0.44
1:B:44:ASP:OD1	1:B:45:VAL:N	2.48	0.44
1:C:156:LEU:CD2	1:C:166:LEU:HB2	2.45	0.44
1:A:234:GLU:HB2	1:A:237:PHE:HD2	1.83	0.44
1:B:145:VAL:N	1:B:149:GLU:OE2	2.51	0.44
1:B:250:ALA:HB1	1:B:342:MET:HE3	1.99	0.43
1:C:6:ILE:HD13	1:C:30:LYS:HG2	2.01	0.43
1:A:250:ALA:HB1	1:A:342:MET:HE3	2.01	0.43
1:A:80:TRP:CD1	1:A:81:THR:HG23	2.52	0.43
1:A:231:ASN:ND2	1:A:431:SER:O	2.52	0.43
1:B:80:TRP:CD1	1:B:81:THR:HG23	2.54	0.42
1:C:179:LYS:HD2	1:C:227:MET:HE3	2.02	0.42
1:C:207:HIS:CG	1:C:208:PRO:HD2	2.54	0.42
1:C:207:HIS:HB2	1:C:263:TYR:CD2	2.54	0.42
1:A:44:ASP:OD1	1:A:45:VAL:N	2.50	0.42
1:B:14:LYS:HD2	1:B:323:ASP:OD2	2.19	0.42
1:B:349:PHE:CE2	1:B:366:ILE:HG12	2.54	0.42
1:A:253:TYR:HB3	1:A:288:ILE:HD11	2.02	0.41
1:C:80:TRP:CD1	1:C:81:THR:HG23	2.55	0.41
1:C:392:LEU:HG	1:C:402:ILE:HG12	2.01	0.41
1:A:261:LEU:HD11	1:A:275:LEU:HD22	2.01	0.41
1:A:207:HIS:CG	1:A:208:PRO:HD2	2.55	0.41
1:C:230:ALA:HB3	1:C:281:MET:HE3	2.02	0.41
1:B:152:TYR:HD1	1:B:165:ILE:HG21	1.85	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:15:ARG:HD3	1:B:22:TRP:CZ2	2.56	0.41
1:B:242:GLU:CD	1:B:249:ARG:HH12	2.29	0.41
1:C:238:HIS:HD2	1:C:282:LEU:HD12	1.85	0.41
1:C:418:GLU:HB2	1:C:420:LYS:HG2	2.03	0.41
1:A:430:THR:HG22	1:C:152:TYR:CE1	2.56	0.40
1:C:322:GLU:HG3	1:C:324:ASN:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/445 (78%)	330 (96%)	15 (4%)	0	100	100
1	B	329/445 (74%)	317 (96%)	12 (4%)	0	100	100
1	C	323/445 (73%)	313 (97%)	10 (3%)	0	100	100
All	All	997/1335 (75%)	960 (96%)	37 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/389 (72%)	276 (99%)	2 (1%)	76	89

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	285/389 (73%)	284 (100%)	1 (0%)	84	93
1	C	254/389 (65%)	253 (100%)	1 (0%)	84	93
All	All	817/1167 (70%)	813 (100%)	4 (0%)	81	92

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	89	HIS
1	A	416	VAL
1	B	89	HIS
1	C	89	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	104	GLN
1	C	61	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	6S1	B	501	-	38,38,38	2.11	11 (28%)	46,56,56	1.84	7 (15%)
4	GOL	A	503	-	5,5,5	0.91	0	5,5,5	1.01	0
2	6S1	A	501	-	38,38,38	2.11	11 (28%)	46,56,56	1.87	6 (13%)
3	EDO	A	502	-	3,3,3	0.46	0	2,2,2	0.32	0
2	6S1	C	501	-	38,38,38	2.12	11 (28%)	46,56,56	1.87	6 (13%)
4	GOL	B	502	-	5,5,5	0.91	0	5,5,5	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	6S1	B	501	-	-	0/16/26/26	0/6/6/6
4	GOL	A	503	-	-	0/4/4/4	-
2	6S1	A	501	-	-	0/16/26/26	0/6/6/6
3	EDO	A	502	-	-	0/1/1/1	-
2	6S1	C	501	-	-	0/16/26/26	0/6/6/6
4	GOL	B	502	-	-	0/4/4/4	-

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	501	6S1	C17-C18	-5.97	1.30	1.40
2	B	501	6S1	C17-C18	-5.92	1.30	1.40
2	A	501	6S1	C17-C18	-5.91	1.30	1.40
2	B	501	6S1	C12-C13	-5.33	1.38	1.42
2	C	501	6S1	C12-C13	-5.29	1.38	1.42
2	A	501	6S1	C12-C13	-5.26	1.38	1.42
2	C	501	6S1	C13-N3	4.07	1.44	1.34
2	B	501	6S1	C13-N3	4.05	1.44	1.34
2	A	501	6S1	C13-N3	4.05	1.44	1.34
2	A	501	6S1	C4-C1	-3.87	1.51	1.55
2	C	501	6S1	C4-C1	-3.87	1.51	1.55
2	B	501	6S1	C4-C1	-3.72	1.51	1.55
2	C	501	6S1	C2-C1	-3.33	1.51	1.55
2	C	501	6S1	C12-C11	3.33	1.53	1.47
2	A	501	6S1	C12-C11	3.26	1.53	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	6S1	C2-C1	-3.25	1.52	1.55
2	C	501	6S1	C11-N2	-3.20	1.35	1.38
2	B	501	6S1	C11-N2	-3.19	1.35	1.38
2	B	501	6S1	C2-C1	-3.14	1.52	1.55
2	A	501	6S1	C11-N2	-3.12	1.35	1.38
2	B	501	6S1	C12-C11	3.12	1.52	1.47
2	C	501	6S1	C1-C5	-2.93	1.47	1.51
2	A	501	6S1	C1-C5	-2.86	1.47	1.51
2	B	501	6S1	C22-C21	2.83	1.53	1.48
2	B	501	6S1	C1-C5	-2.73	1.47	1.51
2	C	501	6S1	C11-N5	2.71	1.36	1.32
2	B	501	6S1	C11-N5	2.68	1.36	1.32
2	A	501	6S1	C11-N5	2.68	1.36	1.32
2	C	501	6S1	C22-C21	2.60	1.53	1.48
2	A	501	6S1	C22-C21	2.55	1.53	1.48
2	C	501	6S1	C19-C17	2.27	1.43	1.39
2	A	501	6S1	C19-C17	2.25	1.43	1.39
2	B	501	6S1	C19-C17	2.23	1.43	1.39

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	6S1	C19-C17-C18	6.95	123.56	117.91
2	C	501	6S1	C19-C17-C18	6.88	123.50	117.91
2	B	501	6S1	C19-C17-C18	6.67	123.33	117.91
2	C	501	6S1	C17-C18-N2	5.54	109.11	105.48
2	A	501	6S1	C17-C18-N2	5.51	109.10	105.48
2	B	501	6S1	C17-C18-N2	5.44	109.05	105.48
2	C	501	6S1	C20-C19-C17	-4.75	112.00	120.02
2	A	501	6S1	C20-C19-C17	-4.74	112.01	120.02
2	B	501	6S1	C20-C19-C17	-4.66	112.15	120.02
2	B	501	6S1	C12-C13-N3	-4.15	119.49	122.52
2	C	501	6S1	C12-C13-N3	-4.01	119.59	122.52
2	A	501	6S1	C12-C13-N3	-3.94	119.64	122.52
2	C	501	6S1	N2-C11-N5	-3.16	109.89	112.39
2	A	501	6S1	N2-C11-N5	-3.05	109.98	112.39
2	B	501	6S1	N2-C11-N5	-2.78	110.19	112.39
2	C	501	6S1	C14-N4-C13	2.38	121.10	117.61
2	B	501	6S1	C16-C12-C13	2.38	119.20	117.44
2	A	501	6S1	C14-N4-C13	2.32	121.01	117.61
2	B	501	6S1	C14-N4-C13	2.13	120.73	117.61

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	503	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	353/445 (79%)	0.59	18 (5%) 33 29	28, 75, 113, 136	2 (0%)
1	B	341/445 (76%)	0.43	18 (5%) 32 28	36, 65, 116, 138	2 (0%)
1	C	336/445 (75%)	0.77	43 (12%) 7 6	31, 82, 127, 148	1 (0%)
All	All	1030/1335 (77%)	0.60	79 (7%) 19 17	28, 73, 119, 148	5 (0%)

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	113	GLN	4.7
1	A	365	GLU	4.5
1	B	438	PHE	4.4
1	C	68	PRO	4.2
1	A	366	ILE	4.0
1	C	288	ILE	3.9
1	A	438	PHE	3.8
1	C	175	TYR	3.8
1	A	68	PRO	3.7
1	B	68	PRO	3.7
1	C	313	PRO	3.6
1	C	152	TYR	3.5
1	C	67	ARG	3.5
1	A	368	PHE	3.5
1	C	367	ARG	3.5
1	C	204	ASN	3.4
1	B	161	PHE	3.4
1	C	372	LEU	3.4
1	C	370	ARG	3.3
1	A	182	LYS	3.3
1	C	369	PRO	3.3
1	C	3	ASP	3.1
1	C	112	LYS	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	350	TYR	3.1
1	B	218	GLN	3.1
1	C	274	ASP	3.0
1	C	232	GLY	2.9
1	B	4	VAL	2.9
1	B	67	ARG	2.8
1	A	371	THR	2.8
1	A	372	LEU	2.8
1	C	293	PHE	2.8
1	B	204	ASN	2.7
1	A	184	GLU	2.7
1	C	221	ASP	2.7
1	B	47	GLN	2.7
1	C	277	LEU	2.6
1	C	223	LEU	2.6
1	B	112	LYS	2.6
1	A	204	ASN	2.6
1	A	144	ARG	2.5
1	C	428	GLN	2.5
1	C	206	ARG	2.5
1	B	3	ASP	2.5
1	C	185	VAL	2.5
1	A	4	VAL	2.4
1	C	224	CYS	2.4
1	C	381	SER	2.4
1	C	321	LEU	2.4
1	C	4	VAL	2.4
1	A	67	ARG	2.3
1	C	157	GLY	2.3
1	A	205	THR	2.3
1	C	174	ARG	2.3
1	C	219	THR	2.2
1	C	368	PHE	2.2
1	A	369	PRO	2.2
1	C	145	VAL	2.2
1	C	47	GLN	2.2
1	C	371	THR	2.2
1	C	231	ASN	2.2
1	C	421	LEU	2.2
1	C	290	ILE	2.1
1	C	220	HIS	2.1
1	C	287	HIS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	161	PHE	2.1
1	C	349	PHE	2.1
1	C	186	ILE	2.1
1	B	147	MET	2.1
1	C	425	PHE	2.1
1	B	156	LEU	2.1
1	B	205	THR	2.1
1	B	315	TYR	2.1
1	C	396	SER	2.1
1	B	167	VAL	2.0
1	B	370[A]	ARG	2.0
1	B	151	GLU	2.0
1	A	219	THR	2.0
1	B	152	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	B	502	6/6	0.71	0.15	81,83,91,94	0
3	EDO	A	502	4/4	0.78	0.18	64,68,71,76	0
4	GOL	A	503	6/6	0.79	0.13	85,88,92,92	0
2	6S1	A	501	33/33	0.95	0.09	47,51,56,60	0
2	6S1	B	501	33/33	0.95	0.08	52,59,62,68	0
2	6S1	C	501	33/33	0.95	0.09	51,57,65,68	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers


as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around 6S1 B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.