



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 1, 2026 – 10:32 am BST

PDB ID : 29US / pdb_000029us
Title : KAT6A SURFACE MUTANT IN COMPLEX WITH QUINOLINE INHIBITOR COMPOUND-1
Authors : Hillig, R.C.; Puetter, V.
Deposited on : 2026-04-09
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	6% 79% 19% ..
1	B	272	9% 75% 22% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4625 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone acetyltransferase KAT6A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	270	Total	C	N	O	S	0	1	0
			2265	1470	381	401	13			
1	B	270	Total	C	N	O	S	0	1	0
			2263	1467	383	400	13			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	507	GLY	-	expression tag	UNP Q92794
A	508	SER	-	expression tag	UNP Q92794
A	638	SER	CYS	engineered mutation	UNP Q92794
A	646	SER	CYS	engineered mutation	UNP Q92794
A	723	SER	CYS	engineered mutation	UNP Q92794
A	773	SER	CYS	engineered mutation	UNP Q92794
B	507	GLY	-	expression tag	UNP Q92794
B	508	SER	-	expression tag	UNP Q92794
B	638	SER	CYS	engineered mutation	UNP Q92794
B	646	SER	CYS	engineered mutation	UNP Q92794
B	723	SER	CYS	engineered mutation	UNP Q92794
B	773	SER	CYS	engineered mutation	UNP Q92794

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 6-(dimethylamino)-N-(2-methylquinolin-8-yl)sulfonyl-1-benzofuran-2-carboxamide (CCD ID: A1J4W) (formula: C₂₁H₁₉N₃O₄S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			29	21	3	4	1		
3	B	1	Total	C	N	O	S	0	0
			29	21	3	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	18	Total	O	0	0
			18	18		
4	B	19	Total	O	0	0
			19	19		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Histone acetyltransferase KAT6A



• Molecule 1: Histone acetyltransferase KAT6A



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	37.03Å 58.91Å 64.92Å 93.02° 91.25° 102.89°	Depositor
Resolution (Å)	34.19 – 2.70 34.19 – 2.70	Depositor EDS
% Data completeness (in resolution range)	91.4 (34.19-2.70) 91.4 (34.19-2.70)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.96 (at 1.82Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.235 , 0.309 0.238 , 0.310	Depositor DCC
R_{free} test set	1093 reflections (3.22%)	wwPDB-VP
Wilson B-factor (Å ²)	38.7	Xtriage
Anisotropy	0.162	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 72.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4625	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ALY, ZN, A1J4W

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.12	3/2318 (0.1%)	1.14	3/3129 (0.1%)
1	B	1.10	4/2316 (0.2%)	1.13	5/3126 (0.2%)
All	All	1.11	7/4634 (0.2%)	1.14	8/6255 (0.1%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	569	GLU	C-O	-9.23	1.13	1.24
1	A	580	GLU	C-O	-6.54	1.15	1.23
1	A	586	SER	CA-C	6.11	1.58	1.53
1	B	586	SER	CA-C	5.40	1.58	1.53
1	B	632	PHE	N-CA	-5.22	1.39	1.46
1	B	596	LEU	CA-C	5.20	1.60	1.52
1	B	580	GLU	C-O	-5.17	1.17	1.23

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	625	GLY	N-CA-C	7.45	119.82	110.96
1	B	574	ASN	CB-CA-C	6.48	123.32	110.42
1	A	765	ARG	CB-CA-C	5.43	115.87	110.33
1	B	539	LEU	N-CA-C	5.19	117.42	109.07
1	A	519	ILE	CB-CA-C	-5.16	102.80	110.33
1	B	584	ASN	N-CA-C	-5.11	106.90	113.23
1	B	611	GLU	CA-C-N	-5.05	113.52	119.84
1	B	611	GLU	C-N-CA	-5.05	113.52	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2265	0	2265	21	0
1	B	2263	0	2263	28	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	29	0	0	1	0
3	B	29	0	0	3	0
4	A	18	0	0	0	0
4	B	19	0	0	1	0
All	All	4625	0	4528	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:660:ARG:HB2	3:B:801:A1J4W:C20	2.05	0.86
1:B:549:SER:HB2	1:B:552:ILE:HG12	1.73	0.71
1:A:706:GLN:HB3	1:A:709:LYS:HG2	1.76	0.67
1:A:683:LEU:O	1:A:688:ARG:NH2	2.28	0.66
1:B:716:LEU:O	1:B:720:THR:HG22	2.00	0.61
1:A:716:LEU:O	1:A:720:THR:HG22	2.01	0.61
1:B:655:ARG:C	3:B:801:A1J4W:C29	2.77	0.58
1:B:683:LEU:HB2	1:B:688:ARG:HD2	1.86	0.56
1:B:550:ARG:HD3	1:B:554:GLN:NE2	2.22	0.54
1:A:514:PHE:CE1	1:A:541:GLU:HA	2.43	0.54
1:B:550:ARG:HD3	1:B:554:GLN:HE22	1.73	0.53
1:B:549:SER:CB	1:B:552:ILE:HG12	2.38	0.53
1:A:685:ASP:OD1	1:A:685:ASP:N	2.38	0.52
1:B:668:LEU:CD2	1:B:774:LEU:HD22	2.40	0.52
1:A:650:LEU:HD13	1:A:652:GLN:HE22	1.74	0.52
1:B:650:LEU:HD13	1:B:652:GLN:HE22	1.75	0.52
1:B:569:GLU:OE1	1:B:572:ARG:NE	2.30	0.52
1:B:667:TYR:CE2	1:B:722:ILE:HG12	2.45	0.51
1:B:655:ARG:O	3:B:801:A1J4W:C29	2.59	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:668:LEU:CD2	1:A:774:LEU:HD22	2.41	0.50
1:B:511:VAL:HG13	1:B:536:LYS:HG3	1.93	0.50
1:B:514:PHE:CE1	1:B:541:GLU:HA	2.47	0.50
1:B:553:LEU:O	1:B:557:MET:HG2	2.13	0.49
1:A:589:TYR:C	1:A:589:TYR:CD1	2.91	0.49
1:A:553:LEU:O	1:A:557:MET:HG2	2.13	0.48
1:B:667:TYR:HE2	1:B:722:ILE:HA	1.77	0.48
1:A:667:TYR:CE1	1:A:722:ILE:HA	2.50	0.47
1:A:667:TYR:HE1	1:A:722:ILE:HA	1.80	0.47
1:B:703:LEU:HD22	1:B:748:ARG:HH11	1.79	0.46
1:A:704:TYR:CD2	1:A:757:MET:HG2	2.51	0.46
1:B:589:TYR:CD1	1:B:589:TYR:C	2.94	0.46
1:B:725:GLN:HG3	1:B:726:ASP:N	2.30	0.44
1:B:700:LEU:HD21	1:B:736:MET:HE1	2.00	0.44
1:A:704:TYR:CE2	1:A:757:MET:HG2	2.53	0.43
1:B:660:ARG:O	1:B:661:PHE:C	2.62	0.43
1:A:660:ARG:O	1:A:661:PHE:C	2.61	0.42
1:B:628:LEU:C	1:B:628:LEU:HD23	2.45	0.42
1:A:613:PHE:CE2	1:A:635:GLU:HG2	2.55	0.41
1:A:539:LEU:HD23	1:A:596:LEU:HD13	2.02	0.41
1:A:632:PHE:CD2	1:A:662:LEU:HD22	2.54	0.41
1:B:578:VAL:HG11	1:B:665:PHE:CZ	2.54	0.41
1:B:648:MET:SD	1:B:648:MET:C	3.04	0.41
1:A:659:GLY:HA3	3:A:801:A1J4W:N14	2.35	0.41
1:B:513:GLU:OE1	1:B:550:ARG:NH2	2.54	0.41
1:A:513:GLU:HG3	1:A:538:TYR:CD2	2.56	0.41
1:B:567:ALA:HB1	1:B:580:GLU:O	2.21	0.41
1:A:578:VAL:HG11	1:A:665:PHE:CZ	2.56	0.40
1:A:642:TYR:HA	1:A:678:SER:O	2.21	0.40
1:B:735:ARG:NH2	4:B:905:HOH:O	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	268/272 (98%)	249 (93%)	17 (6%)	2 (1%)	18	41
1	B	268/272 (98%)	249 (93%)	17 (6%)	2 (1%)	18	41
All	All	536/544 (98%)	498 (93%)	34 (6%)	4 (1%)	18	41

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	735	ARG
1	B	705	HIS
1	A	705	HIS
1	B	735	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/254 (100%)	237 (93%)	17 (7%)	15	36
1	B	254/254 (100%)	234 (92%)	20 (8%)	11	28
All	All	508/508 (100%)	471 (93%)	37 (7%)	13	32

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	510	SER
1	A	511	VAL
1	A	532	SER
1	A	589	TYR
1	A	620	GLN
1	A	624	LYS
1	A	685	ASP
1	A	686	LEU
1	A	709	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	710	GLN
1	A	711	ILE
1	A	749	ARG
1	A	761	GLN
1	A	763	ASN
1	A	765	ARG
1	A	767	VAL
1	A	777	THR
1	B	516	LYS
1	B	532	SER
1	B	550	ARG
1	B	620	GLN
1	B	681	LYS
1	B	686	LEU
1	B	688	ARG
1	B	689	LEU
1	B	690	SER
1	B	692	MET
1	B	710	GLN
1	B	725	GLN
1	B	738	ASP
1	B	742	ASP
1	B	749	ARG
1	B	754	GLN
1	B	762	LEU
1	B	765	ARG
1	B	767	VAL
1	B	777	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	555	GLN
1	A	564	HIS
1	A	568	ASN
1	A	637	HIS
1	A	640	GLN
1	A	675	GLN
1	A	706	GLN
1	A	754	GLN
1	B	554	GLN
1	B	568	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	591	GLN
1	B	603	HIS
1	B	620	GLN
1	B	637	HIS
1	B	640	GLN
1	B	725	GLN
1	B	743	GLN
1	B	754	GLN
1	B	761	GLN
1	B	763	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	ALY	A	604	1	10,11,12	0.91	0	7,12,14	1.02	0
1	ALY	B	604	1	10,11,12	1.27	1 (10%)	7,12,14	1.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	ALY	A	604	1	-	2/9/10/12	-
1	ALY	B	604	1	-	1/9/10/12	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	604	ALY	CB-CA	3.05	1.57	1.53

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	604	ALY	CE-CD-CG-CB
1	A	604	ALY	CE-CD-CG-CB
1	A	604	ALY	CG-CD-CE-NZ

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	A1J4W	B	801	-	32,32,32	2.99	8 (25%)	46,48,48	2.80	10 (21%)
3	A1J4W	A	801	-	32,32,32	3.32	10 (31%)	46,48,48	2.44	13 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1J4W	B	801	-	-	2/19/19/19	0/4/4/4
3	A1J4W	A	801	-	-	4/19/19/19	0/4/4/4

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	801	A1J4W	C18-S15	-11.86	1.65	1.77
3	A	801	A1J4W	C18-S15	-11.00	1.66	1.77
3	A	801	A1J4W	C18-C23	-7.97	1.37	1.42
3	A	801	A1J4W	C13-N14	-6.52	1.32	1.38
3	B	801	A1J4W	O16-S15	6.18	1.50	1.43
3	A	801	A1J4W	O16-S15	5.98	1.50	1.43
3	B	801	A1J4W	C13-N14	-5.52	1.33	1.38
3	A	801	A1J4W	O17-S15	5.42	1.49	1.43
3	B	801	A1J4W	O17-S15	5.39	1.49	1.43
3	A	801	A1J4W	C2-C3	-3.60	1.32	1.44
3	B	801	A1J4W	C18-C23	-3.40	1.40	1.42
3	A	801	A1J4W	C1-C2	-2.80	1.36	1.41
3	B	801	A1J4W	C2-C3	-2.64	1.35	1.44
3	B	801	A1J4W	O5-C6	-2.59	1.34	1.38
3	A	801	A1J4W	C8-N9	-2.46	1.31	1.37
3	B	801	A1J4W	C8-N9	-2.40	1.31	1.37
3	A	801	A1J4W	O5-C4	-2.27	1.34	1.38
3	A	801	A1J4W	C27-C26	2.08	1.41	1.36

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	801	A1J4W	O17-S15-O16	-10.08	107.16	119.55
3	B	801	A1J4W	C13-N14-S15	-9.93	110.65	123.78
3	B	801	A1J4W	O17-S15-O16	-8.79	108.75	119.55
3	B	801	A1J4W	C18-C23-N24	8.02	122.58	119.55
3	A	801	A1J4W	C13-N14-S15	-5.15	116.98	123.78
3	B	801	A1J4W	C18-S15-N14	4.59	115.11	107.29
3	A	801	A1J4W	C18-C23-C22	4.36	120.11	117.68
3	A	801	A1J4W	C19-C18-S15	4.07	121.81	116.98
3	B	801	A1J4W	C6-C2-C3	3.76	109.71	105.65
3	A	801	A1J4W	C12-C1-C2	-3.61	116.85	121.74
3	A	801	A1J4W	C23-C18-S15	-3.58	116.95	120.76
3	B	801	A1J4W	C4-C13-N14	3.16	120.21	115.49
3	A	801	A1J4W	O17-S15-C18	3.05	113.46	108.08
3	B	801	A1J4W	C29-C25-N24	3.02	122.17	117.23
3	B	801	A1J4W	C22-C23-N24	-2.71	119.04	122.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	801	A1J4W	O28-C13-C4	-2.68	117.04	121.07
3	A	801	A1J4W	O5-C6-C7	2.65	130.63	124.97
3	A	801	A1J4W	C1-C2-C6	2.61	122.03	118.57
3	B	801	A1J4W	C29-C25-C26	-2.56	116.39	121.63
3	A	801	A1J4W	O17-S15-N14	2.53	113.86	106.74
3	A	801	A1J4W	C6-C2-C3	2.15	107.97	105.65
3	A	801	A1J4W	O28-C13-N14	2.05	124.59	121.67
3	A	801	A1J4W	C6-O5-C4	-2.04	103.31	105.97

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	801	A1J4W	C13-N14-S15-O17
3	A	801	A1J4W	C13-N14-S15-C18
3	B	801	A1J4W	C13-N14-S15-C18
3	A	801	A1J4W	C23-C18-S15-N14
3	B	801	A1J4W	C23-C18-S15-N14
3	A	801	A1J4W	C19-C18-S15-N14

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	801	A1J4W	3	0
3	A	801	A1J4W	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	269/272 (98%)	0.52	15 (5%)	30 26	30, 60, 121, 142	1 (0%)
1	B	269/272 (98%)	0.74	25 (9%)	14 12	31, 58, 133, 161	1 (0%)
All	All	538/544 (98%)	0.63	40 (7%)	20 18	30, 59, 129, 161	2 (0%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	577	SER	4.1
1	B	745	VAL	3.1
1	B	689	LEU	3.0
1	B	551	THR	2.9
1	B	682	PRO	2.9
1	A	630	GLY	2.8
1	B	776	TRP	2.8
1	A	745	VAL	2.8
1	B	753	ILE	2.7
1	B	775	ARG	2.6
1	A	531	TYR	2.6
1	A	770	ASP	2.6
1	A	549	SER	2.5
1	A	746	ILE	2.5
1	A	747	ILE	2.5
1	B	771	PRO	2.5
1	B	691	TYR	2.5
1	B	741	SER	2.5
1	B	767	VAL	2.5
1	B	722	ILE	2.4
1	B	690	SER	2.4
1	B	695	TRP	2.4
1	B	683	LEU	2.4
1	B	710	GLN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	749	ARG	2.3
1	B	608	TYR	2.3
1	B	748	ARG	2.2
1	A	711	ILE	2.2
1	B	762	LEU	2.2
1	B	729	SER	2.2
1	A	631	TYR	2.2
1	B	685	ASP	2.2
1	A	683	LEU	2.1
1	A	762	LEU	2.1
1	B	756	HIS	2.1
1	A	534	LEU	2.0
1	A	668	LEU	2.0
1	B	699	ILE	2.0
1	B	769	VAL	2.0
1	A	537	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	ALY	B	604	12/13	0.87	0.13	46,52,63,64	0
1	ALY	A	604	12/13	0.91	0.10	36,40,50,51	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	A1J4W	B	801	29/29	0.92	0.11	16,26,86,101	0
2	ZN	B	800	1/1	0.94	0.08	49,49,49,49	0
2	ZN	A	800	1/1	0.94	0.06	47,47,47,47	0
3	A1J4W	A	801	29/29	0.95	0.10	10,27,75,96	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around A1J4W B 801:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around A1J4W A 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.