



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 1, 2026 – 10:33 am BST

PDB ID : 29UT / pdb_000029ut
Title : KAT8 MUTANT (KAT6A SURROGATE, MYST-CRYST) IN COMPLEX
WITH KAT6A INHIBITOR COMPOUND 2
Authors : Hillig, R.C.; Puetter, V.
Deposited on : 2026-04-09
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	278	6% 82% 15% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	AAA	809	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2734 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Histone acetyltransferase KAT8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	AAA	273	2385	1562	386	424	13	0	19	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	502	GLY	-	expression tag	UNP Q9H7Z6
AAA	503	SER	-	expression tag	UNP Q9H7Z6
AAA	579	HIS	TYR	engineered mutation	UNP Q9H7Z6
AAA	645	SER	ALA	engineered mutation	UNP Q9H7Z6
AAA	648	MET	LEU	engineered mutation	UNP Q9H7Z6
AAA	649	ILE	THR	engineered mutation	UNP Q9H7Z6
AAA	660	ARG	LYS	engineered mutation	UNP Q9H7Z6
AAA	697	SER	TRP	engineered mutation	UNP Q9H7Z6
AAA	702	ASN	ILE	engineered mutation	UNP Q9H7Z6

- Molecule 2 is 6-(dimethylamino)-N-(2-methylquinolin-8-yl)sulfonyl-4-(trifluoromethyl)-1-benzofuran-2-carboxamide (CCD ID: A1J4V) (formula: C₂₂H₁₈F₃N₃O₄S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	AAA	1	Total	C	F	N	O	S	0	1
			61	42	3	5	8	3		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	AAA	1	Total Zn 1 1	0	0

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	AAA	5	Total Cl 5 5	0	0

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $\text{C}_3\text{H}_8\text{O}_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	AAA	1	Total	C	O	0	0
			6	3	3		
5	AAA	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	AAA	270	Total	O	0	0
			270	270		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histone acetyltransferase KAT8



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	46.27Å 57.35Å 121.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.65 – 1.40 41.65 – 1.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (41.65-1.40) 99.9 (41.65-1.40)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.25 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.129 , 0.178 0.152 , 0.188	Depositor DCC
R_{free} test set	2100 reflections (3.26%)	wwPDB-VP
Wilson B-factor (Å ²)	18.6	Xtriage
Anisotropy	0.332	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	2734	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, CL, A1J4V, ALY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.19	5/2493 (0.2%)	1.28	6/3371 (0.2%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	547	MET	CA-C	7.60	1.61	1.52
1	AAA	714	ASP	CG-OD1	7.50	1.39	1.25
1	AAA	531	TYR	C-O	5.30	1.30	1.24
1	AAA	579	HIS	CE1-NE2	5.28	1.37	1.32
1	AAA	740	LYS	C-O	5.16	1.29	1.23

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	534	GLN	CB-CG-CD	-6.53	101.51	112.60
1	AAA	534	GLN	N-CA-CB	6.23	118.38	109.78
1	AAA	585	ASP	CA-CB-CG	5.41	118.00	112.60
1	AAA	617	ILE	CA-CB-CG1	5.22	119.27	110.40
1	AAA	596	LEU	N-CA-C	-5.11	105.60	111.07
1	AAA	651	PRO	N-CA-C	5.08	116.90	110.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2385	0	2402	35	1
2	AAA	61	0	0	3	0
3	AAA	1	0	0	0	0
4	AAA	5	0	0	0	0
5	AAA	12	0	16	5	0
6	AAA	270	0	0	9	1
All	All	2734	0	2418	37	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:778:PRO:HG3	5:AAA:809:GOL:H32	1.42	0.97
1:AAA:660[A]:ARG:NH1	6:AAA:905:HOH:O	1.97	0.96
1:AAA:778:PRO:HG3	5:AAA:809:GOL:C3	2.16	0.76
1:AAA:620:GLU:OE2	1:AAA:658[B]:TYR:OH	2.04	0.75
1:AAA:614[A]:VAL:HG21	5:AAA:809:GOL:H2	1.66	0.74
1:AAA:530[A]:ASP:OD1	1:AAA:533:LYS:CE	2.41	0.68
1:AAA:576:ILE:HD11	1:AAA:658[B]:TYR:OH	1.96	0.66
1:AAA:738[B]:TYR:HE2	1:AAA:743:HIS:CE1	2.15	0.65
1:AAA:757[B]:LYS:HA	1:AAA:757[B]:LYS:HE2	1.78	0.65
1:AAA:757[B]:LYS:HA	1:AAA:757[B]:LYS:CE	2.30	0.62
1:AAA:712:ILE:HG13	1:AAA:745[A]:ILE:HG13	1.82	0.60
1:AAA:660[B]:ARG:NH2	6:AAA:909:HOH:O	2.37	0.58
1:AAA:778:PRO:CG	5:AAA:809:GOL:H32	2.26	0.58
1:AAA:774:LYS:HD2	6:AAA:961:HOH:O	2.02	0.58
1:AAA:660[B]:ARG:HD3	6:AAA:1010:HOH:O	2.07	0.54
1:AAA:658[B]:TYR:CE1	1:AAA:766[B]:ILE:HD13	2.45	0.52
1:AAA:511:LYS:HD2	1:AAA:518[B]:GLU:OE2	2.10	0.50
1:AAA:611:GLU:HG3	6:AAA:963:HOH:O	2.11	0.50
1:AAA:530[A]:ASP:OD1	1:AAA:533:LYS:HE2	2.11	0.49
1:AAA:530[A]:ASP:OD1	1:AAA:533:LYS:HE3	2.14	0.48
1:AAA:511:LYS:HG3	1:AAA:536[B]:LYS:HB3	1.95	0.48
1:AAA:738[B]:TYR:CE2	1:AAA:743:HIS:ND1	2.82	0.47
1:AAA:738[B]:TYR:CE2	1:AAA:743:HIS:CE1	2.99	0.46
1:AAA:656:ARG:NH1	6:AAA:907:HOH:O	2.35	0.45
1:AAA:533:LYS:HE3	1:AAA:533:LYS:HB3	1.78	0.45
1:AAA:673:GLU:HG2	1:AAA:777:PRO:HB3	1.98	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:603:HIS:CG	1:AAA:682:PRO:HG2	2.53	0.44
1:AAA:606:LEU:C	1:AAA:606:LEU:HD23	2.42	0.44
1:AAA:614[A]:VAL:HG21	5:AAA:809:GOL:C2	2.43	0.44
1:AAA:655[A]:ARG:NH2	6:AAA:906:HOH:O	2.22	0.43
1:AAA:511:LYS:HE2	6:AAA:942:HOH:O	2.19	0.42
1:AAA:747:VAL:HG23	1:AAA:751:LEU:HD23	2.02	0.42
1:AAA:662:LEU:HD23	1:AAA:662:LEU:HA	1.93	0.41
1:AAA:655[A]:ARG:HA	2:AAA:801[A]:A1J4V:C29	2.50	0.41
1:AAA:557:LEU:HD21	6:AAA:1043:HOH:O	2.21	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:774:LYS:NZ	6:AAA:984:HOH:O[3_645]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	AAA	290/278 (104%)	279 (96%)	10 (3%)	1 (0%)	36 16

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	707	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	AAA	270/256 (106%)	266 (98%)	4 (2%)	57 26

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	568	LYS
1	AAA	631	TYR
1	AAA	734	ASN
1	AAA	774	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	ALY	AAA	604	1	10,11,12	0.78	0	7,12,14	1.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	ALY	AAA	604	1	-	1/9/10/12	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	AAA	604	ALY	CG-CD-CE-NZ

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 6 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	A1J4V	AAA	801[A]	-	36,36,36	1.92	6 (16%)	52,56,56	2.59	15 (28%)
5	GOL	AAA	808	-	5,5,5	0.12	0	5,5,5	0.41	0
2	A1J4V	AAA	801[B]	-	36,36,36	2.11	8 (22%)	52,56,56	2.61	16 (30%)
5	GOL	AAA	809	-	5,5,5	0.10	0	5,5,5	0.71	0
2	A1J4V	AAA	801[C]	-	36,36,36	2.08	8 (22%)	52,56,56	2.30	15 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1J4V	AAA	801[A]	-	-	3/25/25/25	0/4/4/4
5	GOL	AAA	808	-	-	2/4/4/4	-
2	A1J4V	AAA	801[B]	-	-	3/25/25/25	0/4/4/4
5	GOL	AAA	809	-	-	4/4/4/4	-
2	A1J4V	AAA	801[C]	-	-	2/25/25/25	0/4/4/4

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	AAA	801[A]	A1J4V	O18-S16	7.26	1.51	1.43
2	AAA	801[C]	A1J4V	O18-S16	6.65	1.51	1.43
2	AAA	801[B]	A1J4V	O18-S16	5.48	1.49	1.43
2	AAA	801[A]	A1J4V	C13-N15	-5.27	1.33	1.38
2	AAA	801[B]	A1J4V	C13-N15	-5.27	1.33	1.38
2	AAA	801[C]	A1J4V	C13-N15	-5.27	1.33	1.38
2	AAA	801[C]	A1J4V	C19-C29	-4.88	1.39	1.42
2	AAA	801[B]	A1J4V	C19-S16	-4.85	1.72	1.77
2	AAA	801[B]	A1J4V	S16-N15	-4.46	1.55	1.64
2	AAA	801[B]	A1J4V	O17-S16	4.10	1.48	1.43
2	AAA	801[C]	A1J4V	C19-S16	3.68	1.81	1.77
2	AAA	801[B]	A1J4V	C19-C29	-3.53	1.40	1.42
2	AAA	801[A]	A1J4V	C20-C19	3.14	1.41	1.37
2	AAA	801[C]	A1J4V	C20-C19	3.03	1.41	1.37
2	AAA	801[C]	A1J4V	O17-S16	2.71	1.46	1.43
2	AAA	801[A]	A1J4V	O17-S16	2.57	1.46	1.43
2	AAA	801[A]	A1J4V	C27-C26	2.50	1.56	1.50
2	AAA	801[B]	A1J4V	C27-C26	2.13	1.55	1.50
2	AAA	801[A]	A1J4V	C12-C4	-2.06	1.35	1.39
2	AAA	801[B]	A1J4V	C12-C4	-2.06	1.35	1.39
2	AAA	801[C]	A1J4V	C12-C4	-2.06	1.35	1.39
2	AAA	801[C]	A1J4V	C23-C29	-2.00	1.39	1.42

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	AAA	801[A]	A1J4V	O18-S16-O17	-10.00	107.25	119.55
2	AAA	801[B]	A1J4V	O18-S16-O17	-9.79	107.52	119.55
2	AAA	801[C]	A1J4V	O18-S16-O17	-6.86	111.12	119.55
2	AAA	801[A]	A1J4V	C13-N15-S16	-6.62	115.03	123.78
2	AAA	801[A]	A1J4V	C19-C29-N28	6.38	121.95	119.55

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	AAA	801[C]	A1J4V	C20-C19-C29	-5.94	116.34	121.04
2	AAA	801[C]	A1J4V	C20-C19-S16	5.55	123.56	116.98
2	AAA	801[B]	A1J4V	C13-N15-S16	5.34	130.85	123.78
2	AAA	801[B]	A1J4V	C20-C19-S16	5.32	123.29	116.98
2	AAA	801[B]	A1J4V	C19-S16-N15	-5.24	98.37	107.29
2	AAA	801[A]	A1J4V	O14-C13-N15	5.01	128.81	121.67
2	AAA	801[B]	A1J4V	O14-C13-N15	5.01	128.81	121.67
2	AAA	801[C]	A1J4V	O14-C13-N15	5.01	128.81	121.67
2	AAA	801[B]	A1J4V	C29-C19-S16	-4.98	115.47	120.76
2	AAA	801[A]	A1J4V	O18-S16-N15	4.82	120.30	106.74
2	AAA	801[C]	A1J4V	C21-C20-C19	4.58	125.09	120.37
2	AAA	801[B]	A1J4V	C19-C29-C23	4.01	119.91	117.68
2	AAA	801[C]	A1J4V	C19-S16-N15	3.97	114.06	107.29
2	AAA	801[B]	A1J4V	O17-S16-C19	3.29	113.90	108.08
2	AAA	801[A]	A1J4V	C9-C13-N15	-3.07	110.92	115.49
2	AAA	801[B]	A1J4V	C9-C13-N15	-3.07	110.92	115.49
2	AAA	801[C]	A1J4V	C9-C13-N15	-3.07	110.92	115.49
2	AAA	801[A]	A1J4V	C19-C29-C23	2.99	119.34	117.68
2	AAA	801[A]	A1J4V	C23-C29-N28	-2.97	118.70	122.64
2	AAA	801[A]	A1J4V	O18-S16-C19	-2.96	102.85	108.08
2	AAA	801[B]	A1J4V	O18-S16-N15	2.93	114.99	106.74
2	AAA	801[B]	A1J4V	C21-C20-C19	-2.90	117.38	120.37
2	AAA	801[C]	A1J4V	C19-C29-C23	2.81	119.25	117.68
2	AAA	801[C]	A1J4V	C22-C23-C29	2.70	122.67	118.26
2	AAA	801[A]	A1J4V	C20-C19-S16	2.69	120.17	116.98
2	AAA	801[C]	A1J4V	C19-C29-N28	2.65	120.55	119.55
2	AAA	801[A]	A1J4V	O17-S16-C19	2.63	112.72	108.08
2	AAA	801[B]	A1J4V	C27-C26-N28	2.58	121.44	117.23
2	AAA	801[A]	A1J4V	C4-C5-C6	2.56	124.31	119.86
2	AAA	801[B]	A1J4V	C4-C5-C6	2.56	124.31	119.86
2	AAA	801[C]	A1J4V	C4-C5-C6	2.56	124.31	119.86
2	AAA	801[C]	A1J4V	C22-C21-C20	-2.55	116.92	120.99
2	AAA	801[A]	A1J4V	C1-N2-C4	-2.34	116.54	120.54
2	AAA	801[B]	A1J4V	C1-N2-C4	-2.34	116.54	120.54
2	AAA	801[C]	A1J4V	C1-N2-C4	-2.34	116.54	120.54
2	AAA	801[B]	A1J4V	O17-S16-N15	2.28	113.15	106.74
2	AAA	801[C]	A1J4V	C13-N15-S16	2.23	126.73	123.78
2	AAA	801[A]	A1J4V	C20-C19-C29	-2.13	119.35	121.04
2	AAA	801[A]	A1J4V	F33-C30-C6	-2.09	109.06	112.70
2	AAA	801[B]	A1J4V	F33-C30-C6	-2.09	109.06	112.70
2	AAA	801[C]	A1J4V	F33-C30-C6	-2.09	109.06	112.70

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	AAA	801[B]	A1J4V	C13-N15-S16-C19
2	AAA	801[B]	A1J4V	C13-N15-S16-O17
2	AAA	801[B]	A1J4V	C13-N15-S16-O18
5	AAA	808	GOL	C1-C2-C3-O3
5	AAA	809	GOL	O1-C1-C2-O2
5	AAA	809	GOL	O1-C1-C2-C3
2	AAA	801[C]	A1J4V	C13-N15-S16-O17
2	AAA	801[A]	A1J4V	C13-N15-S16-O17
2	AAA	801[C]	A1J4V	C13-N15-S16-C19
5	AAA	809	GOL	C1-C2-C3-O3
5	AAA	808	GOL	O2-C2-C3-O3
5	AAA	809	GOL	O2-C2-C3-O3
2	AAA	801[A]	A1J4V	C13-N15-S16-O18
2	AAA	801[A]	A1J4V	C13-N15-S16-C19


There are no ring outliers.

3 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	AAA	801[A]	A1J4V	1	0
2	AAA	801[B]	A1J4V	2	0
5	AAA	809	GOL	5	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand A1J4V AAA 801 (A)



Ligand A1J4V AAA 801 (B)





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	AAA	272/278 (97%)	-0.04	16 (5%)	28 30	9, 23, 63, 187	19 (6%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	708	GLY	4.5
1	AAA	763	LYS	4.1
1	AAA	778	PRO	4.1
1	AAA	779	LYS	3.6
1	AAA	741	GLY	3.3
1	AAA	762	LYS	3.2
1	AAA	759	ALA	3.1
1	AAA	761	TYR	2.9
1	AAA	739	TRP	2.7
1	AAA	706	PHE	2.6
1	AAA	533	LYS	2.4
1	AAA	757[A]	LYS	2.2
1	AAA	709	THR	2.1
1	AAA	707	ARG	2.1
1	AAA	777	PRO	2.1
1	AAA	710	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
1	ALY	AAA	604	12/13	0.98	0.04	17,19,22,25	0

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	AAA	809	6/6	0.87	0.16	33,43,46,56	6
5	GOL	AAA	808	6/6	0.95	0.09	16,25,28,34	6
2	A1J4V	AAA	801[C]	33/33	0.99	0.07	9,15,20,21	14
4	CL	AAA	805	1/1	0.99	0.06	26,26,26,26	1
4	CL	AAA	806	1/1	0.99	0.05	28,28,28,28	1
2	A1J4V	AAA	801[A]	33/33	0.99	0.07	14,18,38,43	14
2	A1J4V	AAA	801[B]	33/33	0.99	0.07	14,16,21,24	14
4	CL	AAA	803	1/1	1.00	0.02	17,17,17,17	0
4	CL	AAA	807	1/1	1.00	0.05	31,31,31,31	0
4	CL	AAA	804	1/1	1.00	0.04	22,22,22,22	0
3	ZN	AAA	802	1/1	1.00	0.01	15,15,15,15	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1J4V AAA 801 (C):

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around A1J4V AAA 801 (A):**

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around A1J4V AAA 801 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.