



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 1, 2026 – 10:31 am BST

PDB ID : 29UU / pdb_000029uu
Title : KAT8 MUTANT (KAT6A SURROGATE, MYST-CRYST) IN COMPLEX
WITH KAT6A INHIBITOR Example-41
Authors : Hillig, R.C.; Puetter, V.
Deposited on : 2026-04-09
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1553 (1.30-1.30)
Clashscore	190562	1595 (1.30-1.30)
Ramachandran outliers	187476	1551 (1.30-1.30)
Sidechain outliers	187428	1551 (1.30-1.30)
RSRZ outliers	180081	1549 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	278	5% 87% 10%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 2765 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone acetyltransferase KAT8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	269	Total	C	N	O	S	0	25	0
			2366	1554	386	413	13			


There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	502	GLY	-	expression tag	UNP Q9H7Z6
AAA	503	SER	-	expression tag	UNP Q9H7Z6
AAA	579	HIS	TYR	engineered mutation	UNP Q9H7Z6
AAA	645	SER	ALA	engineered mutation	UNP Q9H7Z6
AAA	648	MET	LEU	engineered mutation	UNP Q9H7Z6
AAA	649	ILE	THR	engineered mutation	UNP Q9H7Z6
AAA	660	ARG	LYS	engineered mutation	UNP Q9H7Z6
AAA	697	SER	TRP	engineered mutation	UNP Q9H7Z6
AAA	702	ASN	ILE	engineered mutation	UNP Q9H7Z6

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	AAA	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 7-iodanyl-N-[(2R)-2-(1,3-oxazol-2-yl)-2-phenyl-ethyl]-1,1-bis(oxidanylidene)-2H-1⁶,2,4-benzothiadiazine-3-carboxamide (CCD ID: A1J4U) (formula: C₁₉H₁₅IN₄O₄S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	AAA	1	Total	C	I	N	O	S	0	0
			29	19	1	4	4	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $\text{C}_3\text{H}_8\text{O}_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	AAA	1	Total C O 6 3 3	0	0
4	AAA	1	Total C O 6 3 3	0	0
4	AAA	1	Total C O 6 3 3	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	AAA	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	AAA	5	Total	Cl		0	0
			5	5			

- Molecule 6 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	AAA	1	Total	C	O	0	0
			7	4	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	AAA	333	Total	O		0	0
			333	333			

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histone acetyltransferase KAT8



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	46.46Å 57.72Å 121.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.38 – 1.30 43.38 – 1.30	Depositor EDS
% Data completeness (in resolution range)	97.7 (43.38-1.30) 97.7 (43.38-1.30)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 1.30Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.124 , 0.161 0.140 , 0.170	Depositor DCC
R_{free} test set	2101 reflections (2.65%)	wwPDB-VP
Wilson B-factor (Å ²)	16.6	Xtriage
Anisotropy	0.122	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 50.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	2765	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, GOL, PEG, ZN, ALY, A1J4U

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.00	3/2491 (0.1%)	1.06	5/3363 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	701	GLU	CD-OE1	6.27	1.37	1.25
1	AAA	701	GLU	CD-OE2	5.33	1.35	1.25
1	AAA	764	PRO	CA-C	5.29	1.57	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	764	PRO	CB-CA-C	8.67	121.49	110.92
1	AAA	620	GLU	CB-CG-CD	-6.48	101.58	112.60
1	AAA	651	PRO	N-CA-C	6.18	118.25	110.70
1	AAA	764	PRO	O-C-N	-5.25	115.27	121.46
1	AAA	714	ASP	CA-CB-CG	5.17	117.77	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2366	0	2426	24	1
2	AAA	1	0	0	0	0
3	AAA	29	0	0	0	0
4	AAA	24	0	32	0	0
5	AAA	5	0	0	0	0
6	AAA	7	0	10	0	0
7	AAA	333	0	0	9	2
All	All	2765	0	2468	24	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:585:ASP:OD2	7:AAA:905:HOH:O	1.73	1.03
1:AAA:656:ARG:NH1	7:AAA:906:HOH:O	1.89	0.99
1:AAA:587[B]:LYS:HZ3	1:AAA:611[B]:GLU:CD	1.74	0.95
1:AAA:587[B]:LYS:NZ	1:AAA:611[B]:GLU:OE1	1.99	0.95
1:AAA:587[B]:LYS:NZ	1:AAA:611[B]:GLU:CD	2.22	0.95
1:AAA:747:VAL:HG23	1:AAA:751:LEU:HD23	1.70	0.73
1:AAA:611[B]:GLU:HG3	7:AAA:1012:HOH:O	1.90	0.72
1:AAA:606:LEU:HD21	1:AAA:609:ASP:HB3	1.71	0.71
1:AAA:587[B]:LYS:HZ2	1:AAA:611[B]:GLU:CD	1.99	0.70
1:AAA:689[B]:LEU:HG	7:AAA:1057:HOH:O	1.95	0.67
1:AAA:717[B]:GLN:HG3	7:AAA:985:HOH:O	1.99	0.62
1:AAA:692[B]:ARG:HG3	7:AAA:987:HOH:O	2.02	0.60
1:AAA:745[B]:ILE:HD13	1:AAA:746:CYS:N	2.18	0.58
1:AAA:730:LEU:HB3	1:AAA:736[B]:VAL:HG13	1.85	0.58
1:AAA:603[B]:HIS:CD2	7:AAA:1125:HOH:O	2.60	0.54
1:AAA:745[B]:ILE:HD11	1:AAA:747:VAL:HG12	1.93	0.50
1:AAA:606:LEU:C	1:AAA:606:LEU:HD13	2.37	0.50
1:AAA:730:LEU:CB	1:AAA:736[B]:VAL:HG13	2.42	0.49
1:AAA:511:LYS:HE2	1:AAA:536[B]:LYS:HD3	1.97	0.46
1:AAA:584:LYS:HE2	1:AAA:611[A]:GLU:CG	2.47	0.44
1:AAA:745[B]:ILE:CD1	1:AAA:747:VAL:HG12	2.47	0.44
1:AAA:587[B]:LYS:HD2	7:AAA:1012:HOH:O	2.18	0.43
1:AAA:689[B]:LEU:HD12	7:AAA:1181:HOH:O	2.18	0.43
1:AAA:584:LYS:HE2	1:AAA:611[A]:GLU:OE2	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:545[B]:LYS:NZ	7:AAA:1083:HOH:O[4_575]	1.94	0.26
7:AAA:1091:HOH:O	7:AAA:1187:HOH:O[4_475]	2.04	0.16

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	289/278 (104%)	278 (96%)	11 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	272/256 (106%)	269 (99%)	3 (1%)	65	33

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	607	TYR
1	AAA	707[A]	ARG
1	AAA	707[B]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	ALY	AAA	604	1	10,11,12	0.81	0	7,12,14	1.33	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	ALY	AAA	604	1	-	2/9/10/12	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	604	ALY	CG-CD-CE	2.86	127.10	113.56

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	AAA	604	ALY	CG-CD-CE-NZ
1	AAA	604	ALY	CE-CD-CG-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	AAA	808	-	5,5,5	0.11	0	5,5,5	0.33	0
6	PEG	AAA	810	-	6,6,6	0.21	0	5,5,5	0.22	0
4	GOL	AAA	812	-	5,5,5	0.17	0	5,5,5	0.34	0
4	GOL	AAA	803	-	5,5,5	0.59	0	5,5,5	0.81	0
4	GOL	AAA	811	-	5,5,5	0.15	0	5,5,5	0.19	0
3	A1J4U	AAA	802	-	28,32,32	2.36	9 (32%)	34,46,46	2.59	14 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	AAA	808	-	-	2/4/4/4	-
6	PEG	AAA	810	-	-	2/4/4/4	-
4	GOL	AAA	812	-	-	3/4/4/4	-
4	GOL	AAA	803	-	-	0/4/4/4	-
4	GOL	AAA	811	-	-	0/4/4/4	-
3	A1J4U	AAA	802	-	-	0/13/32/32	0/4/4/4

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	AAA	802	A1J4U	S25-N22	-6.14	1.56	1.62
3	AAA	802	A1J4U	C26-S25	-4.89	1.71	1.75
3	AAA	802	A1J4U	O8-C7	4.78	1.52	1.37
3	AAA	802	A1J4U	C14-C15	-3.79	1.32	1.39
3	AAA	802	A1J4U	O8-C9	-3.43	1.31	1.36
3	AAA	802	A1J4U	C12-C11	-3.09	1.29	1.38
3	AAA	802	A1J4U	O23-S25	2.74	1.46	1.43
3	AAA	802	A1J4U	C21-N4	2.58	1.36	1.30
3	AAA	802	A1J4U	C11-C10	-2.03	1.34	1.38

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	AAA	802	A1J4U	O24-S25-O23	-7.35	110.96	118.46
3	AAA	802	A1J4U	C16-C9-N5	-5.89	118.74	127.93
3	AAA	802	A1J4U	C26-S25-N22	4.55	106.01	102.13
3	AAA	802	A1J4U	C27-C26-S25	3.49	122.68	119.82
3	AAA	802	A1J4U	C11-C10-C15	3.48	124.95	120.65
3	AAA	802	A1J4U	O24-S25-N22	3.37	111.00	108.26
3	AAA	802	A1J4U	C3-N4-C21	3.09	122.35	117.55
3	AAA	802	A1J4U	O8-C9-C16	3.00	124.43	118.38
3	AAA	802	A1J4U	O23-S25-C26	-2.87	105.74	109.29
3	AAA	802	A1J4U	C7-C6-N5	-2.79	110.81	115.95
3	AAA	802	A1J4U	O23-S25-N22	2.74	110.49	108.26
3	AAA	802	A1J4U	C12-C11-C10	-2.73	116.03	120.19
3	AAA	802	A1J4U	O24-S25-C26	2.52	112.41	109.29
3	AAA	802	A1J4U	C26-C27-C28	-2.52	118.31	121.28

There are no chirality outliers.


All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	AAA	808	GOL	O1-C1-C2-C3
4	AAA	812	GOL	C1-C2-C3-O3
4	AAA	812	GOL	O2-C2-C3-O3
4	AAA	812	GOL	O1-C1-C2-C3
4	AAA	808	GOL	O1-C1-C2-O2
6	AAA	810	PEG	C1-C2-O2-C3
6	AAA	810	PEG	O2-C3-C4-O4

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	AAA	268/278 (96%)	0.12	14 (5%)	33 37	8, 18, 43, 96	25 (9%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	707[A]	ARG	6.9
1	AAA	708	GLY	4.5
1	AAA	562[A]	TRP	4.5
1	AAA	608	PHE	4.0
1	AAA	706	PHE	3.5
1	AAA	778	PRO	3.3
1	AAA	607	TYR	3.1
1	AAA	606	LEU	2.9
1	AAA	709	THR	2.8
1	AAA	751	LEU	2.6
1	AAA	605	THR	2.4
1	AAA	758	SER	2.2
1	AAA	777	PRO	2.2
1	AAA	757	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	ALY	AAA	604	12/13	0.96	0.07	15,20,23,28	0

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	AAA	812	6/6	0.85	0.14	34,49,72,82	0
4	GOL	AAA	808	6/6	0.87	0.14	58,61,65,66	0
4	GOL	AAA	811	6/6	0.89	0.13	24,40,57,59	0
6	PEG	AAA	810	7/7	0.92	0.12	42,54,60,61	0
4	GOL	AAA	803	6/6	0.97	0.05	18,19,23,24	0
5	CL	AAA	804	1/1	0.98	0.10	33,33,33,33	0
2	ZN	AAA	801	1/1	1.00	0.01	13,13,13,13	0
5	CL	AAA	805	1/1	1.00	0.01	15,15,15,15	0
5	CL	AAA	806	1/1	1.00	0.14	30,30,30,30	0
5	CL	AAA	807	1/1	1.00	0.10	27,27,27,27	0
5	CL	AAA	809	1/1	1.00	0.03	16,16,16,16	1
3	A1J4U	AAA	802	29/29	1.00	0.03	12,15,20,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1J4U AAA 802:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.