Full wwPDB EM Validation Report ⓘ

Nov 19, 2026 – 03:17 pm GMT

PDB ID	:	29XV / pdb_000029xv
EMDB ID	:	EMD-57438
Title	:	Vaccinia Virus A10 trimer structure
Authors	:	Arragain, B.; Schur, F.K.M.
Deposited on	:	2026-04-14
Resolution	:	2.60 Å(reported)
Based on initial model	:	.

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev133
MolProbity	:	4-5-2 with Phenix2.2
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	8728 (2.10 - 3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	622	13% 87% 6% 7%
1	B	622	12% 88% 5% 7%
1	C	622	12% 88% 5% 7%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 14346 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Core protein OPG136.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	B	581	Total	C	N	O	S	0	0
			4710	3036	776	874	24		
1	A	581	Total	C	N	O	S	0	0
			4710	3036	776	874	24		
1	C	581	Total	C	N	O	S	0	0
			4710	3036	776	874	24		

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	615	GLY	-	expression tag	UNP P16715
B	616	SER	-	expression tag	UNP P16715
B	617	HIS	-	expression tag	UNP P16715
B	618	HIS	-	expression tag	UNP P16715
B	619	HIS	-	expression tag	UNP P16715
B	620	HIS	-	expression tag	UNP P16715
B	621	HIS	-	expression tag	UNP P16715
B	622	HIS	-	expression tag	UNP P16715
A	615	GLY	-	expression tag	UNP P16715
A	616	SER	-	expression tag	UNP P16715
A	617	HIS	-	expression tag	UNP P16715
A	618	HIS	-	expression tag	UNP P16715
A	619	HIS	-	expression tag	UNP P16715
A	620	HIS	-	expression tag	UNP P16715
A	621	HIS	-	expression tag	UNP P16715
A	622	HIS	-	expression tag	UNP P16715
C	615	GLY	-	expression tag	UNP P16715
C	616	SER	-	expression tag	UNP P16715
C	617	HIS	-	expression tag	UNP P16715
C	618	HIS	-	expression tag	UNP P16715
C	619	HIS	-	expression tag	UNP P16715
C	620	HIS	-	expression tag	UNP P16715
C	621	HIS	-	expression tag	UNP P16715
C	622	HIS	-	expression tag	UNP P16715

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	AltConf
2	B	72	Total O 72 72	0
2	A	72	Total O 72 72	0
2	C	72	Total O 72 72	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Core protein OPG136



• Molecule 1: Core protein OPG136



• Molecule 1: Core protein OPG136




MET M2 R21 K35 I69 T81 S82 R83 E84 F119 L122 M156 L174 D181 V182 I183 N184 K188 I194 V195 N196 G197 V198 I208 R211 Q212 I216 Q217 D218 R219 P220 D223 D224 A229 D230 A231 A232 R233 G236 V237 N238 L239 N240
T241 L242 P243 L244 P245 N246 V247 D248 L249 T250 T251 M252 Y267 D270 D273 I274 Y275 Y276 N277 G278 N279 K280 M281 L282 F283 D284 D285 E286 I287 H319 T320 R321 D322 P323 Q324 N325 A326 A327 V328 N329 V330 L334 P335 N336 N343 K347 L386 M390 I416
K421 R425 E456 R459 K460 D461 K462 F478 L483 D486 K508 D521 T536 R537 ARG SER ILE MET GLY SER ASN ASN ILE SER I548 I549 R552 P553 R554 S590 V591 S592 ASP PHE ARG ASP TYR GLN SER TYR ARG GLN TYR ARG ASN TYR CYS PRO ARG
TYR PHE TYR ALA GLY GLY SER HIS HIS HIS HIS HIS HIS

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C3	Depositor
Number of particles used	182668	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	40	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.734	Depositor
Minimum map value	-0.547	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.017	Depositor
Recommended contour level	0.1	Depositor
Map size (Å)	282.24, 282.24, 282.24	wwPDB
Map dimensions	336, 336, 336	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.84, 0.84, 0.84	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	0.09	0/4810	0.22	0/6522
1	B	0.09	0/4810	0.22	0/6522
1	C	0.09	0/4810	0.22	0/6522
All	All	0.09	0/14430	0.22	0/19566

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4710	0	4746	22	0
1	B	4710	0	4746	18	0
1	C	4710	0	4746	20	0
2	A	72	0	0	0	0
2	B	72	0	0	0	0
2	C	72	0	0	0	0
All	All	14346	0	14238	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:284:ASP:HA	1:C:287:ILE:HD12	1.87	0.56
1:B:284:ASP:HA	1:B:287:ILE:HD12	1.87	0.56
1:A:284:ASP:HA	1:A:287:ILE:HD12	1.87	0.55
1:C:281:MET:HG3	1:C:320:THR:HG22	1.88	0.55
1:A:183:ILE:HG13	1:A:194:ILE:HD11	1.90	0.54
1:A:281:MET:HG3	1:A:320:THR:HG22	1.88	0.54
1:B:281:MET:HG3	1:B:320:THR:HG22	1.88	0.54
1:B:183:ILE:HG13	1:B:194:ILE:HD11	1.90	0.53
1:C:183:ILE:HG13	1:C:194:ILE:HD11	1.90	0.52
1:B:81:THR:HG23	1:B:347:LYS:HE2	1.91	0.52
1:A:81:THR:HG23	1:A:347:LYS:HE2	1.91	0.52
1:C:81:THR:HG23	1:C:347:LYS:HE2	1.91	0.52
1:A:416:ILE:HG23	1:A:421:LYS:HB2	1.94	0.50
1:C:416:ILE:HG23	1:C:421:LYS:HB2	1.94	0.49
1:B:416:ILE:HG23	1:B:421:LYS:HB2	1.94	0.48
1:C:35:LYS:HD3	1:C:35:LYS:HA	1.63	0.48
1:A:156:MET:HB2	1:A:425:ARG:HD2	1.96	0.48
1:C:119:PHE:HB3	1:C:122:LEU:HD12	1.96	0.48
1:A:119:PHE:HB3	1:A:122:LEU:HD12	1.96	0.47
1:B:119:PHE:HB3	1:B:122:LEU:HD12	1.96	0.47
1:B:156:MET:HB2	1:B:425:ARG:HD2	1.96	0.47
1:C:156:MET:HB2	1:C:425:ARG:HD2	1.96	0.46
1:B:212:GLN:HB2	1:B:245:PRO:HB3	1.98	0.46
1:C:21:ARG:NH2	1:C:483:LEU:HB2	2.32	0.45
1:C:212:GLN:HB2	1:C:245:PRO:HB3	1.98	0.45
1:A:21:ARG:NH2	1:A:483:LEU:HB2	2.32	0.45
1:B:260:MET:HE2	1:B:260:MET:HB2	1.82	0.45
1:A:212:GLN:HB2	1:A:245:PRO:HB3	1.98	0.45
1:C:83:ARG:NH2	1:C:347:LYS:HD3	2.32	0.45
1:A:83:ARG:NH2	1:A:347:LYS:HD3	2.32	0.44
1:A:35:LYS:HD3	1:A:35:LYS:HA	1.63	0.44
1:B:21:ARG:NH2	1:B:483:LEU:HB2	2.32	0.44
1:B:83:ARG:NH2	1:B:347:LYS:HD3	2.33	0.44
1:B:478:PHE:HZ	1:C:390:MET:HG2	1.83	0.44
1:C:508:LYS:HB2	1:C:508:LYS:HE3	1.79	0.44
1:A:390:MET:HG2	1:C:478:PHE:HZ	1.83	0.43
1:B:390:MET:HG2	1:A:478:PHE:HZ	1.83	0.43
1:B:386:LEU:O	1:B:390:MET:HG3	2.19	0.43
1:A:386:LEU:O	1:A:390:MET:HG3	2.19	0.43
1:C:208:ILE:HD13	1:C:208:ILE:HA	1.84	0.43
1:C:386:LEU:O	1:C:390:MET:HG3	2.19	0.43
1:C:536:THR:HG22	1:C:549:ILE:HG12	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:536:THR:HG22	1:A:549:ILE:HG12	2.01	0.42
1:B:536:THR:HG22	1:B:549:ILE:HG12	2.01	0.42
1:A:508:LYS:HE3	1:A:508:LYS:HB2	1.79	0.42
1:A:453:THR:O	1:A:579:THR:HG21	2.20	0.41
1:B:249:LEU:HD23	1:B:252:MET:SD	2.61	0.41
1:B:271:ARG:HE	1:B:271:ARG:HB3	1.78	0.41
1:A:69:ILE:HG23	1:A:174:LEU:HD12	2.02	0.41
1:A:260:MET:HB2	1:A:260:MET:HE2	1.82	0.41
1:C:69:ILE:HG23	1:C:174:LEU:HD12	2.02	0.41
1:A:77:LEU:HD23	1:A:77:LEU:HA	1.94	0.41
1:A:82:SER:O	1:A:83:ARG:HG2	2.21	0.41
1:A:249:LEU:HD23	1:A:252:MET:SD	2.61	0.41
1:B:453:THR:O	1:B:579:THR:HG21	2.20	0.40
1:C:82:SER:O	1:C:83:ARG:HG2	2.21	0.40
1:C:249:LEU:HD23	1:C:252:MET:SD	2.61	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	577/622 (93%)	568 (98%)	9 (2%)	0	100	100
1	B	577/622 (93%)	568 (98%)	9 (2%)	0	100	100
1	C	577/622 (93%)	568 (98%)	9 (2%)	0	100	100
All	All	1731/1866 (93%)	1704 (98%)	27 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	537/574 (94%)	533 (99%)	4 (1%)	76 89
1	B	537/574 (94%)	534 (99%)	3 (1%)	78 91
1	C	537/574 (94%)	535 (100%)	2 (0%)	84 93
All	All	1611/1722 (94%)	1602 (99%)	9 (1%)	76 91

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	208	ILE
1	B	486	ASP
1	B	590	SER
1	A	208	ILE
1	A	486	ASP
1	A	510	SER
1	A	590	SER
1	C	208	ILE
1	C	590	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	105	ASN
1	B	115	ASN
1	B	168	ASN
1	B	279	ASN
1	B	572	ASN
1	A	93	ASN
1	A	96	ASN
1	A	115	ASN
1	A	168	ASN
1	A	279	ASN
1	A	572	ASN
1	C	115	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	168	ASN
1	C	279	ASN
1	C	338	GLN
1	C	572	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Map visualisation

This section contains visualisations of the EMDB entry EMD-57438. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections

6.1.1 Primary map



X



Y



Z

6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map



X Index: 168



Y Index: 168



Z Index: 168

6.2.2 Raw map



X Index: 168



Y Index: 168



Z Index: 168

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 151



Y Index: 188



Z Index: 170

6.3.2 Raw map



X Index: 149



Y Index: 161



Z Index: 170

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



X



Y



Z

6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map



X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.1. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

6.6.1 emd_57438_msk_1.map (i)



X



Y



Z

7 Map analysis

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate (i)



The volume at the recommended contour level is 60 nm^3; this corresponds to an approximate mass of 54 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.385 Å^{-1}

8 Fourier-Shell correlation

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC



*Reported resolution corresponds to spatial frequency of 0.385 Å^{-1}

8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.60	-	-
Author-provided FSC curve	2.60	2.90	2.63
Unmasked-calculated*	3.28	3.68	3.33

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.28 differs from the reported value 2.6 by more than 10 %

9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-57438 and PDB model 29XV. Per-residue inclusion information can be found in section 3 on page 5.

9.1 Map-model overlay



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.1 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.1).

9.4 Atom inclusion (i)



At the recommended contour level, 78% of all backbone atoms, 76% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.1) and Q-score for the entire model and for each chain.



Chain	Atom inclusion	Q-score
All	0.7590	0.6050
A	0.7610	0.6050
B	0.7620	0.6050
C	0.7620	0.6040