



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 24, 2025 – 03:38 am BST

PDB ID : 2BXZ / pdb_00002bxz
Title : Is radiation damage dependent on the dose-rate used during macromolecular crystallography data collection
Authors : Leiros, H.-K.S.; Timmins, J.; Ravelli, R.B.G.; McSweeney, S.M.
Deposited on : 2005-07-28
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2888 (1.76-1.76)
Clashscore	180529	3097 (1.76-1.76)
Ramachandran outliers	177936	3072 (1.76-1.76)
Sidechain outliers	177891	3072 (1.76-1.76)
RSRZ outliers	164620	2887 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	602	 84% 11%
2	B	2	 100%
2	C	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GLC	A	702	X	-	-	-

2 Entry composition


There are 7 unique types of molecules in this entry. The entry contains 5804 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MALTOOLIGOSYLTREHALOSE TREHALOHYDROLASE.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	580	4865	3076	868	909	2	10	31	29	0

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	2	23	12	11	0	0	0
2	C	2	23	12	11	0	0	0

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
3	A	1	8	4	1	3	0	0

- Molecule 4 is alpha-D-glucopyranose (CCD ID: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	12	6	6	0	0
4	A	1	12	6	6	0	0

- Molecule 5 is BETA-MERCAPTOETHANOL (CCD ID: BME) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
5	A	1	4	2	1	1	0	0

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
6	A	1	1	1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	856	856	856	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MALTOOLIGOSYLTREHALOSE TREHALOHYDROLASE



- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose



- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.42Å 66.53Å 152.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.75 20.00 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.5 (20.00-1.75) 99.4 (20.00-1.75)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 1.76Å)	Xtrriage
Refinement program	REFMAC 5.2.0003	Depositor
R, R_{free}	0.149 , 0.188 0.149 , 0.190	Depositor DCC
R_{free} test set	2479 reflections (4.04%)	wwPDB-VP
Wilson B-factor (Å ²)	13.6	Xtrriage
Anisotropy	0.011	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 70.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5804	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, GLC, MG, BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.02	8/4992 (0.2%)	0.97	12/6780 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	580	ARG	CZ-NH1	-22.11	1.01	1.32
1	A	519	ARG	CZ-NH1	16.95	1.56	1.32
1	A	470	LYS	CG-CD	14.34	1.95	1.52
1	A	579	ARG	CD-NE	9.21	1.59	1.46
1	A	469	LYS	CD-CE	8.74	1.78	1.52
1	A	243	MSE	SE-CE	-5.17	1.79	1.95
1	A	459	GLU	CG-CD	5.14	1.64	1.52
1	A	46	ARG	CG-CD	5.08	1.67	1.52

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	570	GLU	OE1-CD-OE2	-19.31	76.56	122.90
1	A	482	ASP	CA-CB-CG	-14.58	98.02	112.60
1	A	462	GLN	OE1-CD-NE2	-13.84	108.76	122.60
1	A	470	LYS	CB-CG-CD	-11.16	85.63	111.30
1	A	519	ARG	NE-CZ-NH1	-8.46	113.04	121.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	579	ARG	CG-CD-NE	6.39	126.06	112.00
1	A	504	GLY	N-CA-C	5.96	120.36	111.18
1	A	459	GLU	CG-CD-OE1	5.87	131.90	118.40
1	A	459	GLU	CG-CD-OE2	-5.74	105.19	118.40
1	A	519	ARG	NH1-CZ-NH2	5.60	126.58	119.30
1	A	459	GLU	CB-CG-CD	-5.40	103.41	112.60
1	A	466	GLU	CB-CG-CD	5.39	121.77	112.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	462	GLN	Sidechain
1	A	512[A]	ARG	Sidechain
1	A	570	GLU	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4865	0	4599	47	0
2	B	23	0	21	0	0
2	C	23	0	21	2	0
3	A	8	0	12	4	0
4	A	24	0	24	0	0
5	A	4	0	6	1	0
6	A	1	0	0	0	0
7	A	856	0	0	10	0
All	All	5804	0	4683	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:239:SER:HB2	7:A:823:HOH:O	1.47	1.10
1:A:204:ARG:HG2	7:A:1162:HOH:O	1.58	1.03
1:A:296[B]:ILE:HD13	1:A:305:LEU:HD21	1.44	0.97
1:A:416[A]:GLY:O	3:A:701:TRS:O2	1.96	0.82
1:A:239:SER:CB	7:A:823:HOH:O	2.13	0.81
1:A:388:GLU:H	1:A:391:ASN:HD22	1.28	0.77
1:A:395:CYS:SG	5:A:704:BME:S2	2.56	0.76
1:A:456:HIS:H	1:A:487:GLN:HE22	1.37	0.72
1:A:296[B]:ILE:CD1	1:A:305:LEU:HD21	2.20	0.69
1:A:547:ARG:HE	1:A:554[B]:GLU:CD	2.06	0.64
1:A:314:PRO:HG2	2:C:2:GLC:H2	1.82	0.61
1:A:388:GLU:H	1:A:391:ASN:ND2	1.96	0.61
1:A:236:ARG:NH1	7:A:811:HOH:O	2.38	0.56
1:A:397:GLN:NE2	1:A:411:LEU:H	2.04	0.55
1:A:421:GLU:OE2	3:A:701:TRS:H22	2.05	0.55
1:A:78:ARG:CZ	1:A:97:PRO:HG2	2.38	0.54
1:A:549:VAL:HG22	1:A:554[B]:GLU:HG3	1.91	0.52
1:A:109:PHE:HD1	1:A:196[A]:MSE:HE1	1.74	0.52
1:A:217:HIS:CD2	1:A:241:TRP:HD1	2.28	0.52
1:A:236:ARG:NH2	7:A:810:HOH:O	2.32	0.51
1:A:264:LEU:HD12	1:A:296[B]:ILE:HG12	1.92	0.51
1:A:103:GLU:OE1	7:A:805:HOH:O	2.19	0.51
1:A:516:ASP:OD1	1:A:519:ARG:NH2	2.45	0.49
1:A:490:GLN:NE2	1:A:494[A]:ASN:OD1	2.43	0.49
1:A:297[A]:HIS:HE1	7:A:1132:HOH:O	1.96	0.48
1:A:397:GLN:HE22	1:A:411:LEU:H	1.61	0.48
1:A:153:LEU:HD13	1:A:161[B]:ILE:HD12	1.96	0.47
1:A:276:ALA:HB1	1:A:279:TYR:HD2	1.79	0.47
1:A:354[A]:GLU:OE2	7:A:806:HOH:O	2.20	0.46
1:A:159:THR:HB	7:A:870:HOH:O	2.16	0.46
1:A:370:PHE:CZ	1:A:377:GLU:HB3	2.51	0.45
1:A:563:GLN:NE2	3:A:701:TRS:H32	2.31	0.45
1:A:581[B]:LEU:CD2	1:A:599:VAL:CG2	2.94	0.44
1:A:338:LEU:O	3:A:701:TRS:H21	2.18	0.43
1:A:75:PRO:HG3	1:A:108:ASP:HB2	2.00	0.43
1:A:167:ALA:HB3	7:A:894:HOH:O	2.19	0.42
1:A:388:GLU:N	1:A:391:ASN:HD22	2.08	0.42
1:A:412:HIS:H	1:A:412:HIS:CD2	2.37	0.42
1:A:327[B]:THR:HG21	1:A:392:PHE:HD1	1.85	0.42
1:A:430:LEU:HD21	1:A:438:LEU:HD21	2.01	0.42
1:A:581[B]:LEU:HD23	1:A:599:VAL:CG2	2.49	0.41
1:A:220:PRO:HG2	1:A:460:LEU:HD21	2.01	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:309:ASP:CG	1:A:316:LEU:HD11	2.45	0.41
1:A:277:THR:N	1:A:278:PRO:CD	2.83	0.41
1:A:314:PRO:HG2	2:C:2:GLC:C2	2.49	0.41
1:A:371:TRP:O	1:A:377:GLU:HA	2.21	0.41
1:A:80:LEU:HD12	1:A:80:LEU:C	2.46	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	607/602 (101%)	602 (99%)	5 (1%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	497/475 (105%)	497 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	382	HIS
1	A	391	ASN
1	A	397	GLN
1	A	412	HIS
1	A	487	GLN
1	A	539	HIS
1	A	563	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	B	1	2	11,11,12	1.63	2 (18%)	15,15,17	1.09	1 (6%)
2	GLC	B	2	2	12,12,12	1.08	1 (8%)	17,17,17	0.43	0
2	GLC	C	1	2	11,11,12	0.83	0	15,15,17	0.94	1 (6%)
2	GLC	C	2	2	12,12,12	1.00	0	17,17,17	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	0/2/19/22	0/1/1/1
2	GLC	B	2	2	-	0/2/22/22	0/1/1/1
2	GLC	C	1	2	-	2/2/19/22	0/1/1/1
2	GLC	C	2	2	-	2/2/22/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	GLC	C2-C3	3.93	1.58	1.52
2	B	1	GLC	O5-C5	2.30	1.48	1.43
2	B	2	GLC	O1-C1	2.15	1.46	1.39

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	GLC	O5-C5-C6	2.65	111.36	107.20
2	B	1	GLC	C1-O5-C5	2.18	115.14	112.19

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	GLC	O5-C5-C6-O6
2	C	1	GLC	C4-C5-C6-O6
2	C	1	GLC	O5-C5-C6-O6
2	C	2	GLC	C4-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2	GLC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	BME	A	704	-	3,3,3	0.26	0	1,2,2	1.56	0
3	TRS	A	701	-	7,7,7	0.52	0	9,9,9	1.11	0
4	GLC	A	702	-	12,12,12	0.53	0	17,17,17	1.65	4 (23%)
4	GLC	A	703	-	12,12,12	0.71	0	17,17,17	1.24	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BME	A	704	-	-	0/1/1/1	-
3	TRS	A	701	-	-	5/9/9/9	-
4	GLC	A	702	-	1/1/5/5	0/2/22/22	0/1/1/1
4	GLC	A	703	-	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	702	GLC	C1-C2-C3	-3.76	102.51	110.31
4	A	703	GLC	O4-C4-C5	2.84	116.36	109.30
4	A	702	GLC	O5-C1-C2	-2.41	105.98	110.28
4	A	702	GLC	O5-C5-C4	2.24	113.76	109.69
4	A	703	GLC	C4-C3-C2	2.12	114.53	110.82
4	A	702	GLC	C1-O5-C5	-2.01	109.88	113.66

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	702	GLC	C1

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	701	TRS	C1-C-C3-O3
3	A	701	TRS	C2-C-C3-O3
3	A	701	TRS	N-C-C3-O3
3	A	701	TRS	N-C-C2-O2
3	A	701	TRS	C1-C-C2-O2

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	704	BME	1	0
3	A	701	TRS	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	572/602 (95%)	-0.69	3 (0%) 87 91	3, 11, 21, 38	64 (11%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	241	TRP	3.3
1	A	14	SER	2.6
1	A	240	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	C	1	11/12	0.86	0.08	29,30,31,31	11
2	GLC	C	2	12/12	0.90	0.10	14,28,30,33	12
2	GLC	B	1	11/12	0.97	0.04	7,9,10,10	0
2	GLC	B	2	12/12	0.97	0.04	10,14,19,19	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain B:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GLC	A	702	12/12	0.72	0.21	17,22,24,25	12
3	TRS	A	701	8/8	0.76	0.15	31,34,35,38	2
4	GLC	A	703	12/12	0.89	0.09	13,19,21,21	12
5	BME	A	704	4/4	0.96	0.09	15,15,17,17	2
6	MG	A	705	1/1	0.98	0.09	21,21,21,21	1

6.5 Other polymers [i](#)

There are no such residues in this entry.