



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 5, 2025 – 10:03 AM EDT

PDB ID : 2FX6 / pdb\_00002fx6  
Title : bovine trypsin complexed with 2-aminobenzamidazole  
Authors : Katz, B.A.  
Deposited on : 2006-02-03  
Resolution : 1.57 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.45.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7165 (1.60-1.56)
Clashscore	180529	1026 (1.58-1.58)
Ramachandran outliers	177936	1005 (1.58-1.58)
Sidechain outliers	177891	1004 (1.58-1.58)
RSRZ outliers	164620	7163 (1.60-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1829 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Trypsin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	223	1672	1042	283	333	14	0	13	0

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is 1H-benzimidazol-2-amine (CCD ID: AX7) (formula: C<sub>7</sub>H<sub>7</sub>N<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			10	7	3		


- Molecule 4 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	A	146	Total 146	O 146	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Trypsin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.71Å 63.20Å 69.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.00 – 1.57 7.00 – 1.57	Depositor EDS
% Data completeness (in resolution range)	(Not available) (7.00-1.57) 97.9 (7.00-1.57)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.52 (at 1.57Å)	Xtrriage
Refinement program	X-PLOR 3.851	Depositor
R, $R_{free}$	0.202 , 0.215 0.189 , 0.193	Depositor DCC
$R_{free}$ test set	3993 reflections (10.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.4	Xtrriage
Anisotropy	0.365	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 74.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.023 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	1829	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AX7, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.73	11/1755 (0.6%)	1.75	41/2375 (1.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	91	HIS	CD2-NE2	-7.22	1.29	1.37
1	A	40	HIS	CD2-NE2	-6.60	1.30	1.37
1	A	64	GLN	CA-C	6.10	1.59	1.52
1	A	57	HIS	CG-ND1	-5.87	1.31	1.38
1	A	40	HIS	CG-ND1	-5.25	1.32	1.38
1	A	237	TRP	N-CA	5.22	1.52	1.46
1	A	231	VAL	CA-CB	5.12	1.60	1.54
1	A	51	TRP	NE1-CE2	-5.10	1.31	1.37
1	A	57	HIS	CD2-NE2	-5.07	1.32	1.37
1	A	87[A]	LYS	N-CA	5.00	1.52	1.45
1	A	87[B]	LYS	N-CA	5.00	1.52	1.45

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	190	SER	N-CA-C	-7.94	99.07	110.59
1	A	199	VAL	N-CA-C	-7.94	93.90	107.24

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	150	SER	N-CA-C	-7.33	96.09	108.26
1	A	33	LEU	N-CA-C	-7.21	97.48	109.24
1	A	71	ASP	CA-CB-CG	-7.20	105.40	112.60
1	A	86	SER	N-CA-C	-6.91	102.85	112.45
1	A	71	ASP	N-CA-C	-6.77	96.39	110.80
1	A	181	PHE	CA-CB-CG	6.68	120.48	113.80
1	A	76	VAL	N-CA-C	-6.66	97.93	107.98
1	A	125	THR	N-CA-C	-6.63	105.48	113.97
1	A	48	ASN	CA-CB-CG	6.40	119.00	112.60
1	A	78	GLY	N-CA-C	-6.37	103.38	111.72
1	A	237	TRP	NE1-CE2-CZ2	6.32	139.57	130.10
1	A	231	VAL	N-CA-C	6.22	116.96	110.62
1	A	79	ASN	CA-CB-CG	-6.08	106.52	112.60
1	A	91	HIS	CA-CB-CG	-6.00	107.80	113.80
1	A	235	VAL	N-CA-C	5.99	116.17	110.42
1	A	210	GLN	OE1-CD-NE2	-5.91	116.69	122.60
1	A	64	GLN	N-CA-C	-5.71	99.59	108.90
1	A	237	TRP	NE1-CE2-CD2	-5.64	100.07	107.40
1	A	215	TRP	NE1-CE2-CZ2	5.58	138.47	130.10
1	A	159[A]	LYS	N-CA-C	-5.57	100.17	109.24
1	A	159[B]	LYS	N-CA-C	-5.57	100.17	109.24
1	A	162	ILE	N-CA-C	-5.51	101.72	109.21
1	A	81	GLN	N-CA-C	-5.48	99.00	108.69
1	A	115	ASN	OD1-CG-ND2	-5.46	117.14	122.60
1	A	237	TRP	CG-CD1-NE1	-5.46	103.10	110.20
1	A	100	ASN	OD1-CG-ND2	-5.39	117.21	122.60
1	A	25	ASN	OD1-CG-ND2	-5.35	117.25	122.60
1	A	100	ASN	N-CA-C	-5.21	102.56	110.28
1	A	141	TRP	CG-CD1-NE1	-5.20	103.44	110.20
1	A	123	LEU	N-CA-C	-5.19	102.66	110.24
1	A	223	ASN	OD1-CG-ND2	-5.18	117.42	122.60
1	A	34	ASN	N-CA-C	5.14	117.78	109.40
1	A	95	ASN	OD1-CG-ND2	-5.13	117.47	122.60
1	A	49	SER	N-CA-C	5.10	119.13	113.01
1	A	192	GLN	OE1-CD-NE2	-5.08	117.53	122.60
1	A	237	TRP	CD1-NE1-CE2	5.07	118.02	108.90
1	A	74	ASN	CA-CB-CG	-5.05	107.55	112.60
1	A	215	TRP	CG-CD1-NE1	-5.04	103.64	110.20
1	A	37	SER	N-CA-C	-5.01	102.90	110.06

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	117	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1672	0	1637	4	0
2	A	1	0	0	0	0
3	A	10	0	7	1	0
4	A	146	0	0	1	0
All	All	1829	0	1644	5	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (5) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31:VAL:HG12	1:A:66[A]:LEU:HD23	1.91	0.52
1:A:45:SER:OG	1:A:198:PRO:HB3	2.14	0.47
1:A:59:TYR:CD1	1:A:59:TYR:C	2.97	0.41
1:A:78:GLY:O	1:A:79:ASN:HB2	2.20	0.41
3:A:246:AX7:CAE	4:A:666:HOH:O	2.69	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	234/223 (105%)	227 (97%)	7 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/184 (107%)	190 (96%)	7 (4%)	30	6

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	VAL
1	A	117	ARG
1	A	125	THR
1	A	166[A]	SER
1	A	166[B]	SER
1	A	236[A]	SER
1	A	236[B]	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	AX7	A	246	-	11,11,11	2.13	5 (45%)	8,15,15	1.78	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AX7	A	246	-	-	-	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	246	AX7	CAC-CAE	2.76	1.42	1.36
3	A	246	AX7	CAI-NAF	-2.69	1.30	1.38
3	A	246	AX7	CAJ-NAG	-2.55	1.30	1.38
3	A	246	AX7	CAE-CAJ	-2.43	1.37	1.41
3	A	246	AX7	CAD-CAI	-2.26	1.38	1.41

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	246	AX7	CAC-CAE-CAJ	-3.41	115.42	120.09
3	A	246	AX7	CAB-CAD-CAI	-3.12	115.83	120.09

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	246	AX7	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	223/223 (100%)	0.76	17 (7%) <b>21</b> <b>23</b>	9, 17, 29, 56	29 (13%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	147	SER	5.1
1	A	149	THR	3.9
1	A	97	ASN	3.1
1	A	148	GLY	3.1
1	A	115	ASN	2.8
1	A	150	SER	2.8
1	A	136	CYS	2.7
1	A	86	SER	2.7
1	A	146	SER	2.6
1	A	151	TYR	2.5
1	A	215	TRP	2.3
1	A	118	VAL	2.2
1	A	162	ILE	2.2
1	A	222	LYS	2.2
1	A	202	SER	2.1
1	A	145	LYS	2.0
1	A	239	LYS	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	AX7	A	246	10/10	0.85	0.09	25,27,28,28	0
2	CA	A	247	1/1	0.94	0.20	27,27,27,27	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.