WORLDWIDE PDB
PROTEIN DATA BANK

Full wwPDB X-ray Structure Validation Report (i)

Aug 26, 2025 – 01:24 pm BST

PDB ID	:	2JI8 / pdb_00002ji8
Title	:	X-ray structure of Oxalyl-CoA decarboxylase in complex with Formyl- CoA
Authors	:	Berthold, C.L.; Toyota, C.G.; Moussatche, P.; Wood, M.D.; Leeper, F.; Richards, N.G.J.; Lindqvist, Y.
Deposited on	:	2007-02-26
Resolution	:	2.15 Å(reported)

The following versions of software and data (see references (i)) were used in the production of this report:

|		
MolProbity	:	4-5-2 with Phenix2.0rc1
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.45.1

1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1881 (2.16-2.16)
Clashscore	180529	2047 (2.16-2.16)
Ramachandran outliers	177936	2027 (2.16-2.16)
Sidechain outliers	177891	2026 (2.16-2.16)
RSRZ outliers	164620	1882 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol	Chain	Length	Quality of chain
1	A	568	2% 89% 9% ··
1	B	568	2% 88% 10% ··

2 Entry composition ⓘ

There are 7 unique types of molecules in this entry. The entry contains 9206 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called OXALYL-COA DECARBOXYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	559	Total	C	N	O	S	0	4	0
			4194	2641	727	789	37			
1	B	556	Total	C	N	O	S	0	4	0
			4170	2625	723	786	36			

- Molecule 2 is THIAMINE DIPHOSPHATE (CCD ID: TPP) (formula: $C_{12}H_{19}N_4O_7P_2S$).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		
2	B	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Mg 1	0	0
3	B	1	Total 1	Mg 1	0	0

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 27	C 10	N 5	O 10	P 2	0	0
4	B	1	Total 27	C 10	N 5	O 10	P 2	0	0

- Molecule 5 is S-{(9R,13S,15R)-17-[(2R,3S,4R,5R)-5-(6-AMINO-9H-PURIN-9-YL)-4-HYDR OXY-3-(PHOSPHONOOXY)TETRAHYDROFURAN-2-YL]-9,13,15-TRIHYDROXY-10,1 0-DIMETHYL-13,15-DIOXIDO-4,8-DIOXO-12,14,16-TRIOXA-3,7-DIAZA-13,15-DIPHOS PHAHEPTADEC-1-YL} THIOFORMATE (CCD ID: FYN) (formula: $C_{22}H_{36}N_7O_{17}P_3S$).

FYN

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	S	0	0
			50	22	7	17	3	1		
5	B	1	Total	C	N	O	P	S	0	0
			50	22	7	17	3	1		

- Molecule 6 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).

PGE

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			10	6	4		
6	B	1	Total	C	O	0	0
			10	6	4		

• Molecule 7 is water.

| Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	316	Total O 316 316		0	0
7	B	298	Total O 298 298		0	0

3 Residue-property plots ⓘ

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: OXALYL-COA DECARBOXYLASE

Chain A:

• Molecule 1: OXALYL-COA DECARBOXYLASE

Chain B:

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α, β, γ	127.57Å 127.57Å 152.03Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.00 – 2.15 25.00 – 2.15	Depositor EDS
% Data completeness (in resolution range)	98.8 (25.00-2.15) 98.7 (25.00-2.15)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.15 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.189 , 0.231 0.193 , 0.236	Depositor DCC
R_{free} test set	3662 reflections (3.51%)	wwPDB-VP
Wilson B-factor (Å^2)	22.6	Xtriage
Anisotropy	1.120	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 44.0	EDS
L-test for twinning[2]	$< |L| > = 0.50, < L^2 > = 0.34$	Xtriage
Estimated twinning fraction	0.021 for -h,-k,l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	9206	wwPDB-VP
Average B, all atoms (Å^2)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.83% of the height of the origin peak. No significant pseudotranslation is detected.*

---

[1]Intensities estimated from amplitudes.
[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, TPP, FYN, PGE, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#|Z| >5	RMSZ	#|Z| >5
1	A	0.64	0/4280	0.84	2/5793 (0.0%)
1	B	0.63	0/4256	0.85	1/5762 (0.0%)
All	All	0.63	0/8536	0.85	3/11555 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	451	GLY	N-CA-C	-6.62	104.31	111.93
1	B	451	GLY	N-CA-C	-6.43	104.31	112.54
1	A	481	GLY	N-CA-C	5.35	118.28	110.63

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4194	0	4246	32	0
1	B	4170	0	4215	39	0
2	A	26	0	16	0	0
2	B	26	0	16	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	27	0	12	0	0
4	B	27	0	12	1	0
5	A	50	0	32	1	0
5	B	50	0	32	2	0
6	A	10	0	14	0	0
6	B	10	0	14	1	0
7	A	316	0	0	2	0
7	B	298	0	0	1	0
All	All	9206	0	8609	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:153:VAL:O	1:B:157:VAL:HG22	1.89	0.73
1:B:288:GLN:HG2	1:B:294:THR:HG21	1.72	0.71
1:A:14:HIS:NE2	1:A:42:MET:HE2	2.09	0.68
1:B:376:ASN:HD22	1:B:376:ASN:C	2.03	0.67
1:B:113:VAL:HG22	1:B:122:GLU:HG3	1.79	0.64
1:B:14:HIS:CE1	1:B:42:MET:CE	2.81	0.63
1:A:153:VAL:O	1:A:157:VAL:HG22	1.99	0.63
1:B:81:ALA:HB3	1:B:82:PRO:CD	2.30	0.61
1:B:14:HIS:CE1	1:B:42:MET:HE1	2.35	0.61
1:B:299:LEU:HD13	1:B:316:GLN:NE2	2.15	0.61
1:A:278:LEU:CD1	1:A:283:LEU:HD13	2.34	0.57
1:B:257:ASN:C	1:B:257:ASN:HD22	2.12	0.57
1:A:556:ILE:HD11	5:A:1569:FYN:S	2.44	0.57
1:A:283:LEU:O	1:A:289:HIS:HA	2.04	0.57
1:B:376:ASN:HD21	1:B:379:ASN:H	1.53	0.57
1:A:81:ALA:HB3	1:A:82:PRO:CD	2.36	0.55
1:B:556:ILE:HD11	5:B:1566:FYN:S	2.48	0.54
1:B:140:ASN:ND2	7:B:2110:HOH:O	2.40	0.54
1:B:81:ALA:CB	1:B:82:PRO:CD	2.87	0.52
1:A:112:ILE:HG23	1:A:117:GLN:HB2	1.90	0.52
1:B:430:ILE:HD12	1:B:430:ILE:C	2.36	0.51
1:A:56:GLU:OE2	2:B:1563:TPP:N1'	2.45	0.50
1:B:63:ALA:O	1:B:74:GLY:HA3	2.11	0.50
1:B:430:ILE:HD12	1:B:431:GLY:N	2.26	0.50
1:A:104:LEU:HD23	1:A:166:VAL:HB	1.93	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:278:LEU:HD11	1:A:283:LEU:HD13	1.93	0.49
1:B:326:ILE:HG13	4:B:1565:ADP:C2	2.46	0.49
1:A:104:LEU:HD22	1:A:168:LEU:HD11	1.95	0.49
1:B:209:LEU:HD21	1:B:301:LYS:HD3	1.95	0.49
1:B:546:ASP:HB3	1:B:549:ALA:HB2	1.92	0.49
1:A:288:GLN:OE1	1:A:294:THR:HG21	2.12	0.49
1:A:81:ALA:HB3	1:A:82:PRO:HD3	1.95	0.48
1:B:81:ALA:HB3	1:B:82:PRO:HD3	1.95	0.48
1:B:540:LEU:C	1:B:540:LEU:HD23	2.39	0.47
1:B:288:GLN:HG2	1:B:294:THR:CG2	2.41	0.47
1:A:483:TYR:CE1	1:A:553:SER:HA	2.49	0.47
1:B:376:ASN:ND2	1:B:379:ASN:H	2.12	0.47
1:A:278:LEU:HD13	1:A:283:LEU:HD13	1.97	0.47
1:A:508:MET:HE3	1:B:508:MET:HE3	1.97	0.46
1:B:264:ALA:HB3	1:B:355:VAL:HG13	1.96	0.46
1:A:546:ASP:HB3	1:A:549:ALA:HB2	1.96	0.46
1:A:278:LEU:HD11	1:A:283:LEU:CD1	2.46	0.45
1:A:432:MET:O	1:A:436:VAL:HG23	2.17	0.45
1:A:224:ALA:HA	1:A:326:ILE:HD13	1.98	0.45
1:A:116:GLN:NE2	7:A:2059:HOH:O	2.49	0.45
1:B:475:ILE:HG22	1:B:477:MET:CE	2.46	0.45
1:A:32:VAL:HA	1:A:36:ILE:HD11	1.97	0.45
1:B:483:TYR:CE1	1:B:553:SER:HA	2.52	0.44
1:A:520:ASN:ND2	7:A:2285:HOH:O	2.51	0.44
1:A:218:ILE:HD12	1:A:237:VAL:HG22	1.99	0.44
1:A:565:VAL:O	1:A:565:VAL:HG12	2.18	0.43
1:A:234:ARG:HA	1:A:254:LEU:HD11	2.00	0.43
1:B:71:GLY:HA3	6:B:1567:PGE:H2	2.00	0.43
1:A:81:ALA:CB	1:A:82:PRO:CD	2.96	0.43
1:B:286:LEU:O	5:B:1566:FYN:HP83	2.18	0.43
1:B:209:LEU:HD11	1:B:319:ALA:HB3	1.99	0.43
1:B:196:GLN:HE21	1:B:196:GLN:HA	1.83	0.43
1:A:399:GLU:HG3	1:A:454:ALA:HB1	2.01	0.42
1:A:508:MET:CE	1:B:508:MET:HE3	2.50	0.42
1:B:427:VAL:C	1:B:428:MET:HE2	2.44	0.42
1:A:101:MET:O	1:A:163:GLY:HA2	2.20	0.42
1:B:281:ALA:HB1	1:B:287:MET:CE	2.50	0.41
1:A:90:SER:O	1:A:93:HIS:HB3	2.20	0.41
1:A:284:ASN:HB2	1:B:118:GLY:O	2.20	0.41
1:B:475:ILE:HG22	1:B:477:MET:HE2	2.02	0.41
1:B:257:ASN:C	1:B:257:ASN:ND2	2.79	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:65:ILE:HD13	1:B:434:TYR:CZ	2.56	0.41
1:B:482:ILE:HG23	2:B:1563:TPP:H62	2.02	0.41

There are no symmetry-related clashes.

5.3　Torsion angles ⓘ

5.3.1　Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	561/568 (99%)	540 (96%)	20 (4%)	1 (0%)	44	44
1	B	558/568 (98%)	541 (97%)	15 (3%)	2 (0%)	30	27
All	All	1119/1136 (98%)	1081 (97%)	35 (3%)	3 (0%)	37	34

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	81	ALA
1	B	81	ALA
1	B	556	ILE

5.3.2　Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	441/445 (99%)	430 (98%)	11 (2%)	42	45

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	438/445 (98%)	422 (96%)	16 (4%)	29	28
All	All	879/890 (99%)	852 (97%)	27 (3%)	36	36

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	MET
1	A	80	SER
1	A	88	VAL
1	A	110	ARG
1	A	147	ILE
1	A	217	VAL
1	A	283	LEU
1	A	294	THR
1	A	338	LYS
1	A	344	ASP
1	A	367	THR
1	B	80	SER
1	B	109	GLU
1	B	110	ARG
1	B	179	VAL
1	B	257	ASN
1	B	293	LYS
1	B	294	THR
1	B	331[A]	SER
1	B	331[B]	SER
1	B	338	LYS
1	B	344	ASP
1	B	354	LYS
1	B	367	THR
1	B	374	MET
1	B	376	ASN
1	B	462	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	HIS
1	A	116	GLN
1	A	125	GLN
1	A	196	GLN

Continued from previous page...

Mol	Chain	Res	Type
1	A	228	GLN
1	A	304	GLN
1	A	308	GLN
1	A	520	ASN
1	B	140	ASN
1	B	183	ASN
1	B	196	GLN
1	B	228	GLN
1	B	257	ASN
1	B	288	GLN
1	B	316	GLN
1	B	376	ASN
1	B	491	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
6	PGE	A	1570	-	9,9,9	0.54	0	8,8,8	0.30	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	#|Z| > 2	Counts	RMSZ	#|Z| > 2
6	PGE	B	1567	-	9,9,9	0.56	0	8,8,8	0.35	0
2	TPP	A	1566	3	22,27,27	1.61	5 (22%)	29,40,40	1.59	9 (31%)
5	FYN	B	1566	-	43,52,52	1.04	1 (2%)	52,77,77	1.45	7 (13%)
4	ADP	A	1568	-	24,29,29	0.97	1 (4%)	29,45,45	1.38	6 (20%)
2	TPP	B	1563	3	22,27,27	1.54	5 (22%)	29,40,40	1.64	8 (27%)
5	FYN	A	1569	-	43,52,52	1.04	1 (2%)	52,77,77	1.45	6 (11%)
4	ADP	B	1565	-	24,29,29	0.98	1 (4%)	29,45,45	1.34	5 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PGE	A	1570	-	-	4/7/7/7	-
6	PGE	B	1567	-	-	4/7/7/7	-
2	TPP	A	1566	3	-	3/16/17/17	0/2/2/2
5	FYN	B	1566	-	-	3/45/66/66	0/3/3/3
4	ADP	A	1568	-	-	2/12/32/32	0/3/3/3
2	TPP	B	1563	3	-	5/16/17/17	0/2/2/2
5	FYN	A	1569	-	-	3/45/66/66	0/3/3/3
4	ADP	B	1565	-	-	2/12/32/32	0/3/3/3

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1569	FYN	CP1-S	-5.33	1.74	1.81
5	B	1566	FYN	CP1-S	-5.18	1.75	1.81
2	A	1566	TPP	C4-N3	-4.58	1.35	1.39
2	B	1563	TPP	C4-N3	-3.81	1.36	1.39
4	A	1568	ADP	C5-C4	2.58	1.47	1.40
2	A	1566	TPP	C4'-N3'	2.54	1.38	1.35
2	A	1566	TPP	C7'-N3	-2.52	1.44	1.48
2	A	1566	TPP	C2'-N1'	2.50	1.38	1.34
2	A	1566	TPP	C2'-N3'	2.49	1.38	1.34
4	B	1565	ADP	C5-C4	2.42	1.47	1.40
2	B	1563	TPP	C4'-N3'	2.33	1.38	1.35
2	B	1563	TPP	C2'-N1'	2.28	1.38	1.34
2	B	1563	TPP	C6'-N1'	2.11	1.38	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1563	TPP	C7'-N3	-2.00	1.45	1.48

All (41) bond angle outliers are listed below:

| Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1569	FYN	N3-C2-N1	-4.99	120.89	128.68
5	B	1566	FYN	N3-C2-N1	-4.79	121.19	128.68
5	A	1569	FYN	CP1-CP2-NP1	-4.51	102.94	112.42
5	B	1566	FYN	CP1-CP2-NP1	-4.15	103.69	112.42
5	B	1566	FYN	CP4-CP5-NP2	-3.93	103.97	111.90
2	B	1563	TPP	CM4-C4-N3	3.80	127.38	122.53
4	B	1565	ADP	N3-C2-N1	-3.77	122.78	128.68
5	A	1569	FYN	CP2-NP1-CP3	3.48	129.31	122.84
4	A	1568	ADP	N3-C2-N1	-3.45	123.29	128.68
5	B	1566	FYN	CP2-NP1-CP3	3.28	128.92	122.84
5	A	1569	FYN	CP4-CP5-NP2	-3.19	105.46	111.90
2	A	1566	TPP	C6'-C5'-C4'	3.19	120.06	115.72
2	B	1563	TPP	N1'-C2'-N3'	-2.97	120.43	125.54
2	A	1566	TPP	N1'-C2'-N3'	-2.82	120.69	125.54
4	A	1568	ADP	C4-C5-N7	-2.81	106.47	109.40
2	A	1566	TPP	CM4-C4-N3	2.78	126.07	122.53
2	B	1563	TPP	CM2-C2'-N1'	2.74	120.15	117.14
2	B	1563	TPP	C6'-C5'-C4'	2.67	119.35	115.72
2	B	1563	TPP	PA-O3A-PB	-2.55	124.07	132.83
2	A	1566	TPP	O3B-PB-O3A	2.49	112.97	104.64
4	B	1565	ADP	C4-C5-N7	-2.47	106.82	109.40
5	B	1566	FYN	CP5-NP2-CP6	2.42	126.91	122.59
4	A	1568	ADP	C1'-N9-C4	-2.41	122.40	126.64
5	A	1569	FYN	CP5-NP2-CP6	2.35	126.78	122.59
4	A	1568	ADP	O3B-PB-O2B	2.28	116.33	107.64
2	A	1566	TPP	C5'-C6'-N1'	-2.27	120.03	123.82
5	B	1566	FYN	C1'-N9-C4	-2.26	122.67	126.64
2	B	1563	TPP	CM4-C4-C5	-2.25	122.67	127.60
2	A	1566	TPP	C6'-N1'-C2'	2.24	119.78	115.96
4	A	1568	ADP	C2-N1-C6	2.23	122.58	118.75
5	B	1566	FYN	C4-C5-N7	-2.20	107.10	109.40
2	A	1566	TPP	N4'-C4'-N3'	2.20	120.14	117.03
4	B	1565	ADP	PA-O3A-PB	-2.12	125.56	132.83
2	A	1566	TPP	CM2-C2'-N3'	2.11	120.45	117.15
4	B	1565	ADP	C1'-N9-C4	-2.10	122.95	126.64
4	B	1565	ADP	C2-N1-C6	2.07	122.29	118.75
2	A	1566	TPP	PA-O3A-PB	-2.07	125.73	132.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	A	1568	ADP	PA-O3A-PB	-2.06	125.76	132.83
2	B	1563	TPP	C6'-N1'-C2'	2.03	119.42	115.96
5	A	1569	FYN	C1'-N9-C4	-2.02	123.08	126.64
2	B	1563	TPP	C5'-C6'-N1'	-2.01	120.47	123.82

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1566	TPP	PA-O3A-PB-O2B
2	A	1566	TPP	PA-O3A-PB-O3B
2	B	1563	TPP	PA-O3A-PB-O3B
4	A	1568	ADP	O4'-C4'-C5'-O5'
4	B	1565	ADP	O4'-C4'-C5'-O5'
4	A	1568	ADP	C3'-C4'-C5'-O5'
4	B	1565	ADP	C3'-C4'-C5'-O5'
6	A	1570	PGE	O1-C1-C2-O2
6	B	1567	PGE	C6-C5-O3-C4
6	B	1567	PGE	O3-C5-C6-O4
6	B	1567	PGE	O2-C3-C4-O3
5	B	1566	FYN	P2-O6-P1-O12
5	A	1569	FYN	CP2-CP1-S-CM1
6	A	1570	PGE	C6-C5-O3-C4
5	A	1569	FYN	P2-O6-P1-O11
5	B	1566	FYN	P2-O6-P1-O11
2	A	1566	TPP	C4-C5-C6-C7
2	B	1563	TPP	C4-C5-C6-C7
6	B	1567	PGE	C4-C3-O2-C2
6	A	1570	PGE	C1-C2-O2-C3
5	B	1566	FYN	CP2-CP1-S-CM1
2	B	1563	TPP	PA-O3A-PB-O2B
5	A	1569	FYN	P2-O6-P1-O12
6	A	1570	PGE	O3-C5-C6-O4
2	B	1563	TPP	PA-O3A-PB-O1B
2	B	1563	TPP	C6-C7-O7-PA

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	1567	PGE	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1566	FYN	2	0
2	B	1563	TPP	2	0
5	A	1569	FYN	1	0
4	B	1565	ADP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand TPP A 1566

Bond lengths

Bond angles

Torsions

Rings

Ligand FYN B 1566

Bond lengths

Bond angles

Torsions

Rings

Ligand ADP A 1568

Bond lengths

Bond angles

Torsions

Rings

Ligand TPP B 1563

Bond lengths

Bond angles

Torsions

Rings

Ligand FYN A 1569

Bond lengths

Bond angles

Torsions

Rings

Ligand ADP B 1565

Bond lengths

Bond angles

Torsions

Rings

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	559/568 (98%)	-0.04	13 (2%)	61	65	17, 29, 37, 63	4 (0%)
1	B	556/568 (97%)	-0.01	14 (2%)	58	63	17, 29, 38, 64	4 (0%)
All	All	1115/1136 (98%)	-0.02	27 (2%)	59	64	17, 29, 38, 64	8 (0%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	561	VAL	4.5
1	A	553	SER	4.2
1	B	556	ILE	4.1
1	B	560	ASN	3.7
1	B	554	GLY	3.6
1	A	565	VAL	3.6
1	A	563	SER	3.6
1	A	562	VAL	3.5
1	A	560	ASN	3.5
1	B	561	VAL	3.5
1	B	558	SER	3.3
1	B	555	ARG	3.3
1	B	562	VAL	3.1
1	B	553	SER	3.1
1	A	558	SER	2.9
1	A	7	VAL	2.8
1	A	559	LEU	2.8
1	A	556	ILE	2.7
1	A	554	GLY	2.5
1	B	559	LEU	2.5
1	B	296	GLY	2.5
1	A	564	LYS	2.4
1	A	557	LYS	2.4
1	B	493	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	264	ALA	2.1
1	B	483	TYR	2.1
1	B	492	PRO	2.0

6.2   Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3   Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4   Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
6	PGE	B	1567	10/10	0.61	0.18	49,50,50,50	0
6	PGE	A	1570	10/10	0.71	0.16	54,55,55,56	0
5	FYN	B	1566	50/50	0.80	0.15	32,40,45,46	50
5	FYN	A	1569	50/50	0.86	0.12	32,40,42,46	50
2	TPP	B	1563	26/26	0.94	0.08	21,28,30,32	0
2	TPP	A	1566	26/26	0.95	0.08	19,27,28,29	0
4	ADP	A	1568	27/27	0.96	0.07	19,22,24,24	0
4	ADP	B	1565	27/27	0.96	0.07	20,22,26,28	0
3	MG	A	1567	1/1	0.97	0.07	26,26,26,26	0
3	MG	B	1564	1/1	0.98	0.04	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FYN B 1566:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around FYN A 1569:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around TPP B 1563:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around TPP A 1566:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around ADP A 1568:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around ADP B 1565:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

6.5 Other polymers ⓘ

There are no such residues in this entry.