



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 10, 2025 – 10:04 PM EDT

PDB ID : 2ZWG / pdb_00002zwg
Title : Crystal structure of the copper-bound tyrosinase in complex with a caddie protein from streptomyces castaneoglobisporus obtained by soaking the deoxy-form crystal in dioxygen-saturated solution for 12 hours
Authors : Matoba, Y.; Sugiyama, M.
Deposited on : 2008-12-03
Resolution : 1.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2202 (1.34-1.30)
Clashscore	180529	2378 (1.34-1.30)
Ramachandran outliers	177936	2325 (1.34-1.30)
Sidechain outliers	177891	2325 (1.34-1.30)
RSRZ outliers	164620	2199 (1.34-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	281	
2	B	134	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3252 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	276	Total	C	N	O	S	0	4	0
			2237	1407	412	414	4			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	123	SER	PHE	conflict	UNP Q83WS2
A	274	LEU	-	expression tag	UNP Q83WS2
A	275	GLU	-	expression tag	UNP Q83WS2
A	276	HIS	-	expression tag	UNP Q83WS2
A	277	HIS	-	expression tag	UNP Q83WS2
A	278	HIS	-	expression tag	UNP Q83WS2
A	279	HIS	-	expression tag	UNP Q83WS2
A	280	HIS	-	expression tag	UNP Q83WS2
A	281	HIS	-	expression tag	UNP Q83WS2

- Molecule 2 is a protein called MelC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	72	Total	C	N	O	S	0	4	0
			568	357	105	105	1			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	60	ARG	GLY	conflict	UNP Q83WS1
B	62	ALA	GLY	conflict	UNP Q83WS1
B	127	LEU	-	expression tag	UNP Q83WS1
B	128	GLU	-	expression tag	UNP Q83WS1
B	129	HIS	-	expression tag	UNP Q83WS1
B	130	HIS	-	expression tag	UNP Q83WS1
B	131	HIS	-	expression tag	UNP Q83WS1

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	132	HIS	-	expression tag	UNP Q83WS1
B	133	HIS	-	expression tag	UNP Q83WS1
B	134	HIS	-	expression tag	UNP Q83WS1

- Molecule 3 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Cu	0	1
			4	4		
3	B	1	Total	Cu	0	1
			2	2		

- Molecule 4 is NITRATE ION (CCD ID: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	N	O	0	0
			4	1	3		
4	A	1	Total	N	O	0	0
			4	1	3		
4	A	1	Total	N	O	0	0
			4	1	3		
4	B	1	Total	N	O	0	0
			4	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	307	Total 307	O 307	0	0
5	B	118	Total 118	O 118	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tyrosinase



• Molecule 2: MelC



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	65.24Å 97.76Å 55.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.32 30.00 – 1.32	Depositor EDS
% Data completeness (in resolution range)	90.5 (30.00-1.32) 90.3 (30.00-1.32)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 1.32Å)	Xtriage
Refinement program	CNS, SHELXL-97	Depositor
R, R_{free}	0.179 , 0.225 0.168 , 0.211	Depositor DCC
R_{free} test set	3820 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	10.3	Xtriage
Anisotropy	0.234	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3252	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: DAH, NO3, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	0/2325	1.53	23/3173 (0.7%)
2	B	0.70	0/585	1.51	4/794 (0.5%)
All	All	0.75	0/2910	1.53	27/3967 (0.7%)

There are no bond length outliers.

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	153	ARG	CD-NE-CZ	12.74	142.24	124.40
2	B	105	ARG	CD-NE-CZ	11.82	140.94	124.40
1	A	136	ARG	NH1-CZ-NH2	7.99	129.69	119.30
2	B	105	ARG	NE-CZ-NH1	-7.57	113.93	121.50
1	A	136	ARG	NE-CZ-NH1	-7.49	114.01	121.50
1	A	216	HIS	CA-CB-CG	-7.17	106.63	113.80
1	A	13	ASP	CA-CB-CG	-6.93	105.67	112.60
1	A	109	ARG	CA-C-N	6.57	129.41	120.54
1	A	109	ARG	C-N-CA	6.57	129.41	120.54
1	A	45	ASP	CA-CB-CG	6.48	119.08	112.60
1	A	41	PHE	CA-CB-CG	-6.30	107.50	113.80
1	A	258	ARG	NE-CZ-NH1	-6.21	115.29	121.50
2	B	85	ARG	CD-NE-CZ	6.01	132.82	124.40
1	A	76	SER	O-C-N	-5.84	115.08	122.27
1	A	242	PRO	CA-C-O	-5.83	114.69	121.34
1	A	273	ALA	O-C-N	5.70	130.45	123.44
1	A	18	PHE	O-C-N	5.45	127.97	122.09
2	B	109	ARG	NE-CZ-NH1	-5.44	116.06	121.50
1	A	229	ARG	NE-CZ-NH2	-5.32	114.41	119.20
1	A	150	LEU	CA-C-O	-5.32	115.02	120.87
1	A	271	PHE	CA-CB-CG	-5.26	108.54	113.80
1	A	178	ARG	NE-CZ-NH2	-5.20	114.52	119.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	40	GLU	CB-CG-CD	5.06	121.21	112.60
1	A	91	ASP	CA-CB-CG	-5.06	107.54	112.60
1	A	135	SER	CA-C-N	-5.05	114.28	121.50
1	A	135	SER	C-N-CA	-5.05	114.28	121.50
1	A	145	GLY	O-C-N	-5.04	117.30	122.19

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2237	0	2113	14	0
2	B	568	0	543	6	0
3	A	4	0	0	0	0
3	B	2	0	0	0	0
4	A	12	0	0	0	0
4	B	4	0	0	0	0
5	A	307	0	0	4	0
5	B	118	0	0	1	0
All	All	3252	0	2656	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:53[A]:ARG:HD3	5:B:787:HOH:O	1.85	0.74
1:A:43:MET:HE3	2:B:82[A]:HIS:CG	2.37	0.60
1:A:43:MET:HE3	2:B:82[B]:HIS:CG	2.39	0.58
1:A:157:GLU:HB2	5:A:918:HOH:O	2.08	0.54
1:A:47:ASP:OD1	1:A:171:ASN:HB2	2.08	0.53
1:A:132:ARG:HG3	1:A:134:ASP:O	2.12	0.50
1:A:228:ARG:O	1:A:277:HIS:HB2	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:SER:O	1:A:178:ARG:HD3	2.15	0.47
1:A:157:GLU:OE1	1:A:229:ARG:NH2	2.49	0.46
2:B:82[B]:HIS:CE1	2:B:84:MET:HG3	2.50	0.46
1:A:140:ARG:NH2	5:A:874:HOH:O	2.48	0.46
1:A:153:ARG:NH1	5:A:710:HOH:O	2.49	0.46
1:A:94:VAL:O	1:A:94:VAL:HG12	2.16	0.44
1:A:228:ARG:NH1	1:A:275:GLU:O	2.50	0.44
2:B:94:VAL:HA	2:B:97:HIS:HD2	1.83	0.44
2:B:81:LEU:HB2	2:B:120:LEU:HD11	2.00	0.43
1:A:36:ARG:O	1:A:40:GLU:HG3	2.19	0.43
1:A:153:ARG:NH1	5:A:700:HOH:O	2.50	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	278/281 (99%)	267 (96%)	11 (4%)	0	100	100
2	B	71/134 (53%)	69 (97%)	2 (3%)	0	100	100
All	All	349/415 (84%)	336 (96%)	13 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/240 (100%)	238 (100%)	1 (0%)	89	74
2	B	59/94 (63%)	56 (95%)	3 (5%)	20	1
All	All	298/334 (89%)	294 (99%)	4 (1%)	73	31

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	275	GLU
2	B	53[A]	ARG
2	B	53[B]	ARG
2	B	92	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	80	GLN
2	B	116	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	DAH	B	98	2,3	12,13,14	0.90	0	12,17,19	1.57	4 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DAH	B	98	2,3	-	1/5/6/8	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	98	DAH	OE2-CE2-CD2	2.95	127.41	119.45
2	B	98	DAH	CD1-CE1-CZ	-2.85	117.64	120.50
2	B	98	DAH	CE1-CZ-CE2	2.20	122.01	119.66
2	B	98	DAH	OE2-CE2-CZ	-2.08	112.88	118.42

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	98	DAH	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NO3	A	508	-	1,3,3	0.64	0	0,3,3	-	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NO3	A	505	-	1,3,3	0.32	0	0,3,3	-	-
4	NO3	A	507	-	1,3,3	0.51	0	0,3,3	-	-
4	NO3	B	506	-	1,3,3	0.72	0	0,3,3	-	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	276/281 (98%)	0.24	17 (6%) 28 32	7, 13, 26, 60	4 (1%)
2	B	71/134 (52%)	0.37	6 (8%) 18 19	8, 13, 24, 52	4 (5%)
All	All	347/415 (83%)	0.27	23 (6%) 26 29	7, 13, 26, 60	8 (2%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	40	ALA	5.7
1	A	277	HIS	5.5
2	B	122	PRO	5.5
2	B	121	LEU	4.2
1	A	276	HIS	3.6
1	A	48	SER	3.3
1	A	94	VAL	3.2
1	A	2	THR	3.1
1	A	274	LEU	2.9
1	A	135	SER	2.6
2	B	71	GLY	2.6
1	A	229	ARG	2.5
2	B	59	ALA	2.5
1	A	133	VAL	2.5
1	A	96	ALA	2.4
1	A	110	SER	2.4
2	B	41	ALA	2.4
1	A	90	ALA	2.3
1	A	148	ALA	2.2
1	A	187	VAL	2.2
1	A	95	ARG	2.2
1	A	240	GLY	2.1
1	A	275	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DAH	B	98	13/14	0.98	0.05	9,9,13,15	1

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NO3	A	508	4/4	0.88	0.09	22,24,28,30	0
4	NO3	B	506	4/4	0.93	0.15	17,21,22,25	0
4	NO3	A	505	4/4	0.94	0.11	14,19,21,26	0
4	NO3	A	507	4/4	0.94	0.08	17,24,29,29	0
3	CU	A	504	1/1	0.98	0.15	37,37,37,37	1
3	CU	B	503[A]	1/1	0.99	0.03	14,14,14,14	1
3	CU	B	503[B]	1/1	0.99	0.03	24,24,24,24	1
3	CU	A	502	1/1	0.99	0.03	15,15,15,15	0
3	CU	A	501[A]	1/1	1.00	0.03	11,11,11,11	1
3	CU	A	501[B]	1/1	1.00	0.03	9,9,9,9	1

6.5 Other polymers [i](#)

There are no such residues in this entry.