Full wwPDB X-ray Structure Validation Report (i)

Oct 26, 2026 – 03:14 pm GMT

PDB ID	:	30PO / pdb_000030po
Title	:	Apo Structure of a TrkB Agonist
Authors	:	Briggs, D.C.; McDonald, N.Q.
Deposited on	:	2026-05-07
Resolution	:	1.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1220 (1.88-1.88)
Clashscore	190562	1234 (1.88-1.88)
Ramachandran outliers	187476	1222 (1.88-1.88)
Sidechain outliers	187428	1222 (1.88-1.88)
RSRZ outliers	180081	1220 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	248	5% 83% 9% 8%
1	B	248	5% 88% 5% 7%

2 Entry composition


There are 6 unique types of molecules in this entry. The entry contains 7259 atoms, of which 3456 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ZEB85.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	229	Total	C	H	N	O	S	0	3	0
			3503	1116	1718	315	349	5			
1	B	230	Total	C	H	N	O	S	0	0	0
			3455	1102	1690	309	349	5			

- Molecule 2 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	S	0	0
			32	8	17	2	4	1		
2	B	1	Total	C	H	N	O	S	0	0
			32	8	17	2	4	1		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Ca 1 1	0	0
3	B	1	Total Ca 1 1	0	0

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C H O 14 3 8 3	0	0
4	B	1	Total C H O 12 3 6 3	0	0

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	2	Total Cl 2 2	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	122	Total O 122 122	0	0
6	B	85	Total O 85 85	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ZEB85

Chain A:



- Molecule 1: ZEB85

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants a, b, c, α, β, γ	110.56Å 110.56Å 184.30Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	52.95 – 1.88 52.95 – 1.88	Depositor EDS
% Data completeness (in resolution range)	99.9 (52.95-1.88) 100.0 (52.95-1.88)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	0.96 (at 1.88Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.193 , 0.232 0.190 , 0.227	Depositor DCC
R_{free} test set	2669 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å^2)	48.4	Xtriage
Anisotropy	0.129	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.41 , 47.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	7259	wwPDB-VP
Average B, all atoms (Å^2)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.75% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, EPE, CA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.52	0/1840	0.63	0/2504
1	B	0.50	0/1807	0.63	0/2462
All	All	0.51	0/3647	0.63	0/4966

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1785	1718	1704	19	0
1	B	1765	1690	1690	12	0
2	A	15	17	17	1	0
2	B	15	17	17	4	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	B	12	14	14	2	0
5	B	2	0	0	0	0
6	A	122	0	0	9	2
6	B	85	0	0	2	1
All	All	3803	3456	3442	31	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:51:ASP:OD1	6:A:401:HOH:O	1.95	0.85
1:A:143:SER:O	6:A:402:HOH:O	2.06	0.74
1:A:183:ALA:O	6:A:404:HOH:O	2.13	0.66
1:A:152:SER:OG	6:A:403:HOH:O	2.12	0.65
1:A:131:PHE:CE2	1:A:249:LEU:HD22	2.36	0.60
1:B:63:ILE:HD11	1:B:131:PHE:CE2	2.38	0.59
1:A:226:THR:HG23	6:A:482:HOH:O	2.02	0.59
1:B:126:GLY:HA3	2:B:302:EPE:O3S	2.02	0.58
1:A:170[A]:ARG:NH2	6:A:406:HOH:O	2.33	0.55
1:A:63:ILE:HD11	1:A:131:PHE:CE2	2.42	0.55
1:A:152:SER:CB	6:A:403:HOH:O	2.56	0.53
1:B:226:THR:HG23	6:B:419:HOH:O	2.09	0.52
4:B:301:GOL:O3	6:B:401:HOH:O	2.17	0.51
1:A:131:PHE:CZ	1:A:249:LEU:HD22	2.47	0.50
1:B:129:TYR:H	2:B:302:EPE:H102	1.76	0.50
1:A:96:THR:HG22	1:A:98:ILE:HG13	1.96	0.48
1:B:61:ASN:HB2	1:B:124:VAL:CG2	2.43	0.48
1:B:59:ALA:HB3	2:B:302:EPE:O2S	2.14	0.48
1:A:61:ASN:HB2	1:A:124:VAL:CG2	2.44	0.48
1:A:109:HIS:CD2	6:A:423:HOH:O	2.66	0.47
1:A:202:ASN:HB2	1:B:201:LYS:HB3	1.96	0.47
1:B:178:LEU:O	4:B:303:GOL:O3	2.34	0.46
1:B:153:SER:O	1:B:153:SER:OG	2.32	0.45
1:B:129:TYR:HD2	1:B:249:LEU:HD11	1.82	0.45
1:B:76:ARG:NE	2:B:302:EPE:O2S	2.50	0.44
1:A:197:VAL:HG12	1:A:208:ILE:HD12	2.00	0.43
1:A:202:ASN:O	1:A:204:ARG:N	2.52	0.42
1:A:65:GLN:OE1	6:A:405:HOH:O	2.22	0.42
1:A:59:ALA:HB3	2:A:301:EPE:O1S	2.20	0.42
1:B:63:ILE:HD11	1:B:131:PHE:CD2	2.55	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:446:HOH:O	6:B:446:HOH:O[12_554]	1.52	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:505:HOH:O	6:A:514:HOH:O[4_455]	1.80	0.40
6:A:519:HOH:O	6:A:521:HOH:O[4_455]	2.05	0.15

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	228/248 (92%)	217 (95%)	11 (5%)	0	100	100
1	B	226/248 (91%)	220 (97%)	6 (3%)	0	100	100
All	All	454/496 (92%)	437 (96%)	17 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/205 (96%)	196 (100%)	1 (0%)	81	76
1	B	194/205 (95%)	194 (100%)	0	100	100
All	All	391/410 (95%)	390 (100%)	1 (0%)	86	82

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	247	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	105	GLN
1	B	105	GLN
1	B	132	HIS
1	B	248	HIS

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	GOL	B	303	3	5,5,5	0.76	0	5,5,5	0.82	0
4	GOL	B	301	-	5,5,5	1.13	0	5,5,5	0.63	0
2	EPE	B	302	-	15,15,15	0.99	1 (6%)	18,20,20	2.28	7 (38%)
2	EPE	A	301	-	15,15,15	1.44	1 (6%)	18,20,20	1.61	4 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	303	3	-	1/4/4/4	-
4	GOL	B	301	-	-	4/4/4/4	-
2	EPE	B	302	-	-	7/9/19/19	0/1/1/1
2	EPE	A	301	-	-	2/9/19/19	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	EPE	C10-S	5.09	1.84	1.77
2	B	302	EPE	C10-S	3.44	1.82	1.77

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	302	EPE	O1S-S-C10	4.40	112.22	106.92
2	B	302	EPE	O2S-S-C10	4.06	111.81	106.92
2	B	302	EPE	C7-N4-C3	3.61	120.47	111.23
2	A	301	EPE	C7-N4-C3	3.48	120.13	111.23
2	B	302	EPE	C5-N4-C3	3.28	116.22	108.83
2	B	302	EPE	C7-N4-C5	3.20	119.43	111.23
2	B	302	EPE	O3S-S-C10	3.02	110.66	105.77
2	A	301	EPE	C5-N4-C3	2.74	114.99	108.83
2	B	302	EPE	O2S-S-O1S	-2.34	105.84	113.95
2	A	301	EPE	O3S-S-C10	2.28	109.46	105.77
2	A	301	EPE	O2S-S-C10	-2.09	104.40	106.92

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	302	EPE	C10-C9-N1-C2
4	B	301	GOL	C1-C2-C3-O3
4	B	301	GOL	O1-C1-C2-O2
2	B	302	EPE	C9-C10-S-O3S
4	B	301	GOL	O1-C1-C2-C3
4	B	301	GOL	O2-C2-C3-O3
2	B	302	EPE	S-C10-C9-N1
2	A	301	EPE	C10-C9-N1-C6
2	B	302	EPE	C10-C9-N1-C6
2	B	302	EPE	C9-C10-S-O1S

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	302	EPE	C9-C10-S-O2S
2	B	302	EPE	N4-C7-C8-O8
4	B	303	GOL	C1-C2-C3-O3
2	A	301	EPE	C10-C9-N1-C2

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	303	GOL	1	0
4	B	301	GOL	1	0
2	B	302	EPE	4	0
2	A	301	EPE	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	229/248 (92%)	0.27	12 (5%)	33	34	34, 61, 102, 177	2 (0%)
1	B	230/248 (92%)	0.39	12 (5%)	33	34	48, 65, 103, 128	0
All	All	459/496 (92%)	0.33	24 (5%)	33	34	34, 63, 103, 177	2 (0%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	261	GLY	5.2
1	B	199	ALA	4.0
1	A	201	LYS	3.9
1	A	200	GLY	3.8
1	B	144	SER	3.6
1	B	128	TYR	3.4
1	B	200	GLY	3.4
1	A	202	ASN	3.4
1	A	199	ALA	3.3
1	A	249	LEU	3.2
1	A	128	TYR	3.2
1	B	261	GLY	3.2
1	B	153	SER	3.0
1	A	153	SER	3.0
1	B	152	SER	2.9
1	B	202	ASN	2.9
1	B	247	HIS	2.9
1	B	249	LEU	2.9
1	A	67	PRO	2.4
1	B	67	PRO	2.3
1	A	152	SER	2.3
1	A	69	ARG	2.1
1	A	143	SER	2.1
1	B	81	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	EPE	B	302	15/15	0.84	0.32	48,76,92,100	32
3	CA	A	302	1/1	0.84	0.09	98,98,98,98	0
4	GOL	B	301	6/6	0.86	0.13	56,81,102,123	0
4	GOL	B	303	6/6	0.89	0.13	61,80,95,105	0
2	EPE	A	301	15/15	0.94	0.16	51,73,94,95	32
5	CL	B	304	1/1	0.96	0.10	51,51,51,51	0
5	CL	B	305	1/1	0.97	0.11	67,67,67,67	0
3	CA	B	306	1/1	0.99	0.03	74,74,74,74	0

6.5 Other polymers (i)

There are no such residues in this entry.