Full wwPDB X-ray Structure Validation Report (i)

Oct 5, 2026 – 04:12 pm BST

PDB ID	:	30UF / pdb_000030uf
Title	:	X-ray structure of the adduct formed upon reaction of the organopalladium compound complex 1 with RNase A
Authors	:	Ferraro, G.; Tito, G.; Merlino, A.
Deposited on	:	2026-05-13
Resolution	:	1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.2
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	124	4% 81% 19%
1	BBB	124	16% 90% 10%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 1980 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribonuclease pancreatic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	AAA	124	947	572	170	193	12	0	0	0
1	BBB	124	947	572	170	193	12	0	0	0

- Molecule 2 is PALLADIUM ION (CCD ID: PD) (formula: Pd) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Pd		
2	AAA	1	1	1	0	0
2	BBB	1	1	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	AAA	54	54	54	0	0
3	BBB	30	30	30	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ribonuclease pancreatic

Chain AAA:

- Molecule 1: Ribonuclease pancreatic

Chain BBB:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	101.02Å 32.86Å 74.64Å 90.00° 90.51° 90.00°	Depositor
Resolution (Å)	31.25 – 1.99 31.25 – 1.99	Depositor EDS
% Data completeness (in resolution range)	71.2 (31.25-1.99) 71.2 (31.25-1.99)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.99 (at 1.98Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.211 , 0.234 0.219 , 0.242	Depositor DCC
R_{free} test set	644 reflections (3.72%)	wwPDB-VP
Wilson B-factor (Å^2)	27.0	Xtriage
Anisotropy	0.054	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.40 , 52.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	0.026 for -h,-k,l	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	1980	wwPDB-VP
Average B, all atoms (Å^2)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.06% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	AAA	1.09	1/963 (0.1%)	1.41	2/1300 (0.2%)
1	BBB	1.05	0/963	1.37	0/1300
All	All	1.07	1/1926 (0.1%)	1.39	2/2600 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	60	GLN	N-CA	5.36	1.49	1.46

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	AAA	31	LYS	CA-C-N	5.29	127.90	120.28
1	AAA	31	LYS	C-N-CA	5.29	127.90	120.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	947	0	894	11	0
1	BBB	947	0	894	6	0
2	AAA	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	BBB	1	0	0	0	0
3	AAA	54	0	0	1	0
3	BBB	30	0	0	0	0
All	All	1980	0	1788	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:111:GLU:HG2	1:AAA:118:VAL:HG21	1.75	0.69
1:BBB:48:HIS:NE2	1:BBB:82:THR:OG1	2.40	0.55
1:AAA:11:GLN:HG2	1:AAA:35:LEU:HD21	1.91	0.52
1:BBB:111:GLU:HG2	1:BBB:118:VAL:HG21	1.93	0.49
1:AAA:14:ASP:OD2	1:AAA:25:TYR:OH	2.30	0.48
1:AAA:25:TYR:CZ	1:AAA:29:MET:HG3	2.48	0.48
1:AAA:86:GLU:HG2	1:AAA:90:SER:HB3	1.96	0.48
1:AAA:20:ALA:HB2	1:AAA:82:THR:OG1	2.17	0.45
1:AAA:111:GLU:HG2	1:AAA:118:VAL:CG2	2.46	0.45
1:BBB:111:GLU:CG	1:BBB:118:VAL:HG21	2.46	0.45
1:BBB:17:THR:HG1	1:BBB:48:HIS:HD1	1.64	0.44
1:AAA:98:LYS:HE3	3:AAA:315:HOH:O	2.17	0.44
1:AAA:66:LYS:NZ	1:AAA:122:ALA:HB2	2.34	0.43
1:BBB:61:LYS:O	1:BBB:73:TYR:HA	2.19	0.43
1:AAA:92:TYR:CD1	1:AAA:93:PRO:HA	2.53	0.42
1:AAA:73:TYR:O	1:AAA:107:ILE:HA	2.19	0.42
1:BBB:92:TYR:CG	1:BBB:93:PRO:HA	2.55	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	122/124 (98%)	116 (95%)	6 (5%)	0	100	100
1	BBB	122/124 (98%)	110 (90%)	11 (9%)	1 (1%)	16	11
All	All	244/248 (98%)	226 (93%)	17 (7%)	1 (0%)	30	27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	BBB	90	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	108/109 (99%)	104 (96%)	4 (4%)	30	30
1	BBB	108/109 (99%)	106 (98%)	2 (2%)	50	56
All	All	216/218 (99%)	210 (97%)	6 (3%)	38	41

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	21	SER
1	AAA	24	ASN
1	AAA	77	SER
1	AAA	119	HIS
1	BBB	21	SER
1	BBB	124	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	AAA	124/124 (100%)	0.31	5 (4%)	42	41	17, 30, 58, 78	0
1	BBB	124/124 (100%)	0.97	20 (16%)	4	4	21, 54, 101, 120	0
All	All	248/248 (100%)	0.64	25 (10%)	12	11	17, 37, 94, 120	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BBB	97	TYR	4.0
1	BBB	95	CYS	3.6
1	BBB	46	PHE	3.3
1	BBB	96	ALA	2.8
1	BBB	29	MET	2.8
1	BBB	92	TYR	2.6
1	AAA	21	SER	2.6
1	BBB	87	THR	2.5
1	BBB	27	ASN	2.5
1	BBB	23	SER	2.4
1	BBB	25	TYR	2.4
1	BBB	89	SER	2.4
1	BBB	99	THR	2.3
1	AAA	22	SER	2.3
1	BBB	15	SER	2.3
1	AAA	32	SER	2.3
1	BBB	100	THR	2.3
1	AAA	1	LYS	2.2
1	BBB	14	ASP	2.2
1	BBB	91	LYS	2.2
1	BBB	40	CYS	2.1
1	BBB	101	GLN	2.1
1	BBB	19	ALA	2.1
1	BBB	102	ALA	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	AAA	89	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	PD	BBB	201	1/1	0.95	0.07	62,62,62,62	1
2	PD	AAA	201	1/1	0.98	0.06	37,37,37,37	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around PD BBB 201:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around PD AAA 201:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.