Full wwPDB X-ray Structure Validation Report (i)

Nov 16, 2026 – 12:05 pm GMT

PDB ID	:	31BM / pdb_000031bm
Title	:	Crystal structure of inactive ThomasA-E540A in complex with IgA1 Fab and hinge region (PVPSTPPT residues)
Authors	:	Castillo Calvente, C.; Martinez Gascuena, A.; Marquez-Monino, M.A.; Hernandez-Aleman, H.; Trastoy, B.
Deposited on	:	2026-05-22
Resolution	:	2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	560	2% 87% 12%
2	H	240	5% 80% 14% 5%
3	L	213	3% 87% 13%

2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 7896 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IgA protease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	556	Total	C	N	O	S	0	1	0
			4362	2764	725	856	17			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	319	GLY	-	expression tag	UNP Q9AES2
A	320	GLY	-	expression tag	UNP Q9AES2
A	321	SER	-	expression tag	UNP Q9AES2
A	322	GLY	-	expression tag	UNP Q9AES2
A	540	ALA	GLU	engineered mutation	UNP Q9AES2

- Molecule 2 is a protein called IgA1 Heavy Chain Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	227	Total	C	N	O	S	0	0	0
			1693	1076	282	327	8			


- Molecule 3 is a protein called IgA1 Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	213	Total	C	N	O	S	0	0	0
			1608	1000	272	329	7			

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Zn	0	0
			2	2		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	H	1	Total C O 4 2 2	0	0
5	L	1	Total C O 4 2 2	0	0
5	L	1	Total C O 4 2 2	0	0
5	L	1	Total C O 4 2 2	0	0

- Molecule 6 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	H	1	Total C O 4 2 2	0	0

- Molecule 7 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 7 4 3	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	97	Total O 97 97	0	0
8	H	49	Total O 49 49	0	0
8	L	42	Total O 42 42	0	0


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: IgA protease



• Molecule 2: IgA1 Heavy Chain Fab



• Molecule 3: IgA1 Light Chain



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	253.77Å 53.62Å 79.15Å 90.00° 92.11° 90.00°	Depositor
Resolution (Å)	39.64 – 2.40 39.64 – 2.40	Depositor EDS
% Data completeness (in resolution range)	94.2 (39.64-2.40) 94.2 (39.64-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.57 (at 2.40Å)	Xtriage
Refinement program	PHENIX (1.21_5207)	Depositor
R, R_{free}	0.227 , 0.260 0.226 , 0.258	Depositor DCC
R_{free} test set	1763 reflections (4.19%)	wwPDB-VP
Wilson B-factor (Å^2)	48.0	Xtriage
Anisotropy	0.154	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.30 , 32.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.020 for -h,-k,l	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	7896	wwPDB-VP
Average B, all atoms (Å^2)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, EDO, ACT, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.18	0/4471	0.33	0/6079
2	H	0.14	0/1743	0.31	0/2399
3	L	0.14	0/1642	0.35	0/2233
All	All	0.16	0/7856	0.33	0/10711

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	683	ARG	Sidechain
1	A	696	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4362	0	4063	39	0
2	H	1693	0	1628	21	0
3	L	1608	0	1507	16	0
4	A	2	0	0	0	0
5	A	4	0	6	0	0
5	H	4	0	6	0	0
5	L	12	0	18	0	0
6	A	12	0	9	0	0
6	H	4	0	3	1	0
7	A	7	0	10	0	0
8	A	97	0	0	0	0
8	H	49	0	0	0	0
8	L	42	0	0	0	0
All	All	7896	0	7250	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:142:VAL:HG22	2:H:190:THR:HG22	1.63	0.78
1:A:589:TYR:HB3	1:A:664:TYR:HD1	1.54	0.72
1:A:709:GLN:HE22	1:A:739:ASP:HA	1.62	0.64
1:A:709:GLN:NE2	1:A:739:ASP:HA	2.14	0.62
1:A:361:ARG:HH21	1:A:407:ARG:HG2	1.66	0.61
3:L:200:LEU:HD11	3:L:203:PRO:HA	1.85	0.58
1:A:551:GLU:HA	1:A:568:LEU:HD21	1.86	0.58
1:A:696:ARG:NH2	1:A:768:SER:OG	2.38	0.57
1:A:869:VAL:HG12	1:A:871:THR:HG22	1.86	0.57
1:A:367:MET:HE3	1:A:630:LEU:HD12	1.87	0.56
1:A:824:ILE:HD11	1:A:874:TYR:HB2	1.86	0.56
2:H:27:ASN:HD21	2:H:29:ALA:HB3	1.72	0.53
3:L:162:VAL:HG22	3:L:174:LEU:HD12	1.91	0.52
1:A:827:THR:HG21	1:A:848:LEU:HD21	1.92	0.52
1:A:443:ASN:H	6:H:301:ACT:H3	1.75	0.51
1:A:716:ILE:HG22	1:A:724:ALA:HB3	1.93	0.50
1:A:439:PRO:HD3	1:A:510:PHE:HB2	1.93	0.50
2:H:25:ILE:HD11	2:H:71:THR:HA	1.95	0.49
1:A:367:MET:HE1	1:A:545:LEU:HD21	1.94	0.49
2:H:127:VAL:HB	2:H:216:VAL:HG21	1.94	0.49
3:L:198:GLN:C	3:L:200:LEU:H	2.20	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:35:TRP:CE2	3:L:73:PHE:HB2	2.47	0.48
1:A:553:SER:O	1:A:593:ASN:ND2	2.43	0.48
1:A:758:ILE:HG22	1:A:759:LYS:HD3	1.95	0.48
2:H:125:PRO:HB3	2:H:150:PHE:HB3	1.96	0.47
1:A:473:LYS:HB2	1:A:473:LYS:HE2	1.76	0.47
2:H:100:VAL:HG12	2:H:101:THR:HG23	1.97	0.47
3:L:80:PRO:O	3:L:83:ILE:HD12	2.15	0.47
1:A:340:TRP:CE2	2:H:212:PRO:HG3	2.51	0.46
2:H:36:ARG:HB3	2:H:46:ILE:HD11	1.97	0.46
2:H:31:TYR:HB2	2:H:96:THR:HB	1.97	0.46
2:H:19:SER:HA	2:H:75:GLN:HG2	1.98	0.46
2:H:25:ILE:HG22	2:H:32:TRP:CE2	2.51	0.45
2:H:99:VAL:HG23	2:H:100:VAL:H	1.81	0.45
1:A:561:LYS:NZ	2:H:195:GLN:HA	2.32	0.45
3:L:35:TRP:CZ3	3:L:88:CYS:HB3	2.51	0.45
1:A:643:VAL:HG23	1:A:677:SER:HB3	1.98	0.45
2:H:76:PHE:CZ	2:H:93:CYS:HB2	2.51	0.44
3:L:147:TRP:CE2	3:L:178:LEU:HB2	2.53	0.44
1:A:863:ILE:HD12	1:A:868:THR:HG21	2.00	0.44
2:H:167:VAL:HA	2:H:190:THR:O	2.16	0.44
3:L:54:LEU:HD22	3:L:58:VAL:HB	1.99	0.44
1:A:326:ASN:HD21	1:A:382:GLY:H	1.66	0.44
1:A:861:LYS:HG2	1:A:862:PRO:HD2	1.98	0.44
2:H:142:VAL:HG12	3:L:117:PHE:HZ	1.81	0.44
1:A:393[A]:GLN:HG2	1:A:831:THR:O	2.18	0.43
1:A:583:LEU:O	1:A:696:ARG:NH1	2.50	0.43
2:H:174:PRO:HA	2:H:185:THR:HB	1.99	0.43
3:L:184:ASP:HA	3:L:187:LYS:HG2	2.01	0.43
1:A:352:ILE:HD13	1:A:415:THR:HG22	2.00	0.43
1:A:846:LYS:HD3	1:A:847:THR:HG23	2.01	0.42
2:H:132:LEU:HD11	2:H:142:VAL:HB	2.01	0.42
3:L:83:ILE:HD11	3:L:105:MET:HE2	2.00	0.42
1:A:624:PHE:CD1	1:A:637:TYR:HB2	2.54	0.42
1:A:553:SER:O	1:A:599:MET:HG2	2.19	0.42
2:H:37:GLN:HB2	2:H:43:LEU:HD23	2.02	0.42
1:A:844:PRO:HB3	1:A:872:TYR:CE2	2.55	0.41
2:H:130:LEU:HD11	2:H:146:LEU:HB2	2.02	0.41
1:A:389:LYS:HE3	1:A:828:ALA:HB1	2.01	0.41
1:A:558:LEU:O	1:A:559:ASP:C	2.63	0.41
1:A:565:SER:OG	1:A:568:LEU:HB2	2.20	0.41
3:L:112:PRO:HB3	3:L:138:PHE:HB3	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:200:LEU:HD21	3:L:204:VAL:H	1.84	0.41
3:L:39:ARG:HD2	3:L:84:ALA:HB2	2.03	0.41
2:H:141:VAL:HG21	2:H:196:CYS:HB3	2.02	0.41
3:L:35:TRP:HB2	3:L:48:ILE:HB	2.02	0.41
1:A:343:ASN:HB3	1:A:446:HIS:O	2.21	0.41
1:A:813:SER:O	1:A:869:VAL:HA	2.21	0.41
3:L:139:TYR:CG	3:L:140:PRO:HA	2.55	0.41
1:A:482:PRO:HG2	1:A:486:GLU:HA	2.03	0.40
1:A:814:ILE:O	1:A:829:GLY:HA3	2.20	0.40
1:A:633:ASP:OD1	1:A:633:ASP:N	2.54	0.40
1:A:675:LEU:HD23	1:A:680:SER:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	555/560 (99%)	529 (95%)	25 (4%)	1 (0%)	43 58
2	H	225/240 (94%)	214 (95%)	11 (5%)	0	100 100
3	L	211/213 (99%)	203 (96%)	8 (4%)	0	100 100
All	All	991/1013 (98%)	946 (96%)	44 (4%)	1 (0%)	48 64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	559	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	466/498 (94%)	462 (99%)	4 (1%)	70 85
2	H	188/208 (90%)	186 (99%)	2 (1%)	65 82
3	L	177/187 (95%)	176 (99%)	1 (1%)	78 89
All	All	831/893 (93%)	824 (99%)	7 (1%)	73 86

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	541	PHE
1	A	558	LEU
1	A	673	ASP
1	A	847	THR
2	H	50	TYR
2	H	99	VAL
3	L	33	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	334	GLN
1	A	446	HIS
1	A	587	ASN
1	A	709	GLN
2	H	-3	GLN
2	H	140	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	EDO	L	301	-	3,3,3	0.28	0	2,2,2	0.29	0
6	ACT	A	904	-	3,3,3	1.03	0	3,3,3	1.33	0
6	ACT	A	905	-	3,3,3	1.08	0	3,3,3	1.21	0
5	EDO	L	303	-	3,3,3	0.28	0	2,2,2	0.24	0
6	ACT	H	301	-	3,3,3	1.08	0	3,3,3	1.22	0
6	ACT	A	906	-	3,3,3	1.08	0	3,3,3	1.19	0
5	EDO	A	903	-	3,3,3	0.28	0	2,2,2	0.19	0
5	EDO	L	302	-	3,3,3	0.27	0	2,2,2	0.30	0
7	PEG	A	907	-	6,6,6	0.24	0	5,5,5	0.27	0
5	EDO	H	302	-	3,3,3	0.28	0	2,2,2	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	L	301	-	-	0/1/1/1	-
5	EDO	L	303	-	-	0/1/1/1	-
5	EDO	A	903	-	-	1/1/1/1	-
5	EDO	L	302	-	-	0/1/1/1	-
7	PEG	A	907	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	H	302	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	907	PEG	O2-C3-C4-O4
5	H	302	EDO	O1-C1-C2-O2
7	A	907	PEG	C4-C3-O2-C2
5	A	903	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	H	301	ACT	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	556/560 (99%)	0.52	14 (2%) 58 54	25, 61, 88, 98	1 (0%)
2	H	227/240 (94%)	0.32	11 (4%) 35 32	35, 47, 85, 100	0
3	L	213/213 (100%)	0.40	6 (2%) 55 51	30, 49, 78, 101	0
All	All	996/1013 (98%)	0.45	31 (3%) 51 47	25, 53, 86, 101	1 (0%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	199	GLY	3.6
2	H	198	ALA	3.4
2	H	197	LEU	3.3
2	H	135	THR	3.2
2	H	137	PRO	2.8
1	A	868	THR	2.6
2	H	196	CYS	2.6
2	H	134	SER	2.6
3	L	200	LEU	2.6
3	L	203	PRO	2.5
3	L	198	GLN	2.4
1	A	762	PHE	2.4
3	L	213	CYS	2.4
1	A	773	TYR	2.4
3	L	189	LYS	2.4
2	H	223	PRO	2.3
1	A	822	SER	2.3
2	H	138	ASP	2.3
1	A	686	THR	2.3
1	A	556	TYR	2.2
1	A	789	PHE	2.2
1	A	869	VAL	2.1
2	H	136	GLN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	554	ASN	2.1
1	A	404	TYR	2.1
1	A	635	ASP	2.1
1	A	718	HIS	2.1
1	A	560	ASP	2.0
3	L	208	PHE	2.0
1	A	710	LEU	2.0
2	H	194	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	EDO	H	302	4/4	0.65	0.19	53,58,60,61	0
6	ACT	A	906	4/4	0.68	0.19	58,66,70,72	0
5	EDO	A	903	4/4	0.70	0.20	66,68,70,71	0
5	EDO	L	301	4/4	0.71	0.20	59,63,63,64	0
5	EDO	L	302	4/4	0.75	0.15	56,59,61,62	0
6	ACT	A	904	4/4	0.77	0.22	48,48,48,50	0
6	ACT	A	905	4/4	0.80	0.16	53,57,58,60	0
6	ACT	H	301	4/4	0.84	0.19	44,51,52,52	0
7	PEG	A	907	7/7	0.87	0.15	46,48,52,56	0
5	EDO	L	303	4/4	0.89	0.11	43,43,48,50	0
4	ZN	A	901	1/1	0.98	0.04	76,76,76,76	0
4	ZN	A	902	1/1	0.99	0.02	55,55,55,55	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around ZN A 901:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around ZN A 902:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.