Full wwPDB X-ray Structure Validation Report (i)

Oct 27, 2026 – 03:08 pm GMT

PDB ID	:	32YH / pdb_000032yh
Title	:	Streptomyces sp. TLI_171 GH5_7 Mannanase, Apo
Authors	:	Bloch, Y.
Deposited on	:	2026-07-30
Resolution	:	1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	397	% 89% 7% .
1	B	397	% 88% 8% .

2 Entry composition


There are 4 unique types of molecules in this entry. The entry contains 12854 atoms, of which 5888 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GH5_7 Mannanase.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	381	Total	C	H	N	O	S	85	13	0
			6033	1953	2925	563	581	11			
1	B	381	Total	C	H	N	O	S	88	18	0
			6080	1967	2945	566	591	11			

- Molecule 2 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	O	S	0	0
			10	2	6	1	1		
2	B	1	Total	C	H	O	S	0	0
			10	2	6	1	1		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C H O 10 2 6 2	1	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	355	Total O 355 355	0	0
4	B	355	Total O 356 356	0	2


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: GH5_7 Mannanase



• Molecule 1: GH5_7 Mannanase



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α, β, γ	137.62Å 137.62Å 82.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.71 – 1.55 30.71 – 1.55	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.71-1.55) 99.9 (30.71-1.55)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.41 (at 1.54Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.143 , 0.164 0.142 , 0.161	Depositor DCC
R_{free} test set	2000 reflections (1.73%)	wwPDB-VP
Wilson B-factor (Å^2)	23.2	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.40 , 34.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.98	EDS
Total number of atoms	12854	wwPDB-VP
Average B, all atoms (Å^2)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.42% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.77	0/3217	0.76	1/4378 (0.0%)
1	B	0.78	1/3240 (0.0%)	0.77	1/4408 (0.0%)
All	All	0.78	1/6457 (0.0%)	0.77	2/8786 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	7
1	B	0	3
All	All	0	10

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	198	ARG	NE-CZ	-6.61	1.25	1.33

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	401	ASP	CA-CB-CG	7.00	119.59	112.60
1	B	401	ASP	CA-CB-CG	5.53	118.13	112.60

There are no chirality outliers.

All (10) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	184	ARG	Sidechain
1	A	188[A]	ARG	Sidechain
1	A	188[B]	ARG	Sidechain
1	A	346[A]	ARG	Sidechain
1	A	346[B]	ARG	Sidechain
1	A	359[A]	ARG	Sidechain
1	A	44	ARG	Sidechain
1	B	176	ARG	Sidechain
1	B	184	ARG	Sidechain
1	B	198	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3108	2925	2915	14	0
1	B	3135	2945	2932	24	0
2	A	4	6	6	0	0
2	B	4	6	6	0	0
3	A	4	6	6	0	0
4	A	355	0	0	2	1
4	B	356	0	0	12	0
All	All	6966	5888	5865	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:400:ASN:OD1	1:A:411[A]:GLN:NE2	1.88	1.05
1:B:421[B]:THR:HG21	4:B:664:HOH:O	1.66	0.94
1:B:421[B]:THR:HG22	4:B:720:HOH:O	1.79	0.82
1:B:424[A]:GLY:O	4:B:601:HOH:O	2.01	0.78
1:B:192[A]:GLU:OE1	4:B:602:HOH:O	2.07	0.70
1:B:328:ARG:HD3	4:B:901:HOH:O	1.96	0.66
1:B:421[B]:THR:HG23	4:B:803:HOH:O	1.97	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:310[A]:ASN:ND2	4:B:606:HOH:O	2.34	0.59
1:A:96:ALA:O	1:A:142:PRO:HD2	2.09	0.53
1:A:216[A]:GLU:OE2	1:A:336:GLU:OE1	2.28	0.51
1:B:216[B]:GLU:OE2	1:B:336:GLU:OE1	2.27	0.51
1:B:240:SER:HA	1:B:252[B]:LEU:HD21	1.92	0.50
1:B:199:ASN:HB3	1:B:202:THR:OG1	2.11	0.49
1:B:59:ARG:HG2	4:B:616:HOH:O	2.11	0.49
1:B:216[B]:GLU:HG2	1:B:253:ALA:HB3	1.93	0.49
1:B:258:GLY:HA2	1:B:278:TRP:CD1	2.47	0.49
1:B:96:ALA:O	1:B:142:PRO:HD2	2.13	0.48
1:A:258:GLY:HA2	1:A:278:TRP:CD1	2.49	0.48
1:A:240:SER:HA	1:A:252[A]:LEU:HD21	1.95	0.48
1:A:310[A]:ASN:ND2	4:A:605:HOH:O	2.46	0.48
1:B:59:ARG:CG	4:B:616:HOH:O	2.63	0.47
1:B:394:HIS:HE1	4:B:791:HOH:O	1.98	0.47
1:B:412:LEU:C	1:B:412:LEU:HD23	2.40	0.47
1:A:418:GLU:HG2	4:A:729:HOH:O	2.16	0.46
1:A:412:LEU:C	1:A:412:LEU:HD23	2.41	0.45
1:A:216[A]:GLU:HG2	1:A:253:ALA:HB3	1.98	0.45
1:B:77:TYR:CD1	1:B:77:TYR:C	2.94	0.45
1:B:268:TYR:HB3	1:B:269:PRO:HD3	1.99	0.44
1:A:199:ASN:HB3	1:A:202:THR:OG1	2.18	0.43
1:A:89:MET:HG2	1:A:398:TRP:CE3	2.53	0.43
1:A:221:PRO:HD2	1:A:256:ASP:HA	1.99	0.43
1:B:99:PHE:CE1	1:B:145:ASN:HB2	2.54	0.43
1:A:142:PRO:HA	1:A:216[B]:GLU:HB3	2.01	0.42
1:B:421[B]:THR:CG2	4:B:803:HOH:O	2.63	0.41
1:B:261:GLY:HA2	1:B:272:ASN:HA	2.03	0.41
1:A:77:TYR:CD1	1:A:77:TYR:C	2.99	0.41
1:B:216[A]:GLU:HA	1:B:253:ALA:O	2.21	0.41
1:B:310[A]:ASN:CG	4:B:606:HOH:O	2.64	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:634:HOH:O	4:A:651:HOH:O[7_556]	1.96	0.24

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/397 (99%)	378 (96%)	14 (4%)	0	100	100
1	B	396/397 (100%)	382 (96%)	14 (4%)	0	100	100
All	All	788/794 (99%)	760 (96%)	28 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/308 (100%)	304 (99%)	3 (1%)	68	47
1	B	309/308 (100%)	306 (99%)	3 (1%)	68	47
All	All	616/616 (100%)	610 (99%)	6 (1%)	76	47

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	97	TRP
1	A	232[A]	LEU
1	A	232[B]	LEU
1	B	97	TRP
1	B	232[A]	LEU
1	B	232[B]	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	105	HIS
1	A	305	ASN
1	B	394	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	DMS	B	501	-	3,3,3	0.95	0	3,3,3	0.13	0
3	EDO	A	502	-	3,3,3	0.77	0	2,2,2	0.42	0
2	DMS	A	501	-	3,3,3	1.19	1 (33%)	3,3,3	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	502	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	DMS	O-S	2.03	1.63	1.50

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	381/397 (95%)	-0.50	2 (0%)	87	90	10, 24, 38, 87	13 (3%)
1	B	381/397 (95%)	-0.56	4 (1%)	79	85	10, 23, 37, 61	18 (4%)
All	All	762/794 (95%)	-0.53	6 (0%)	82	87	10, 24, 38, 87	31 (4%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	423[A]	ALA	5.6
1	B	424[A]	GLY	4.7
1	A	424	GLY	3.0
1	A	308	GLY	2.7
1	B	167	SER	2.2
1	B	172	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	EDO	A	502	4/4	0.86	0.13	27,41,44,49	1
2	DMS	B	501	4/4	0.95	0.11	23,39,39,40	0
2	DMS	A	501	4/4	0.95	0.10	23,38,41,41	0

6.5 Other polymers (i)

There are no such residues in this entry.