Full wwPDB X-ray Structure Validation Report ⓘ

Nov 9, 2026 – 10:25 AM EST

PDB ID	:	36DV / pdb_000036dv
Title	:	Borrelia Burgdorferi SwrD
Authors	:	Nair, D.; Crane, B.R.
Deposited on	:	2026-06-02
Resolution	:	1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	75	67% · 29%
1	B	75	71% 29%
1	C	75	% 69% · 29%
1	D	75	% 71% · 28%
1	E	75	% 71% 29%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	75	

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 6075 atoms, of which 3234 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Flagellar protein FlbD.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	53	Total	C	H	N	O	S	0	0	0
			877	277	449	69	79	3			
1	B	53	Total	C	H	N	O	S	0	0	0
			877	277	449	69	79	3			
1	C	53	Total	C	H	N	O	S	0	0	0
			877	277	449	69	79	3			
1	D	54	Total	C	H	N	O	S	0	0	0
			895	283	459	70	80	3			
1	E	53	Total	C	H	N	O	S	0	0	0
			877	277	449	69	79	3			
1	F	53	Total	C	H	N	O	S	0	0	0
			877	277	449	69	79	3			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP O51281
B	0	MET	-	initiating methionine	UNP O51281
C	0	MET	-	initiating methionine	UNP O51281
D	0	MET	-	initiating methionine	UNP O51281
E	0	MET	-	initiating methionine	UNP O51281
F	0	MET	-	initiating methionine	UNP O51281

- Molecule 2 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	41	Total	H	O	2	0
			123	82	41		
2	B	49	Total	H	O	3	0
			147	98	49		
2	C	40	Total	H	O	1	0
			120	80	40		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	55	Total H O 165 110 55	2	0
2	E	39	Total H O 117 78 39	1	0
2	F	41	Total H O 123 82 41	2	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Flagellar protein FlbD

Chain A:




• Molecule 1: Flagellar protein FlbD

Chain B:




• Molecule 1: Flagellar protein FlbD

Chain C:




• Molecule 1: Flagellar protein FlbD

Chain D:




• Molecule 1: Flagellar protein FlbD

Chain E:



• Molecule 1: Flagellar protein FlbD

Chain F:


4%
63%
7%
29%
MO
Y12
A24
N25
P26
T29
V44
I51
K52
LEU
TYR
ARG
LYS
GLU
VAL
ALA
SER
LEU
GLU
LYS
ILE
LEU
GLY
GLU
GLY
ASN
GLY
GLY
VAL
GLU
LEU

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	68.03Å 72.06Å 75.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.99 – 1.80 29.99 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.7 (29.99-1.80) 98.7 (29.99-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	0.95 (at 1.60Å)	Xtriage
Refinement program	PHENIX (2.0_5936: ???)	Depositor
R, R_{free}	0.159 , 0.185 0.160 , 0.185	Depositor DCC
R_{free} test set	1931 reflections (4.42%)	wwPDB-VP
Wilson B-factor (Å^2)	27.0	Xtriage
Anisotropy	0.238	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.38 , 45.3	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	6075	wwPDB-VP
Average B, all atoms (Å^2)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 29.74 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4729e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	0.33	0/434	0.56	0/584
1	B	0.36	0/434	0.60	0/584
1	C	0.38	0/434	0.58	0/584
1	D	0.36	0/442	0.57	0/595
1	E	0.28	0/434	0.52	0/584
1	F	0.34	0/434	0.59	0/584
All	All	0.34	0/2612	0.57	0/3515

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	428	449	451	3	0
1	B	428	449	451	0	0
1	C	428	449	451	1	0
1	D	436	459	462	1	0
1	E	428	449	451	0	0
1	F	428	449	451	5	0
2	A	41	82	0	1	0
2	B	49	98	0	0	0
2	C	40	80	0	0	0
2	D	55	110	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	39	78	0	0	0
2	F	41	82	0	1	0
All	All	2841	3234	2717	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:0:MET:HG2	2:A:116:HOH:O	2.01	0.60
1:A:24:ALA:HB2	1:A:44:VAL:HG23	1.90	0.54
1:F:24:ALA:HB2	1:F:44:VAL:CG2	2.41	0.50
1:F:24:ALA:HB2	1:F:44:VAL:HG21	1.97	0.46
1:A:24:ALA:HB2	1:A:44:VAL:CG2	2.49	0.43
1:F:51:ILE:O	1:F:51:ILE:HG22	2.19	0.42
1:F:25:ASN:ND2	1:F:29:THR:OG1	2.52	0.42
1:D:0:MET:N	2:D:102:HOH:O	2.49	0.41
1:C:25:ASN:O	1:C:25:ASN:CG	2.64	0.41
1:F:0:MET:N	2:F:203:HOH:O	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	51/75 (68%)	50 (98%)	1 (2%)	0	100	100
1	B	51/75 (68%)	49 (96%)	2 (4%)	0	100	100
1	C	51/75 (68%)	51 (100%)	0	0	100	100
1	D	52/75 (69%)	52 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	E	51/75 (68%)	50 (98%)	1 (2%)	0	100 100
1	F	51/75 (68%)	49 (96%)	1 (2%)	1 (2%)	6 1
All	All	307/450 (68%)	301 (98%)	5 (2%)	1 (0%)	36 25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	25	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	49/66 (74%)	49 (100%)	0	100 100
1	B	49/66 (74%)	49 (100%)	0	100 100
1	C	49/66 (74%)	49 (100%)	0	100 100
1	D	50/66 (76%)	50 (100%)	0	100 100
1	E	49/66 (74%)	49 (100%)	0	100 100
1	F	49/66 (74%)	49 (100%)	0	100 100
All	All	295/396 (74%)	295 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	25	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	53/75 (70%)	-0.30	0	100	100	23, 34, 61, 96	0
1	B	53/75 (70%)	-0.43	0	100	100	20, 28, 57, 79	0
1	C	53/75 (70%)	-0.19	1 (1%)	66	66	22, 33, 62, 86	0
1	D	54/75 (72%)	-0.41	1 (1%)	66	66	22, 29, 54, 88	0
1	E	53/75 (70%)	-0.01	1 (1%)	66	66	24, 39, 69, 96	0
1	F	53/75 (70%)	0.25	3 (5%)	29	28	21, 39, 70, 104	0
All	All	319/450 (70%)	-0.18	6 (1%)	66	66	20, 34, 69, 104	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	53	LEU	4.0
1	C	0	MET	2.2
1	E	52	LYS	2.2
1	F	24	ALA	2.2
1	F	12	TYR	2.1
1	F	26	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.