



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 16, 2026 – 02:06 PM EDT

PDB ID : 36UI / pdb_000036ui
Title : Crystal Structure of IDH2 (R140Q) in complex with covalent inhibitor LY3410738
Authors : Antonysamy, S.S.
Deposited on : 2026-07-01
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	461	
1	B	461	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6647 atoms, of which 101 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Isocitrate dehydrogenase [NADP], mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	388	Total	C	N	O	S	0	0	0
			3073	1960	530	565	18			
1	B	412	Total	C	N	O	S	0	0	0
			3232	2054	560	600	18			

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	140	GLN	ARG	variant	UNP P48735
A	453	ALA	-	expression tag	UNP P48735
A	454	ALA	-	expression tag	UNP P48735
A	455	ALA	-	expression tag	UNP P48735
A	456	HIS	-	expression tag	UNP P48735
A	457	HIS	-	expression tag	UNP P48735
A	458	HIS	-	expression tag	UNP P48735
A	459	HIS	-	expression tag	UNP P48735
A	460	HIS	-	expression tag	UNP P48735
A	461	HIS	-	expression tag	UNP P48735
B	140	GLN	ARG	variant	UNP P48735
B	453	ALA	-	expression tag	UNP P48735
B	454	ALA	-	expression tag	UNP P48735
B	455	ALA	-	expression tag	UNP P48735
B	456	HIS	-	expression tag	UNP P48735
B	457	HIS	-	expression tag	UNP P48735
B	458	HIS	-	expression tag	UNP P48735
B	459	HIS	-	expression tag	UNP P48735
B	460	HIS	-	expression tag	UNP P48735
B	461	HIS	-	expression tag	UNP P48735

- Molecule 2 is 7-([(1S)-1-{4-[(1S)-2-cyclopropyl-1-(4-propanoylpiperazin-1-yl)ethyl]phenyl}ethyl]amino)-1-ethyl-1,4-dihydro-2H-pyrimido[4,5-d][1,3]oxazin-2-one (CCD ID: A1DLL) (formula: C₂₈H₃₈N₆O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 75	C 28	H 38	N 6	O 3	0	0
2	B	1	Total 75	C 28	H 38	N 6	O 3	0	0

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: $\text{C}_{21}\text{H}_{28}\text{N}_7\text{O}_{17}\text{P}_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
3	A	1	Total	C	H	N	O	P	0	0
			73	21	25	7	17	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	52	Total 52	O 52	0	0
4	B	67	Total 67	O 67	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Isocitrate dehydrogenase [NADP], mitochondrial



- Molecule 1: Isocitrate dehydrogenase [NADP], mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.72Å 116.64Å 129.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.57 – 2.20 29.57 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.57-2.20) 99.7 (29.57-2.20)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 2.20Å)	Xtriage
Refinement program	BUSTER 2.11.8	Depositor
R, R_{free}	0.224 , 0.247 0.215 , 0.238	Depositor DCC
R_{free} test set	2175 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	44.8	Xtriage
Anisotropy	0.387	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 31.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6647	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1DLL, NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	0/3140	1.02	0/4236
1	B	0.68	0/3302	1.07	2/4464 (0.0%)
All	All	0.65	0/6442	1.05	2/8700 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	65	PHE	CA-CB-CG	5.46	119.27	113.80
1	B	420	HIS	N-CA-C	5.39	119.99	113.41

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3073	0	3051	17	0
1	B	3232	0	3179	15	0
2	A	37	38	0	1	0
2	B	37	38	0	0	0
3	A	48	25	25	1	0
4	A	52	0	0	0	0
4	B	67	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	6546	101	6255	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:169:THR:HG21	1:A:238:TYR:HE2	1.45	0.80
1:A:169:THR:HG21	1:A:238:TYR:CE2	2.33	0.61
1:A:185:VAL:HG23	1:B:257:ALA:HB1	1.85	0.58
1:A:195:VAL:HG22	1:A:206:GLU:HG2	1.86	0.57
1:A:219:MET:HE3	1:A:221:MET:HG3	1.87	0.56
1:A:170:ILE:HD13	1:A:306:TRP:HB3	1.86	0.56
1:A:222:TYR:HB2	1:B:212:PHE:CZ	2.42	0.54
1:B:107:TYR:O	1:B:382:ARG:NH2	2.41	0.54
1:A:107:TYR:O	1:A:382:ARG:NH2	2.41	0.54
1:B:195:VAL:HG22	1:B:206:GLU:HG2	1.91	0.53
1:A:115:THR:O	3:A:502:NAP:H2N	2.09	0.52
1:A:327:LEU:HB3	1:A:348:HIS:HB3	1.91	0.52
1:B:205:LYS:HB3	1:B:207:TRP:CZ2	2.46	0.50
1:B:293:MET:HE3	1:B:297:VAL:HG11	1.94	0.50
1:A:184:PHE:CE1	1:A:218:GLY:HA3	2.48	0.48
1:B:152:ILE:HD13	1:B:369:ILE:HG22	1.95	0.48
1:A:151:PRO:HG3	2:A:501:A1DLL:C1	2.44	0.48
1:B:353:ARG:HG3	1:B:354:HIS:HD2	1.79	0.47
1:B:353:ARG:HG3	1:B:354:HIS:CD2	2.51	0.46
1:A:298:LEU:HD21	1:B:319:ILE:HG23	1.97	0.46
1:B:167:PRO:HG2	1:B:244:TRP:HH2	1.81	0.45
1:B:321:ALA:HB1	1:B:330:MET:HE1	1.98	0.44
1:A:324:PHE:HE2	1:B:161:VAL:HG22	1.83	0.44
1:B:311:TYR:HA	1:B:314:ASP:HB3	2.00	0.43
1:A:288:ARG:HD3	1:A:296:GLN:OE1	2.18	0.42
1:A:65:PHE:CE1	1:A:438:LEU:HB3	2.54	0.42
1:A:143:LEU:HB3	1:A:336:CYS:SG	2.60	0.42
1:B:335:VAL:HG22	1:B:342:ILE:CD1	2.51	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	380/461 (82%)	374 (98%)	6 (2%)	0	100	100
1	B	406/461 (88%)	400 (98%)	6 (2%)	0	100	100
All	All	786/922 (85%)	774 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	330/389 (85%)	320 (97%)	10 (3%)	36	49
1	B	343/389 (88%)	332 (97%)	11 (3%)	34	47
All	All	673/778 (86%)	652 (97%)	21 (3%)	35	48

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	172	ARG
1	A	242	LYS
1	A	262	PHE
1	A	282	LYS
1	A	291	ASP
1	A	293	MET
1	A	309	LYS
1	A	324	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	343	GLU
1	A	351	VAL
1	B	172	ARG
1	B	242	LYS
1	B	262	PHE
1	B	291	ASP
1	B	309	LYS
1	B	315	VAL
1	B	316	GLN
1	B	330	MET
1	B	331	THR
1	B	343	GLU
1	B	351	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	136	ASN
1	A	141	ASN
1	A	223	ASN
1	A	281	ASN
1	B	141	ASN
1	B	223	ASN
1	B	273	HIS
1	B	316	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAP	A	502	-	50,52,52	0.54	0	71,80,80	0.60	0
2	A1DLL	B	501	1	41,41,41	1.05	4 (9%)	52,58,58	0.76	2 (3%)
2	A1DLL	A	501	1	41,41,41	1.02	4 (9%)	52,58,58	0.86	2 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	A	502	-	-	7/35/67/67	0/5/5/5
2	A1DLL	B	501	1	-	0/28/53/53	0/5/5/5
2	A1DLL	A	501	1	-	0/28/53/53	0/5/5/5

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	A1DLL	C16-N4	4.51	1.53	1.48
2	A	501	A1DLL	C16-N4	4.38	1.53	1.48
2	B	501	A1DLL	C2-N2	3.31	1.43	1.39
2	A	501	A1DLL	C2-N2	3.23	1.43	1.39
2	A	501	A1DLL	C17-N4	2.09	1.51	1.47
2	A	501	A1DLL	C20-N4	2.09	1.51	1.47
2	B	501	A1DLL	C17-N4	2.08	1.51	1.47
2	B	501	A1DLL	C20-N4	2.07	1.51	1.47

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	A1DLL	C25-C24-C16	-2.61	111.09	114.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	A1DLL	C4-C3-C	2.55	125.92	120.79
2	B	501	A1DLL	C4-C3-C	2.37	125.56	120.79
2	B	501	A1DLL	O-C5-N2	2.00	119.48	117.14

There are no chirality outliers.

All (7) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	502	NAP	C5D-O5D-PN-O3
3	A	502	NAP	C5D-O5D-PN-O1N
3	A	502	NAP	O4D-C1D-N1N-C2N
3	A	502	NAP	O4D-C1D-N1N-C6N
3	A	502	NAP	C2D-C1D-N1N-C2N
3	A	502	NAP	C5D-O5D-PN-O2N
3	A	502	NAP	C2D-C1D-N1N-C6N


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	NAP	1	0
2	A	501	A1DLL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	388/461 (84%)	0.31	15 (3%) 43 40	33, 56, 81, 105	0
1	B	412/461 (89%)	0.18	11 (2%) 56 53	32, 48, 73, 83	0
All	All	800/922 (86%)	0.25	26 (3%) 49 46	32, 52, 77, 105	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	207	TRP	4.5
1	B	40	ALA	4.2
1	A	181	ALA	4.2
1	A	310	ASN	4.2
1	A	222	TYR	3.6
1	B	258	TYR	3.4
1	B	311	TYR	3.3
1	A	324	PHE	3.3
1	A	173	HIS	3.1
1	B	458	HIS	3.0
1	B	65	PHE	2.6
1	A	295	ALA	2.5
1	B	210	TYR	2.5
1	A	221	MET	2.3
1	B	174	ALA	2.3
1	A	427	LEU	2.3
1	A	160	LEU	2.2
1	A	255	LEU	2.2
1	B	457	HIS	2.2
1	A	422	LEU	2.1
1	A	144	GLY	2.1
1	A	257	ALA	2.1
1	B	180	LYS	2.1
1	A	449	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	212	PHE	2.1
1	A	214	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	A1DLL	A	501	37/37	0.93	0.08	48,51,58,60	0
2	A1DLL	B	501	37/37	0.94	0.07	40,47,60,62	0
3	NAP	A	502	48/48	0.97	0.06	38,41,44,45	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around A1DLL A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around A1DLL B 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.