

# Full wwPDB X-ray Structure Validation Report (i)

#### May 3, 2025 – 12:21 PM EDT

PDB ID : 3BDW / pdb\_00003bdw Title : Human CD94/NKG2A

Title : Human CD94/NKG2A thors : Sullivan, L.C.; Clements, C.S.

Authors : Sullivan, L.C.; Deposited on : 2007-11-15

Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1

Xtriage (Phenix) : 2.0rc1 EDS : 3.0

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.006 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.43.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
$R_{free}$	164625	5504 (2.50-2.50)
Clashscore	180529	6282 (2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)
RSRZ outliers	164620	5504 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	A	123	74%	20%	6%				
1	С	123	72%	21%	7% •				
2	В	120	63%	29%					
2	D	120	66%	26%	5% •				



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3922 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Natural killer cells antigen CD94.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	123		С		0	S	0	0	0
			1007	631	167	199	10	Ů	Ü	
1	C	122	Total	С	N	O	S	0	0	0
		122	1002	628	166	198	10		U	0

• Molecule 2 is a protein called NKG2-A/NKG2-B type II integral membrane protein.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	D	116	Total	С	N	О	S	0	0	0
	Б	110	931	585	166	171	9	0		
9	D	116	Total	С	N	О	S	0	0	0
2	ש	110	931	585	166	171	9	0	U	U

• Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	16	Total O 16 16	0	0
3	В	8	Total O 8 8	0	0
3	С	18	Total O 18 18	0	0
3	D	9	Total O 9 9	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Natural killer cells antigen CD94





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	44.65Å 34.73Å 152.90Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $89.72^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	30.59 - 2.50	Depositor
resolution (A)	30.59 - 2.50	EDS
% Data completeness	98.4 (30.59-2.50)	Depositor
(in resolution range)	97.1 (30.59-2.50)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.12	Depositor
$< I/\sigma(I) > 1$	1.96  (at  2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
$R, R_{free}$	0.234 , $0.270$	Depositor
it, it free	0.235 , $0.271$	DCC
$R_{free}$ test set	826 reflections $(5.16\%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	51.3	Xtriage
Anisotropy	0.200	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.35, 36.0	EDS
L-test for twinning <sup>2</sup>	$< L > = 0.45, < L^2> = 0.28$	Xtriage
Estimated twinning fraction	0.430 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3922	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.52% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.62	0/1033	0.81	1/1398 (0.1%)	
1	С	0.62	0/1028	0.77	0/1391	
2	В	0.57	0/954	0.81	3/1286 (0.2%)	
2	D	0.55	0/954	0.80	2/1286 (0.2%)	
All	All	0.59	0/3969	0.80	6/5361 (0.1%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	В	0	1
2	D	0	1
All	All	0	2

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	$\operatorname{Res}$	Type	Atoms	${f Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
2	В	178	PHE	N-CA-C	5.46	116.57	108.60
2	D	178	PHE	N-CA-C	5.44	116.55	108.60
2	В	184	HIS	N-CA-C	-5.08	101.29	109.82
2	D	183	HIS	N-CA-C	5.07	118.36	107.67
1	A	63	GLU	N-CA-C	-5.00	101.53	109.59
2	В	134	GLY	N-CA-C	5.00	118.54	112.49

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	В	183	HIS	Peptide



Continued from previous page...

Mol	Chain	Res	Type	Group
2	D	183	HIS	Peptide

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1007	0	911	16	0
1	С	1002	0	909	19	0
2	В	931	0	895	18	0
2	D	931	0	895	18	0
3	A	16	0	0	1	0
3	В	8	0	0	0	0
3	С	18	0	0	0	0
3	D	9	0	0	0	0
All	All	3922	0	3610	69	0

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:THR:HG22	1:A:84:GLU:H	1.35	0.88
1:C:81:THR:HG22	1:C:84:GLU:H	1.39	0.87
2:B:139:THR:HG22	2:B:142:GLU:H	1.41	0.85
2:D:139:THR:HG22	2:D:142:GLU:H	1.43	0.83
1:C:148:ASN:HD22	1:C:148:ASN:C	1.92	0.78
1:A:148:ASN:C	1:A:148:ASN:HD22	1.95	0.75
2:D:157:ILE:HG21	2:D:163:MET:HE2	1.69	0.74
1:C:61:CYS:O	1:C:62:GLN:O	2.05	0.74
2:B:144:LEU:HD11	2:B:189:MET:HE2	1.70	0.73
2:D:144:LEU:HD11	2:D:189:MET:HE2	1.73	0.71
2:B:157:ILE:HG21	2:B:163:MET:HE2	1.73	0.70
2:B:174:TRP:CE3	2:B:210:VAL:HG13	2.28	0.68
2:D:174:TRP:CE3	2:D:210:VAL:HG13	2.28	0.68
1:A:178:LEU:O	1:A:179:ILE:HG23	1.96	0.64



 $Continued\ from\ previous\ page...$ 

Continued from previ		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ (\mathring{\rm A})$	overlap (Å)
1:C:177:GLN:O	1:C:178:LEU:HD23	2.06	0.56
1:A:177:GLN:O	1:A:178:LEU:HD23	2.08	0.54
2:B:152:SER:HB3	2:B:231:HIS:CE1	2.43	0.54
1:C:66:VAL:HG22	1:C:73:TYR:HB2	1.90	0.53
2:D:206:LEU:HD22	2:D:219:ALA:HA	1.89	0.53
1:A:66:VAL:HG22	1:A:73:TYR:HB2	1.91	0.52
2:B:163:MET:HG2	2:B:211:LEU:HD11	1.90	0.52
2:B:206:LEU:HD22	2:B:219:ALA:HA	1.90	0.52
2:D:152:SER:HB3	2:D:231:HIS:CE1	2.44	0.52
1:C:61:CYS:O	1:C:62:GLN:C	2.53	0.51
1:C:140:LEU:HD13	1:C:141:PHE:CE2	2.46	0.51
1:A:99:LEU:HD11	1:A:155:TYR:CE2	2.46	0.50
2:D:163:MET:HE1	2:D:216:LEU:HD11	1.93	0.50
1:A:140:LEU:HD13	1:A:141:PHE:CE2	2.47	0.50
2:D:163:MET:HG2	2:D:211:LEU:HD11	1.93	0.49
2:D:157:ILE:HG21	2:D:163:MET:CE	2.41	0.49
1:A:148:ASN:ND2	1:A:150:LYS:H	2.11	0.49
2:D:119:CYS:SG	2:D:125:THR:HG22	2.53	0.48
2:B:158:ASP:OD1	2:B:230:LYS:NZ	2.41	0.48
1:C:99:LEU:HD11	1:C:155:TYR:CE2	2.48	0.48
1:A:119:LEU:HD23	1:A:153:ILE:HG13	1.96	0.48
2:B:119:CYS:SG	2:B:125:THR:HG22	2.53	0.48
2:D:198:ILE:HD13	2:D:198:ILE:C	2.38	0.48
2:D:158:ASP:OD1	2:D:230:LYS:NZ	2.42	0.47
1:C:148:ASN:ND2	1:C:150:LYS:H	2.13	0.47
1:A:68:TYR:HB2	2:B:124:ILE:HG22	1.98	0.46
2:B:198:ILE:C	2:B:198:ILE:HD13	2.40	0.46
1:C:148:ASN:C	1:C:148:ASN:ND2	2.65	0.46
2:B:163:MET:HE1	2:B:216:LEU:HD11	1.97	0.45
1:A:114:PHE:CE2	1:A:162:LEU:HD12	2.51	0.45
2:B:157:ILE:HG21	2:B:163:MET:CE	2.46	0.44
1:C:132:ASN:HD22	1:C:132:ASN:N	2.15	0.44
3:A:184:HOH:O	2:B:168:ILE:HG23	2.17	0.44
1:C:119:LEU:HD23	1:C:153:ILE:HG13	2.00	0.44
1:A:97:LEU:HD23	1:A:117:ILE:HG22	2.00	0.43
1:C:61:CYS:C	1:C:62:GLN:O	2.62	0.43
1:A:148:ASN:HD22	1:A:150:LYS:H	1.64	0.42
2:D:187:VAL:HG12	2:D:188:THR:O	2.18	0.42
1:A:132:ASN:HD22	1:A:132:ASN:N	2.16	0.42
1:A:179:ILE:HD13	1:A:179:ILE:N	2.35	0.42
2:B:133:ILE:HD12	2:B:133:ILE:N	2.35	0.42



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
		$\operatorname{distance}\ ( ext{\AA})$	overlap (Å)
1:C:68:TYR:HB2	2:D:124:ILE:HG22	2.01	0.41
1:A:148:ASN:HD22	1:A:149:THR:N	2.19	0.41
1:C:114:PHE:CE2	1:C:162:LEU:HD12	2.55	0.41
2:B:198:ILE:HD11	2:B:218:SER:CA	2.51	0.41
1:C:97:LEU:HD23	1:C:117:ILE:HG22	2.03	0.41
2:D:210:VAL:HG11	2:D:225:ILE:HD13	2.02	0.41
1:C:65:TRP:CD2	1:C:74:PHE:HB2	2.55	0.41
2:B:187:VAL:HG12	2:B:188:THR:O	2.21	0.40
2:D:198:ILE:HD11	2:D:218:SER:CA	2.51	0.40
2:D:133:ILE:HD12	2:D:133:ILE:N	2.36	0.40
2:B:170:SER:HB2	2:B:228:HIS:HE1	1.87	0.40
2:D:139:THR:HG23	2:D:140:TRP:N	2.36	0.40
1:C:113:GLN:OE1	1:C:171:ARG:NH2	2.55	0.40
1:C:148:ASN:HD22	1:C:149:THR:N	2.20	0.40

There are no symmetry-related clashes.

# 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	121/123 (98%)	111 (92%)	7 (6%)	3 (2%)	4	7
1	С	120/123 (98%)	110 (92%)	8 (7%)	2 (2%)	7	14
2	В	112/120 (93%)	103 (92%)	8 (7%)	1 (1%)	14	28
2	D	112/120 (93%)	103 (92%)	8 (7%)	1 (1%)	14	28
All	All	465/486 (96%)	427 (92%)	31 (7%)	7 (2%)	8	16

#### All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	62	GLN



Continued from previous page...

Mol	Chain	Res	Type
1	A	62	GLN
2	В	213	VAL
1	A	61	CYS
1	С	61	CYS
2	D	213	VAL
1	A	58	CYS

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	113/114 (99%)	100 (88%)	13 (12%)	4 9		
1	$\mathbf{C}$	113/114 (99%)	100 (88%)	13 (12%)	4 9		
2	В	105/109 (96%)	92 (88%)	13 (12%)	4 7		
2	D	105/109~(96%)	92 (88%)	13 (12%)	4 7		
All	All	436/446 (98%)	384 (88%)	52 (12%)	4 8		

All (52) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	66	VAL
1	A	81	THR
1	A	83	ASN
1	A	93	LYS
1	A	112	GLN
1	A	126	THR
1	A	132	ASN
1	A	140	LEU
1	A	143	SER
1	A	145	GLU
1	A	148	ASN
1	A	177	GLN
1	A	179	ILE
2	В	122	GLU
2	В	129	SER



Continued from previous page...

Mol	$\frac{\text{Chain}}{\text{Chain}}$	Res	$egin{array}{c}  ext{rus page} \  ext{Type} \end{array}$
2	В	139	THR
$\frac{2}{2}$	В	159	THR ASN
2	В	160	GLU
2 2	В	166	LEU
$\frac{2}{2}$	В	181	SER
2	В	195	LYS
2 2 2	В	198	ILE
$\frac{2}{2}$	В	210	ILE VAL
2	В	213	VAL
2	В	214	ASN
$\frac{1}{2}$	В	231	ASN HIS
2	C	66	VAL
1	B C C C C C C C C C C C C C C C C C C C	81	THR
1	С	83	ASN LYS
1	С	93	LYS
1	С	112	GLN
1	С	126	GLN THR
1	С	132	ASN
1	С	140	ASN LEU
1	С	143	SER
1	С	145	SER GLU
1	С	148	ASN
1	С	177	ASN GLN
1	С	179	ILE
2	D	122	ILE GLU
2	D	129	SER
2	D	139	THR
2	D	159	ASN
2	D	160	GLU
2	D	166	LEU
2 2	D	181	SER
	D	195	LYS
2	D	198	ILE
2	D	210	VAL
2	D	213	VAL
2	D	214	ASN
2	D	231	HIS

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such side chains are listed below:

Mol	Chain	Res	Type
1	A	83	ASN



Continued from previous page...

Mol	Chain	Res	Type
1	A	87	HIS
1	A	98	GLN
1	A	100	GLN
1	A	101	ASN
1	A	132	ASN
1	A	148	ASN
2	В	212	GLN
2	В	214	ASN
2	В	228	HIS
2	В	231	HIS
1	С	83	ASN
1	С	87	HIS
1	С	98	GLN
1	С	100	GLN
1	С	101	ASN
1	С	132	ASN
1	C C C C C C	148	ASN
1	С	160	ASN
2	D	212	GLN
2	D	214	ASN
2	D	228	HIS
2	D	231	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

# 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry (i)

There are no ligands in this entry.



# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	#	RSR	$\mathbb{Z}>2$	$OWAB(A^2)$	Q<0.9
1	A	123/123 (100%)	-1.26	0	100	100	37, 41, 45, 49	0
1	С	122/123~(99%)	-1.23	0	100	100	37, 41, 45, 49	0
2	В	116/120 (96%)	-1.25	0	100	100	37, 41, 43, 45	0
2	D	116/120~(96%)	-1.22	0	100	100	37, 41, 43, 45	0
All	All	477/486 (98%)	-1.24	0	100	100	37, 41, 45, 49	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

There are no ligands in this entry.

## 6.5 Other polymers (i)

There are no such residues in this entry.

