



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 05:18 AM UTC

PDB ID : 3H5E / pdb_00003h5e
Title : LeuD_1-156 small subunit of isopropylmalate isomerase (Rv2987c) from mycobacterium tuberculosis
Authors : Manikandan, K.; Geerlof, A.; Weiss, M.S.
Deposited on : 2009-04-22
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	
1	B	159	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2522 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-isopropylmalate dehydratase small subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	154	1200	769	211	219	1	9	2	0
1	B	152	1187	762	209	215	1	6	2	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP P65277
A	0	ALA	-	expression tag	UNP P65277
A	1A	SER	-	cloning artifact	UNP P65277
A	122	ILE	LEU	SEE REMARK 999	UNP P65277
B	-1	GLY	-	expression tag	UNP P65277
B	0	ALA	-	expression tag	UNP P65277
B	1A	SER	-	cloning artifact	UNP P65277
B	122	ILE	LEU	SEE REMARK 999	UNP P65277


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	82	Total	O	0	0
			82	82		
2	B	53	Total	O	0	0
			53	53		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3-isopropylmalate dehydratase small subunit



- Molecule 1: 3-isopropylmalate dehydratase small subunit



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	64.62Å 64.62Å 419.82Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.00 30.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.4 (30.00-2.00) 98.4 (30.00-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.4.0069	Depositor
R, R_{free}	0.234 , 0.269 0.236 , 0.267	Depositor DCC
R_{free} test set	1132 reflections (3.08%)	wwPDB-VP
Wilson B-factor (Å ²)	37.9	Xtrriage
Anisotropy	0.421	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2522	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.84	2/1233 (0.2%)	1.00	3/1674 (0.2%)
1	B	0.79	0/1220	0.88	1/1656 (0.1%)
All	All	0.82	2/2453 (0.1%)	0.94	4/3330 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	5	HIS	CB-CG	-7.00	1.40	1.50
1	A	3	ALA	CA-CB	-5.32	1.35	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	5	HIS	CA-CB-CG	10.38	124.18	113.80
1	A	25	ILE	CA-C-N	-6.65	112.91	119.76
1	A	25	ILE	C-N-CA	-6.65	112.91	119.76
1	B	99	ASP	N-CA-C	5.63	117.42	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1200	0	1197	15	0
1	B	1187	0	1188	18	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	82	0	0	2	0
2	B	53	0	0	1	0
All	All	2522	0	2385	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77:ARG:HD3	2:A:218:HOH:O	1.74	0.88
1:B:5:HIS:NE2	1:B:140:GLN:HB2	1.89	0.86
1:B:140:GLN:HA	1:B:140:GLN:HE21	1.44	0.81
1:A:99:ASP:OD2	1:A:99:ASP:N	2.21	0.74
1:A:74:GLY:O	1:A:75:SER:C	2.39	0.65
1:A:21:THR:HG21	1:A:76:SER:HA	1.79	0.64
1:A:103:GLY:O	1:A:107:LYS:HG3	2.03	0.59
1:B:16[B]:ARG:HD2	1:B:46:TRP:CH2	2.37	0.58
1:B:77:ARG:O	1:B:78:GLU:HB2	2.05	0.55
1:B:121[A]:GLU:HG3	1:B:125:LYS:HE3	1.88	0.54
1:B:40:ASP:OD2	2:B:170:HOH:O	2.19	0.51
1:B:4:PHE:CE2	1:B:139:LEU:HD23	2.46	0.51
1:A:76:SER:HB3	1:A:100:ILE:HG22	1.93	0.50
1:A:28:VAL:HG13	2:A:178:HOH:O	2.12	0.50
1:B:25:ILE:O	1:B:25:ILE:HG23	2.13	0.49
1:A:139:LEU:HD11	1:A:155:ILE:HD13	1.96	0.48
1:B:102:ARG:NH1	1:B:155:ILE:HD13	2.29	0.48
1:A:146:ALA:O	1:A:147:ALA:C	2.57	0.47
1:A:16[A]:ARG:HD3	1:A:46:TRP:CH2	2.50	0.47
1:B:10:ILE:HB	1:B:62:GLY:HA2	1.97	0.46
1:B:9:GLY:HA3	1:B:63:SER:OG	2.15	0.45
1:A:10:ILE:HB	1:A:62:GLY:HA2	1.97	0.45
1:A:95:SER:HA	1:A:115:VAL:O	2.16	0.44
1:B:97:PHE:HB2	1:B:102:ARG:HD3	1.99	0.44
1:B:102:ARG:HH12	1:B:155:ILE:HD13	1.82	0.44
1:B:102:ARG:NH1	1:B:112:ALA:O	2.51	0.43
1:A:16[A]:ARG:HE	1:A:16[A]:ARG:HB3	1.68	0.42
1:B:74:GLY:HA3	1:B:75:SER:C	2.44	0.42
1:A:75:SER:O	1:A:76:SER:O	2.37	0.42
1:B:140:GLN:HE21	1:B:140:GLN:CA	2.21	0.41
1:A:138:ASN:OD1	1:A:138:ASN:C	2.63	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:20:ASP:C	1:B:20:ASP:OD2	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	154/159 (97%)	148 (96%)	4 (3%)	2 (1%)	9 5
1	B	152/159 (96%)	144 (95%)	7 (5%)	1 (1%)	18 14
All	All	306/318 (96%)	292 (95%)	11 (4%)	3 (1%)	12 8

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	76	SER
1	A	75	SER
1	B	78	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	126/127 (99%)	117 (93%)	9 (7%)	13 10
1	B	125/127 (98%)	120 (96%)	5 (4%)	28 27

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	251/254 (99%)	237 (94%)	14 (6%)	22	16

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	HIS
1	A	16[A]	ARG
1	A	16[B]	ARG
1	A	99	ASP
1	A	102	ARG
1	A	121[A]	GLU
1	A	121[B]	GLU
1	A	149	VAL
1	A	154	LYS
1	B	25	ILE
1	B	32	ARG
1	B	75	SER
1	B	140	GLN
1	B	155	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	104	ASN
1	A	140	GLN
1	B	117	GLN
1	B	140	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	154/159 (96%)	1.07	15 (9%) 13 12	26, 42, 53, 62	5 (3%)
1	B	152/159 (95%)	1.45	36 (23%) 2 2	25, 42, 56, 66	4 (2%)
All	All	306/318 (96%)	1.26	51 (16%) 4 4	25, 42, 54, 66	9 (2%)

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	156	ASP	5.2
1	B	73	THR	4.8
1	B	75	SER	4.6
1	B	4	PHE	4.5
1	A	76	SER	4.3
1	B	76	SER	3.9
1	A	3	ALA	3.6
1	A	77	ARG	3.3
1	B	44	ALA	3.3
1	A	4	PHE	3.3
1	B	77	ARG	3.1
1	B	139	LEU	2.9
1	B	45	GLY	2.9
1	B	151	LEU	2.9
1	B	149	VAL	2.9
1	A	99	ASP	2.9
1	B	30	LEU	2.8
1	B	98	GLY	2.7
1	B	72	GLY	2.7
1	A	33	VAL	2.7
1	B	148	THR	2.6
1	A	75	SER	2.6
1	B	147	ALA	2.6
1	A	107	LYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	27	ALA	2.5
1	A	46	TRP	2.5
1	B	122	ILE	2.5
1	B	145	THR	2.5
1	B	5	HIS	2.5
1	A	50	PRO	2.4
1	A	5	HIS	2.4
1	B	150	VAL	2.4
1	B	21	THR	2.4
1	B	29	PHE	2.3
1	B	74	GLY	2.3
1	A	48	SER	2.2
1	B	127	ILE	2.2
1	A	108	ALA	2.2
1	B	133	LEU	2.2
1	B	135	ILE	2.2
1	B	143	ILE	2.2
1	B	8	SER	2.2
1	B	26	PRO	2.1
1	B	36	THR	2.2
1	B	25	ILE	2.1
1	B	41	GLY	2.1
1	B	100	ILE	2.1
1	B	126	LEU	2.1
1	B	155	ILE	2.1
1	A	73	THR	2.0
1	B	129	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.