



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 28, 2025 – 01:05 PM EDT

PDB ID : 3SZW / pdb\_00003szw  
Title : Crystal structure of sulfide:quinone oxidoreductase Cys128Ser variant from *Acidithiobacillus ferrooxidans* in complex with decylubiquinone  
Authors : Cherney, M.M.; Zhang, Y.; James, M.N.G.; Weiner, J.H.  
Deposited on : 2011-07-19  
Resolution : 2.20 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0rc1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.43.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	437	

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 3641 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Sulfide-quinone reductase, putative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	421	3233	2076	541	593	23	0	3	0

There are 5 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	expression tag	UNP B7JBP8
A	-1	ARG	-	expression tag	UNP B7JBP8
A	0	GLY	-	expression tag	UNP B7JBP8
A	1	SER	-	expression tag	UNP B7JBP8
A	128	SER	CYS	engineered mutation	UNP B7JBP8

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is DODECYL-BETA-D-MALTOSE (CCD ID: LMT) (formula:  $C_{24}H_{46}O_{11}$ ).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	C O	0	0
			35	24 11		

- Molecule 4 is 2-decyl-5,6-dimethoxy-3-methylcyclohexa-2,5-diene-1,4-dione (CCD ID: DCQ) (formula:  $C_{19}H_{30}O_4$ ).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	C O	0	0
			23	19 4		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	O S	0	0
			5	4 1		

- Molecule 6 is HYDROSULFURIC ACID (CCD ID: H2S) (formula: H<sub>2</sub>S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	S	0	0
			1	1		
6	A	1	Total	S	0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total S 2 2	0	1
6	A	1	Total S 1 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	287	Total O 287 287	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sulfide-quinone reductase, putative



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	149.91Å 149.91Å 81.39Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.75 – 2.20 50.75 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.75-2.20) 94.7 (50.75-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.16	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.78 (at 2.20Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, $R_{free}$	0.168 , 0.219 0.163 , 0.214	Depositor DCC
$R_{free}$ test set	1401 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.5	Xtrriage
Anisotropy	0.462	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 49.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3641	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, FAD, LMT, H2S, DCQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/3324	0.78	2/4497 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	184	ILE	CA-C-N	5.33	124.97	119.05
1	A	184	ILE	C-N-CA	5.33	124.97	119.05

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3233	0	3222	54	0
2	A	53	0	30	2	0
3	A	35	0	46	1	0
4	A	23	0	30	3	0
5	A	5	0	0	1	0
6	A	5	0	0	0	0
7	A	287	0	0	5	1
All	All	3641	0	3328	57	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:407:SER:N	1:A:408:GLU:HB2	1.66	1.10
1:A:407:SER:H	1:A:408:GLU:HB2	1.29	0.95
1:A:166:GLU:HG3	1:A:360:MET:HE1	1.61	0.83
1:A:76:GLN:HG2	1:A:90:LEU:HB3	1.63	0.81
1:A:407:SER:HB2	1:A:408:GLU:HA	1.68	0.74
1:A:341:GLU:OE1	1:A:343:ARG:NH2	2.19	0.74
1:A:6:ILE:HD12	1:A:17:ALA:HA	1.70	0.71
1:A:310:GLU:HG2	1:A:311:THR:N	2.08	0.66
1:A:418:MET:HE3	1:A:418:MET:HA	1.80	0.63
1:A:404:MET:HG3	7:A:837:HOH:O	1.97	0.62
1:A:166:GLU:HG3	1:A:360:MET:CE	2.30	0.61
3:A:501:LMT:H6D	7:A:753:HOH:O	2.03	0.58
1:A:14:MET:HB2	1:A:15:PRO:HD3	1.86	0.57
1:A:355:VAL:HG21	4:A:502:DCQ:H14	1.87	0.57
1:A:385:ARG:HD2	7:A:747:HOH:O	2.05	0.56
1:A:360:MET:HE3	1:A:360:MET:HA	1.86	0.56
1:A:8:GLY:O	1:A:13:GLY:HA3	2.06	0.55
1:A:407:SER:CA	1:A:408:GLU:HB2	2.38	0.54
1:A:412:GLU:C	1:A:414:VAL:H	2.13	0.54
1:A:152:ILE:O	1:A:189:PHE:HA	2.09	0.53
1:A:7:LEU:HD13	1:A:105:THR:CG2	2.38	0.53
1:A:407:SER:HB2	1:A:408:GLU:CA	2.37	0.51
1:A:140:TYR:OH	1:A:175:LEU:HD13	2.11	0.51
1:A:415:LEU:C	1:A:417:LYS:N	2.69	0.51
1:A:412:GLU:O	1:A:415:LEU:HG	2.10	0.50
1:A:23:ALA:HB1	1:A:337:LYS:HD2	1.94	0.49
1:A:227:LYS:NZ	7:A:877:HOH:O	2.44	0.49
1:A:416:PHE:C	1:A:416:PHE:CD1	2.91	0.48
1:A:415:LEU:C	1:A:417:LYS:H	2.20	0.48
1:A:326[B]:GLU:HG3	1:A:411:TYR:HE1	1.78	0.48
1:A:415:LEU:C	1:A:415:LEU:HD12	2.39	0.47
1:A:395:GLU:HG2	7:A:834:HOH:O	2.13	0.47
1:A:248:ILE:O	1:A:249:LYS:HB3	2.13	0.47
1:A:92:ASP:OD1	1:A:94:ASN:HB2	2.14	0.47
2:A:500:FAD:H9	2:A:500:FAD:O3'	2.15	0.47
1:A:278:ASN:HB2	1:A:279:PRO:CD	2.45	0.46
1:A:14:MET:CB	1:A:15:PRO:HD3	2.45	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:153:GLY:HA3	1:A:190:ILE:O	2.16	0.45
1:A:10:GLY:HA3	2:A:500:FAD:O5B	2.17	0.44
1:A:356[B]:CYS:SG	1:A:367:PHE:CZ	3.11	0.44
1:A:76:GLN:HE21	1:A:76:GLN:HB2	1.60	0.43
1:A:359:ASP:O	1:A:360:MET:HE3	2.19	0.43
1:A:194:PRO:HG3	1:A:312:THR:CG2	2.49	0.43
1:A:370:LEU:O	1:A:376:ARG:HB3	2.18	0.43
1:A:326[B]:GLU:HG3	1:A:411:TYR:CE1	2.54	0.43
1:A:184:ILE:HA	1:A:185:PRO:HD3	1.89	0.42
1:A:301:GLY:HA2	1:A:325:ILE:HG12	2.01	0.42
4:A:502:DCQ:H7	4:A:502:DCQ:H1M	1.82	0.42
1:A:335:ASN:OD1	1:A:345:GLY:HA3	2.19	0.42
1:A:-2:MET:N	5:A:503:SO4:O4	2.49	0.41
1:A:311:THR:O	1:A:311:THR:HG23	2.19	0.41
1:A:402:MET:N	1:A:402:MET:HE2	2.36	0.41
1:A:411:TYR:CZ	4:A:502:DCQ:H1MA	2.56	0.41
1:A:410:PHE:C	1:A:410:PHE:CD2	2.98	0.41
1:A:412:GLU:C	1:A:414:VAL:N	2.79	0.41
1:A:14:MET:HG3	1:A:39:PHE:CD1	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:871:HOH:O	7:A:871:HOH:O[11_655]	1.91	0.29

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	422/437 (97%)	402 (95%)	19 (4%)	1 (0%)	44 52

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	410	PHE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	336/347 (97%)	322 (96%)	14 (4%)	25 33

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	LEU
1	A	24	LEU
1	A	94	ASN
1	A	144	LEU
1	A	175	LEU
1	A	177	LYS
1	A	202	GLN
1	A	310	GLU
1	A	339	ASP
1	A	360	MET
1	A	373	LEU
1	A	416	PHE
1	A	417	LYS
1	A	418	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	36	ASN
1	A	64	HIS
1	A	72	HIS
1	A	80	GLN
1	A	86	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	234	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 5 are modelled with single atom - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	LMT	A	501	-	36,36,36	1.28	3 (8%)	47,47,47	1.51	9 (19%)
4	DCQ	A	502	-	23,23,23	2.60	10 (43%)	27,29,29	1.07	1 (3%)
5	SO4	A	503	-	4,4,4	0.25	0	6,6,6	0.05	0
2	FAD	A	500	-	54,58,58	3.66	25 (46%)	71,89,89	2.29	14 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '·' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LMT	A	501	-	-	11/21/61/61	0/2/2/2
4	DCQ	A	502	-	-	5/14/38/38	0/1/1/1
2	FAD	A	500	-	-	4/30/50/50	0/6/6/6

All (38) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	FAD	O2'-C2'	-11.69	1.18	1.43
2	A	500	FAD	O4-C4	9.00	1.40	1.23
2	A	500	FAD	O2-C2	8.81	1.41	1.24
4	A	502	DCQ	O5-C5	7.27	1.39	1.23
2	A	500	FAD	C4X-N5	6.97	1.45	1.30
2	A	500	FAD	C10-N1	6.42	1.46	1.33
2	A	500	FAD	PA-O3P	6.08	1.66	1.59
2	A	500	FAD	P-O3P	5.70	1.65	1.59
4	A	502	DCQ	O2-C2	5.27	1.35	1.23
2	A	500	FAD	C9-C8	4.78	1.46	1.39
2	A	500	FAD	C2A-N3A	4.59	1.39	1.32
2	A	500	FAD	P-O1P	4.43	1.66	1.50
2	A	500	FAD	PA-O1A	4.29	1.65	1.50
2	A	500	FAD	C6-C5X	4.27	1.46	1.40
2	A	500	FAD	O4B-C1B	4.02	1.46	1.40
2	A	500	FAD	C9-C9A	3.87	1.45	1.39
2	A	500	FAD	C6-C7	3.87	1.44	1.39
3	A	501	LMT	C3'-C4'	-3.86	1.41	1.52
3	A	501	LMT	O5B-C1B	3.86	1.51	1.41
2	A	500	FAD	C8A-N7A	3.65	1.41	1.34
2	A	500	FAD	C2-N1	3.57	1.44	1.36
4	A	502	DCQ	C3-C2	-3.48	1.38	1.48
4	A	502	DCQ	C6-C1	3.48	1.41	1.35
2	A	500	FAD	C2A-N1A	3.48	1.40	1.33
2	A	500	FAD	C6A-N6A	3.45	1.46	1.34
4	A	502	DCQ	C4-C5	-3.38	1.39	1.48
2	A	500	FAD	C10-N10	2.96	1.43	1.37
4	A	502	DCQ	C7-C6	2.96	1.58	1.51
4	A	502	DCQ	C6-C5	-2.78	1.38	1.46
2	A	500	FAD	C5'-C4'	2.68	1.55	1.51
3	A	501	LMT	O5'-C5'	2.58	1.50	1.44
2	A	500	FAD	C8M-C8	2.50	1.55	1.51
2	A	500	FAD	C4X-C10	-2.32	1.37	1.44
4	A	502	DCQ	C4-C3	2.24	1.44	1.36
2	A	500	FAD	C4-N3	2.20	1.43	1.38
4	A	502	DCQ	C1-C2	-2.19	1.39	1.47

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	FAD	C5X-N5	2.13	1.43	1.39
4	A	502	DCQ	O4-C4M	-2.01	1.40	1.45

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	FAD	O4B-C1B-N9A	13.65	126.84	108.75
2	A	500	FAD	N3A-C2A-N1A	-5.79	120.82	128.67
3	A	501	LMT	C1-O1'-C1'	4.65	121.62	113.68
3	A	501	LMT	O5'-C5'-C4'	4.21	118.43	109.72
2	A	500	FAD	O2'-C2'-C3'	4.14	118.93	109.25
2	A	500	FAD	C4X-C10-N10	3.86	122.00	116.48
4	A	502	DCQ	C1M-C1-C6	-3.42	118.82	124.45
2	A	500	FAD	O2A-PA-O3P	2.96	115.28	107.27
2	A	500	FAD	C4-N3-C2	-2.83	120.61	125.64
2	A	500	FAD	C10-C4X-N5	-2.79	119.11	124.81
3	A	501	LMT	C6'-C5'-C4'	-2.77	105.60	113.38
3	A	501	LMT	C1B-O5B-C5B	-2.72	108.41	113.72
3	A	501	LMT	C1B-O1B-C4'	-2.62	111.77	117.98
3	A	501	LMT	O1'-C1-C2	2.61	118.21	109.37
2	A	500	FAD	O4B-C4B-C5B	2.57	117.57	109.33
2	A	500	FAD	C4-C4X-N5	2.52	121.69	118.21
2	A	500	FAD	C4X-C4-N3	2.47	119.54	113.25
2	A	500	FAD	C4B-O4B-C1B	-2.35	107.78	109.92
3	A	501	LMT	C3'-C4'-C5'	2.27	115.97	110.93
3	A	501	LMT	C1'-O5'-C5'	2.09	117.79	113.72
3	A	501	LMT	C2'-C3'-C4'	2.04	114.31	109.68
2	A	500	FAD	O3B-C3B-C2B	2.03	118.32	111.82
2	A	500	FAD	O3'-C3'-C4'	2.01	113.50	108.93
2	A	500	FAD	O4-C4-C4X	-2.00	121.25	126.53

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	LMT	O5'-C5'-C6'-O6'
4	A	502	DCQ	C9-C10-C11-C12
3	A	501	LMT	C4'-C5'-C6'-O6'
3	A	501	LMT	C11-C10-C9-C8
3	A	501	LMT	C6-C7-C8-C9
4	A	502	DCQ	C10-C11-C12-C13
4	A	502	DCQ	C11-C12-C13-C14

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
3	A	501	LMT	C4-C5-C6-C7
3	A	501	LMT	C1-C2-C3-C4
4	A	502	DCQ	C6-C7-C8-C9
2	A	500	FAD	O4B-C4B-C5B-O5B
2	A	500	FAD	C2'-C3'-C4'-O4'
4	A	502	DCQ	C7-C8-C9-C10
3	A	501	LMT	C3-C4-C5-C6
3	A	501	LMT	C9-C10-C11-C12
3	A	501	LMT	C2-C1-O1'-C1'
3	A	501	LMT	C5-C6-C7-C8
2	A	500	FAD	C2'-C3'-C4'-C5'
3	A	501	LMT	O1'-C1-C2-C3
2	A	500	FAD	O2'-C2'-C3'-C4'


There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	LMT	1	0
4	A	502	DCQ	3	0
5	A	503	SO4	1	0
2	A	500	FAD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	421/437 (96%)	-0.49	11 (2%) 57 54	16, 32, 65, 110	3 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	415	LEU	6.9
1	A	414	VAL	5.5
1	A	411	TYR	4.6
1	A	418	MET	3.7
1	A	410	PHE	3.4
1	A	413	LYS	3.2
1	A	416	PHE	2.9
1	A	409	PRO	2.6
1	A	412	GLU	2.4
1	A	-1	ARG	2.1
1	A	56	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	DCQ	A	502	23/23	0.82	0.26	73,82,90,97	0
5	SO4	A	503	5/5	0.88	0.12	80,89,92,92	0
6	H2S	A	506[A]	1/1	0.88	0.15	59,59,59,59	1
6	H2S	A	506[B]	1/1	0.88	0.15	24,24,24,24	1
6	H2S	A	505	1/1	0.90	0.17	59,59,59,59	0
3	LMT	A	501	35/35	0.92	0.09	24,38,45,50	0
6	H2S	A	507	1/1	0.93	0.11	60,60,60,60	0
2	FAD	A	500	53/53	0.97	0.06	16,25,33,37	0
6	H2S	A	504	1/1	0.98	0.04	38,38,38,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around LMT A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around FAD A 500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.