Full wwPDB X-ray Structure Validation Report (i)

Mar 28, 2026 – 05:24 AM UTC

PDB ID	:	3TDH / pdb_00003tdh
Title	:	Structure of the regulatory fragment of sccharomyces cerevisiae AMPK in complex with AMP
Authors	:	Mayer, F.V.; Heath, R.; Underwood, E.; Sanders, M.J.; Carmena, D.; McCartney, R.; Leiper, F.C.; Xiao, B.; Jing, C.; Walker, P.A.; Haire, L.F.; Ogrodowicz, R.; Martin, S.R.; Schmidt, M.C.; Gamblin, S.J.; Carling, D.
Deposited on	:	2011-08-11
Resolution	:	2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	179	13% 68% 10% 22%
2	B	113	34% 88% . 8%
3	C	323	14% 85% 11% . .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4599 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carbon catabolite-derepressing protein kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	139	Total	C	N	O	S	0	0	0
			1109	726	184	194	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	455	GLY	-	expression tag	UNP P06782
A	456	PRO	-	expression tag	UNP P06782

- Molecule 2 is a protein called SNF1 protein kinase subunit beta-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	104	Total	C	N	O	S	0	0	0
			828	538	131	157	2			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	303	MET	-	expression tag	UNP P34164

- Molecule 3 is a protein called Nuclear protein SNF4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	316	Total	C	N	O	S	0	0	0
			2471	1577	412	472	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	MET	-	expression tag	UNP P12904
C	1	ALA	-	expression tag	UNP P12904

- Molecule 4 is ADENOSINE MONOPHOSPHATE (CCD ID: AMP) (formula: $C_{10}H_{14}N_5O_7P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	C	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	26	Total	O	0	0
			26	26		
5	B	34	Total	O	0	0
			34	34		
5	C	108	Total	O	0	0
			108	108		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Carbon catabolite-derepressing protein kinase

Chain A:



• Molecule 2: SNF1 protein kinase subunit beta-2

Chain B:



• Molecule 3: Nuclear protein SNF4

Chain C:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α, β, γ	94.97Å 242.38Å 79.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.30 25.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.6 (25.00-2.30) 99.5 (25.00-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.05 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.248 , 0.290 0.244 , 0.283	Depositor DCC
R_{free} test set	2062 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å^2)	45.5	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 35.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	4599	wwPDB-VP
Average B, all atoms (Å^2)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.62% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.42	0/1138	0.76	0/1544
2	B	0.48	0/851	0.73	0/1172
3	C	0.46	0/2506	0.78	0/3398
All	All	0.45	0/4495	0.77	0/6114

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1109	0	1121	15	0
2	B	828	0	802	7	0
3	C	2471	0	2525	22	0
4	C	23	0	12	0	0
5	A	26	0	0	0	0
5	B	34	0	0	0	0
5	C	108	0	0	0	0
All	All	4599	0	4460	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:610:PHE:HZ	2:B:379:ASN:OD1	1.79	0.66
1:A:483:LEU:HG	1:A:490:ALA:HB1	1.80	0.64
3:C:81:THR:O	3:C:85:ILE:HG12	1.98	0.64
3:C:85:ILE:CD1	3:C:241:VAL:HG21	2.32	0.59
3:C:84:PHE:O	3:C:88:ILE:HG22	2.02	0.59
3:C:49:LEU:HD12	3:C:52:LYS:HE2	1.85	0.59
3:C:139:MET:HE1	3:C:165:LEU:O	2.07	0.53
3:C:18:ALA:O	3:C:22:ILE:HG12	2.11	0.51
1:A:609:THR:HG23	2:B:399:TYR:HE1	1.75	0.51
1:A:575:ILE:HD11	1:A:581:LEU:HB2	1.93	0.50
3:C:315:LYS:HG3	3:C:319:LEU:HD22	1.94	0.49
3:C:136:CYS:HB3	3:C:318:LEU:HD21	1.94	0.49
1:A:500:LYS:HA	3:C:59:GLN:OE1	2.12	0.48
1:A:547:ARG:HG2	1:A:565:LEU:HB3	1.94	0.48
3:C:146:ARG:HG3	3:C:166:THR:HG22	1.95	0.48
3:C:44:LEU:HD13	3:C:53:SER:HB2	1.96	0.47
3:C:235:VAL:HG21	3:C:296:PHE:CE2	2.50	0.47
3:C:283:ILE:HG23	3:C:295:PHE:CE2	2.50	0.46
1:A:624:MET:HE3	2:B:391:LEU:HD22	1.97	0.46
1:A:482:MET:CB	3:C:262:MET:HE1	2.47	0.45
3:C:15:GLN:NE2	3:C:184:LEU:HA	2.31	0.45
1:A:482:MET:HB3	3:C:262:MET:HE1	1.99	0.45
3:C:175:ALA:HB2	3:C:284:MET:HE3	1.99	0.43
3:C:85:ILE:HD13	3:C:241:VAL:HG11	2.01	0.43
3:C:149:LEU:HD13	3:C:165:LEU:HB2	2.02	0.42
1:A:489:ALA:HB1	1:A:493:ILE:HD12	2.01	0.42
1:A:514:TYR:HB3	1:A:515:PRO:HD2	2.02	0.42
1:A:609:THR:HG23	2:B:399:TYR:CE1	2.54	0.42
1:A:620:THR:HG23	2:B:391:LEU:HD23	2.02	0.41
1:A:611:SER:HB2	3:C:160:ILE:HG21	2.02	0.41
3:C:285:ASP:CG	3:C:288:ARG:HH21	2.28	0.41
1:A:516:LEU:HD11	2:B:319:MET:HG2	2.04	0.40
2:B:370:LEU:HD23	3:C:37:VAL:HG12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	133/179 (74%)	128 (96%)	5 (4%)	0	100	100
2	B	100/113 (88%)	97 (97%)	3 (3%)	0	100	100
3	C	314/323 (97%)	310 (99%)	4 (1%)	0	100	100
All	All	547/615 (89%)	535 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	121/162 (75%)	121 (100%)	0	100	100
2	B	94/108 (87%)	94 (100%)	0	100	100
3	C	283/297 (95%)	271 (96%)	12 (4%)	26	40
All	All	498/567 (88%)	486 (98%)	12 (2%)	43	62

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	15	GLN
3	C	20	GLU
3	C	88	ILE
3	C	109	LEU
3	C	110	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	159	GLU
3	C	165	LEU
3	C	185	LYS
3	C	216	THR
3	C	249	ILE
3	C	285	ASP
3	C	318	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	574	GLN
3	C	15	GLN
3	C	60	ASN
3	C	127	HIS
3	C	192	ASN
3	C	229	ASN
3	C	286	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	AMP	C	323	-	25,25,25	1.46	4 (16%)	37,38,38	1.89	8 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	AMP	C	323	-	-	0/10/26/26	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	323	AMP	C5-C4	4.88	1.47	1.39
4	C	323	AMP	C5-C6	2.80	1.48	1.41
4	C	323	AMP	C8-N7	2.40	1.36	1.31
4	C	323	AMP	C5-N7	-2.19	1.35	1.39

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	C	323	AMP	C5-C4-N3	-5.61	119.00	126.72
4	C	323	AMP	N3-C4-N9	4.45	134.74	127.17
4	C	323	AMP	C2-N3-C4	3.77	121.04	111.83
4	C	323	AMP	N3-C2-N1	-3.55	123.21	128.58
4	C	323	AMP	C4-C5-N7	-3.39	106.70	110.58
4	C	323	AMP	C4-N9-C8	2.65	108.52	105.74
4	C	323	AMP	C5-N7-C8	2.45	107.31	103.45
4	C	323	AMP	C6-C5-N7	2.30	136.52	132.09

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand AMP C 323



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å^2)	Q<0.9
1	A	139/179 (77%)	1.15	24 (17%)	4 4	34, 54, 70, 73	0
2	B	104/113 (92%)	1.58	38 (36%)	1 1	32, 59, 75, 78	0
3	C	316/323 (97%)	0.84	44 (13%)	6 7	25, 40, 62, 65	0
All	All	559/615 (90%)	1.05	106 (18%)	3 4	25, 47, 69, 78	0

All (106) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	369	ALA	5.9
2	B	368	GLY	5.6
3	C	121	LEU	5.5
2	B	355	TYR	5.3
3	C	218	GLY	5.1
3	C	253	LEU	5.0
1	A	577	THR	4.9
1	A	562	ILE	4.9
3	C	122	ASP	4.8
1	A	609	THR	4.5
1	A	608	SER	4.4
2	B	337	TRP	4.4
3	C	123	THR	4.3
3	C	6	ASP	4.2
3	C	252	ASP	4.1
3	C	119	ASP	4.0
2	B	335	THR	4.0
2	B	362	PHE	4.0
2	B	371	PRO	3.9
1	A	550	TYR	3.8
1	A	493	ILE	3.7
2	B	327	ASP	3.7
3	C	126	ILE	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	155	GLU	3.5
2	B	359	GLN	3.5
2	B	361	GLN	3.5
2	B	303	MET	3.5
2	B	326	LEU	3.5
3	C	217	GLN	3.5
2	B	352	LEU	3.3
3	C	249	ILE	3.3
2	B	358	THR	3.3
2	B	365	ASN	3.2
3	C	157	HIS	3.1
2	B	367	SER	3.1
1	A	490	ALA	3.1
3	C	160	ILE	3.0
3	C	251	ASN	3.0
2	B	375	HIS	3.0
2	B	413	GLU	3.0
2	B	363	ASN	2.9
3	C	254	SER	2.9
1	A	591	SER	2.9
2	B	356	TYR	2.9
3	C	154	GLU	2.8
3	C	120	GLN	2.8
3	C	250	TYR	2.8
1	A	578	ASN	2.7
3	C	195	THR	2.6
1	A	564	ASP	2.6
1	A	502	LYS	2.6
1	A	488	PRO	2.6
3	C	216	THR	2.6
1	A	497	VAL	2.6
2	B	350	VAL	2.6
2	B	323	TYR	2.6
2	B	339	THR	2.5
3	C	98	PHE	2.5
3	C	153	ASP	2.5
2	B	372	ILE	2.5
3	C	255	LEU	2.5
2	B	366	ASN	2.5
1	A	499	LYS	2.4
3	C	74	ARG	2.4
2	B	347	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	322	TYR	2.4
3	C	169	ARG	2.4
2	B	354	LYS	2.4
3	C	213	GLN	2.4
2	B	336	SER	2.4
2	B	353	ASN	2.3
3	C	258	GLY	2.3
1	A	487	SER	2.3
3	C	124	ALA	2.3
3	C	12	SER	2.3
1	A	549	LYS	2.3
3	C	59	GLN	2.2
3	C	156	THR	2.2
3	C	289	LYS	2.2
3	C	13	ILE	2.2
2	B	321	ARG	2.2
1	A	491	SER	2.2
3	C	158	ARG	2.2
3	C	75	PHE	2.2
1	A	485	GLN	2.2
3	C	321	SER	2.2
1	A	548	TRP	2.1
2	B	391	LEU	2.1
2	B	360	ASP	2.1
2	B	348	GLU	2.1
3	C	212	ILE	2.1
3	C	159	GLU	2.1
3	C	198	ASN	2.1
3	C	68	TRP	2.1
1	A	610	PHE	2.1
1	A	630	SER	2.1
3	C	129	SER	2.1
1	A	492	LYS	2.1
2	B	324	TYR	2.1
2	B	389	ASN	2.1
1	A	500	LYS	2.0
1	A	479	ARG	2.0
3	C	92	PHE	2.0
2	B	306	THR	2.0
2	B	364	GLU	2.0
3	C	206	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	AMP	C	323	23/23	0.94	0.07	45,45,49,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around AMP C 323:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.