Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2026 – 10:09 PM JST

PDB ID	:	44YD / pdb_000044yd
Title	:	Crystal structure of D-lactate dehydrogenase from Lacticaseibacillus rhamnosus in its apoform
Authors	:	Matyuta, I.O.; Pometun, A.A.; Loginova, A.A.; Popov, V.O.; Boyko, K.M.
Deposited on	:	2026-08-17
Resolution	:	2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	341	6% 83% 14%
1	B	341	5% 77% 17%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 5210 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called D-lactate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	332	Total	C	N	O	S	0	0	0
			2553	1637	418	488	10			
1	B	330	Total	C	N	O	S	0	2	0
			2548	1644	411	483	10			

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP A0A7S7JGB1
A	2	GLY	-	expression tag	UNP A0A7S7JGB1
A	3	MET	-	expression tag	UNP A0A7S7JGB1
A	4	HIS	-	expression tag	UNP A0A7S7JGB1
A	5	HIS	-	expression tag	UNP A0A7S7JGB1
A	6	HIS	-	expression tag	UNP A0A7S7JGB1
A	7	HIS	-	expression tag	UNP A0A7S7JGB1
A	8	HIS	-	expression tag	UNP A0A7S7JGB1
A	9	HIS	-	expression tag	UNP A0A7S7JGB1
A	234	SER	GLY	conflict	UNP A0A7S7JGB1
B	1	MET	-	initiating methionine	UNP A0A7S7JGB1
B	2	GLY	-	expression tag	UNP A0A7S7JGB1
B	3	MET	-	expression tag	UNP A0A7S7JGB1
B	4	HIS	-	expression tag	UNP A0A7S7JGB1
B	5	HIS	-	expression tag	UNP A0A7S7JGB1
B	6	HIS	-	expression tag	UNP A0A7S7JGB1
B	7	HIS	-	expression tag	UNP A0A7S7JGB1
B	8	HIS	-	expression tag	UNP A0A7S7JGB1
B	9	HIS	-	expression tag	UNP A0A7S7JGB1
B	234	SER	GLY	conflict	UNP A0A7S7JGB1

- Molecule 2 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 7 4 3	0	0
2	A	1	Total C O 7 4 3	0	0
2	B	1	Total C O 7 4 3	0	0
2	B	1	Total C O 7 4 3	0	0

- Molecule 3 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O 10 6 4	0	0

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 6 3 3	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	27	Total O 27 27	0	0
5	B	38	Total O 38 38	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: D-lactate dehydrogenase

Chain A:



• Molecule 1: D-lactate dehydrogenase

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	65.33Å 80.25Å 70.22Å 90.00° 110.85° 90.00°	Depositor
Resolution (Å)	48.59 – 2.10 48.59 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.2 (48.59-2.10) 99.2 (48.59-2.10)	Depositor EDS
R_{merge}	0.37	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.33 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.249 , 0.324 0.249 , 0.324	Depositor DCC
R_{free} test set	1945 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å^2)	29.4	Xtriage
Anisotropy	0.302	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 46.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.51$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	5210	wwPDB-VP
Average B, all atoms (Å^2)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.64% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, PEG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	1.34	6/2612 (0.2%)	1.71	16/3556 (0.4%)
1	B	1.32	7/2614 (0.3%)	1.69	22/3558 (0.6%)
All	All	1.33	13/5226 (0.2%)	1.70	38/7114 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	94	ALA	C-O	7.62	1.33	1.24
1	A	238	VAL	C-O	6.93	1.31	1.24
1	B	194	PHE	C-O	6.84	1.31	1.23
1	A	174	GLY	C-O	-6.49	1.15	1.23
1	A	184	PRO	C-O	-6.34	1.16	1.24
1	A	149	LYS	C-O	5.82	1.30	1.23
1	B	180	ILE	C-O	5.74	1.30	1.24
1	B	56	ILE	C-O	5.70	1.30	1.24
1	B	208	ILE	C-O	5.64	1.29	1.24
1	B	157	GLY	C-O	5.19	1.30	1.23
1	A	161	THR	C-O	5.17	1.30	1.24
1	A	152	GLY	C-O	-5.15	1.17	1.24
1	B	265	ILE	C-O	5.05	1.29	1.24

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	161	THR	CA-C-N	9.66	131.03	122.43
1	A	161	THR	C-N-CA	9.66	131.03	122.43
1	A	66	ALA	CA-C-N	6.47	127.17	119.98
1	A	66	ALA	C-N-CA	6.47	127.17	119.98
1	B	69	PHE	CB-CA-C	6.45	121.09	110.90
1	B	8	HIS	CB-CA-C	6.38	121.48	110.45
1	B	66	ALA	CA-C-N	6.29	126.97	119.98
1	B	66	ALA	C-N-CA	6.29	126.97	119.98
1	A	325	ASP	CA-CB-CG	6.19	118.79	112.60
1	A	28	ALA	CA-C-N	6.17	128.87	120.54
1	A	28	ALA	C-N-CA	6.17	128.87	120.54
1	B	103	SER	CA-C-N	6.01	132.25	122.81
1	B	103	SER	C-N-CA	6.01	132.25	122.81
1	A	8	HIS	CB-CA-C	5.79	119.42	109.51
1	B	170	LYS	N-CA-C	-5.73	105.12	111.36
1	B	170	LYS	CA-C-O	-5.66	114.42	120.42
1	B	102	LEU	CA-C-O	-5.62	114.20	120.66
1	B	190	ASP	CB-CA-C	5.61	119.00	109.80
1	A	335	GLU	CB-CG-CD	5.52	121.98	112.60
1	A	155	THR	CA-CB-OG1	-5.45	101.42	109.60
1	B	168	ALA	CA-C-N	5.45	127.63	120.60
1	B	168	ALA	C-N-CA	5.45	127.63	120.60
1	A	10	LYS	CA-C-O	-5.44	114.45	120.38
1	B	318	PHE	CA-C-O	-5.42	114.68	120.42
1	B	285	LYS	N-CA-C	-5.41	103.92	111.39
1	B	225	GLU	CB-CA-C	5.37	119.40	110.81
1	A	310	THR	CA-CB-OG1	-5.34	101.59	109.60
1	B	310	THR	CA-CB-OG1	-5.28	101.68	109.60
1	A	73	HIS	N-CA-C	-5.27	105.43	111.07
1	A	336	VAL	CA-C-O	-5.24	115.79	121.92
1	A	161	THR	O-C-N	5.17	129.47	122.59
1	B	163	HIS	N-CA-C	-5.17	104.91	111.11
1	B	48	GLU	CA-C-O	-5.16	115.06	120.63
1	A	69	PHE	CB-CA-C	5.12	119.56	110.85
1	B	249	GLN	O-C-N	5.06	127.28	122.07
1	B	248	THR	N-CA-C	-5.06	105.86	112.23
1	B	186	PRO	CA-C-O	-5.05	115.58	121.34
1	B	218	GLN	CB-CA-C	-5.00	102.17	110.68

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	281	HIS	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2553	0	2441	18	0
1	B	2548	0	2452	26	0
2	A	14	0	20	0	0
2	B	14	0	20	0	0
3	B	10	0	14	0	0
4	B	6	0	8	0	0
5	A	27	0	0	1	0
5	B	38	0	0	2	0
All	All	5210	0	4955	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:89:ASN:H	1:B:89:ASN:HD22	1.33	0.74
1:B:11:ILE:HG12	1:B:323:LEU:HD13	1.83	0.60
1:B:80:LEU:C	1:B:80:LEU:HD23	2.29	0.56
1:B:158:VAL:HG21	1:B:169:ILE:HG13	1.91	0.53
1:A:42:LEU:HD22	1:A:56:ILE:HG21	1.90	0.53
1:B:102:LEU:CD2	1:B:336:VAL:HG23	2.39	0.52
1:B:80:LEU:HD23	1:B:80:LEU:O	2.11	0.50
1:B:191:HIS:HB2	1:B:194:PHE:CE1	2.46	0.50
1:B:81:THR:HG22	1:B:103:SER:OG	2.14	0.48
1:B:80:LEU:O	1:B:102:LEU:HA	2.13	0.48
1:A:215:GLY:CA	1:A:242:ARG:HG3	2.44	0.47
1:B:263:VAL:HG22	1:B:298:VAL:HG22	1.95	0.47
1:A:284:PHE:O	1:A:286:ASP:N	2.48	0.47
1:B:166:ARG:O	1:B:169:ILE:HB	2.15	0.47
1:A:21:ILE:HD12	1:A:37:TYR:CE2	2.50	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:13:ALA:HA	1:B:57:ASN:O	2.15	0.46
1:B:231:MET:O	1:B:259:LYS:CE	2.63	0.46
1:A:13:ALA:HA	1:A:57:ASN:O	2.16	0.46
1:A:21:ILE:HG22	1:B:30:GLU:HA	1.98	0.46
1:A:134:GLN:OE1	1:A:134:GLN:HA	2.16	0.45
1:B:134:GLN:HA	1:B:134:GLN:OE1	2.16	0.45
1:A:215:GLY:N	1:A:242:ARG:HG3	2.32	0.45
1:B:113:ILE:HG21	1:B:164:ILE:HG21	1.98	0.45
1:B:89:ASN:HD22	1:B:89:ASN:N	2.07	0.45
1:A:24:PHE:CD1	1:A:35:LEU:HD21	2.51	0.44
1:A:38:HIS:ND1	5:A:503:HOH:O	2.36	0.44
1:B:129:GLY:N	5:B:508:HOH:O	2.43	0.43
1:A:181:ALA:O	1:A:196:TYR:HA	2.17	0.43
1:B:158:VAL:HG21	1:B:169:ILE:CG1	2.46	0.43
1:B:191:HIS:HD2	1:B:194:PHE:O	2.01	0.43
1:B:24:PHE:CD1	1:B:35:LEU:HD21	2.54	0.42
1:B:108:TYR:CD2	1:B:109:SER:N	2.88	0.42
1:A:73:HIS:HB2	1:A:98:TYR:HB3	2.00	0.42
1:B:200:GLU:HG2	1:B:230:LEU:HD13	2.01	0.42
1:B:80:LEU:HD21	1:B:82:ILE:HG23	2.01	0.42
1:B:127:ASN:HA	5:B:538:HOH:O	2.20	0.41
1:A:80:LEU:O	1:A:102:LEU:HA	2.19	0.41
1:B:73:HIS:HB2	1:B:98:TYR:HB3	2.02	0.40
1:A:169:ILE:HG23	1:A:179:VAL:HG11	2.03	0.40
1:A:235:VAL:HG13	1:A:260:LEU:HD23	2.02	0.40
1:A:215:GLY:HA3	1:A:242:ARG:HG3	2.03	0.40
1:A:267:THR:HG22	1:A:303:HIS:CE1	2.57	0.40
1:A:267:THR:HG22	1:A:303:HIS:ND1	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	330/341 (97%)	309 (94%)	19 (6%)	2 (1%)	21 18
1	B	330/341 (97%)	317 (96%)	9 (3%)	4 (1%)	10 7
All	All	660/682 (97%)	626 (95%)	28 (4%)	6 (1%)	14 10

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	285	LYS
1	B	283	SER
1	B	280	LYS
1	B	66	ALA
1	B	241	ALA
1	A	286	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	261/284 (92%)	256 (98%)	5 (2%)	50 58
1	B	261/284 (92%)	249 (95%)	12 (5%)	24 25
All	All	522/568 (92%)	505 (97%)	17 (3%)	33 37

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	103	SER
1	A	204	LYS
1	A	259	LYS
1	A	277	ASN
1	A	280	LYS
1	B	8	HIS
1	B	80	LEU
1	B	89	ASN
1	B	92	MET
1	B	102	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	104	ASN
1	B	178	LYS
1	B	204	LYS
1	B	276	LEU
1	B	281	HIS
1	B	323	LEU
1	B	329	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	HIS
1	A	96	ASN
1	A	153	GLN
1	B	89	ASN
1	B	191	HIS
1	B	212	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	PEG	A	402	-	6,6,6	0.40	0	5,5,5	0.41	0
4	GOL	B	403	-	5,5,5	0.22	0	5,5,5	0.79	0
2	PEG	B	404	-	6,6,6	0.22	0	5,5,5	0.52	0
2	PEG	B	401	-	6,6,6	0.75	0	5,5,5	0.57	0
3	PGE	B	402	-	9,9,9	0.34	0	8,8,8	0.45	0
2	PEG	A	401	-	6,6,6	0.96	0	5,5,5	0.79	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	A	402	-	-	1/4/4/4	-
4	GOL	B	403	-	-	4/4/4/4	-
2	PEG	B	404	-	-	3/4/4/4	-
2	PEG	B	401	-	-	2/4/4/4	-
3	PGE	B	402	-	-	4/7/7/7	-
2	PEG	A	401	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	PEG	C1-C2-O2-C3
2	A	401	PEG	O2-C3-C4-O4
2	B	401	PEG	O2-C3-C4-O4
3	B	402	PGE	O2-C3-C4-O3
4	B	403	GOL	O1-C1-C2-C3
4	B	403	GOL	C1-C2-C3-O3
4	B	403	GOL	O2-C2-C3-O3
3	B	402	PGE	O1-C1-C2-O2
2	A	401	PEG	O1-C1-C2-O2
2	B	404	PEG	C1-C2-O2-C3
3	B	402	PGE	C3-C4-O3-C5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	404	PEG	C4-C3-O2-C2
2	B	404	PEG	O2-C3-C4-O4
3	B	402	PGE	C6-C5-O3-C4
4	B	403	GOL	O1-C1-C2-O2
2	A	401	PEG	C4-C3-O2-C2
2	A	402	PEG	C4-C3-O2-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	332/341 (97%)	0.60	20 (6%)	27	29	19, 35, 64, 95	0
1	B	330/341 (96%)	0.71	16 (4%)	35	38	14, 35, 54, 76	2 (0%)
All	All	662/682 (97%)	0.66	36 (5%)	31	33	14, 35, 57, 95	2 (0%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	279	ALA	5.4
1	B	336	VAL	4.8
1	A	6	HIS	4.4
1	A	283	SER	4.1
1	A	278	LEU	4.1
1	A	185	TYR	3.8
1	B	108	TYR	3.7
1	B	102	LEU	3.5
1	A	287	PRO	3.4
1	A	7	HIS	3.2
1	B	225	GLU	3.2
1	B	279	ALA	3.2
1	B	276	LEU	3.1
1	A	32	GLY	3.0
1	B	202	LEU	2.8
1	A	288	LEU	2.8
1	B	281	HIS	2.7
1	A	282	GLY	2.7
1	A	284	PHE	2.6
1	B	288	LEU	2.6
1	A	285	LYS	2.5
1	B	280	LYS	2.4
1	A	45	HIS	2.4
1	A	8	HIS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	310	THR	2.4
1	B	7	HIS	2.3
1	A	281	HIS	2.3
1	A	188	LYS	2.3
1	B	278	LEU	2.3
1	B	282	GLY	2.2
1	A	184	PRO	2.2
1	B	105	VAL	2.2
1	A	294	ALA	2.2
1	A	253	SER	2.1
1	B	27	TRP	2.0
1	B	157	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	PEG	A	402	7/7	0.76	0.16	42,50,60,66	0
2	PEG	B	404	7/7	0.80	0.15	48,51,58,58	0
2	PEG	A	401	7/7	0.83	0.16	39,43,48,48	0
3	PGE	B	402	10/10	0.85	0.14	33,48,52,62	0
2	PEG	B	401	7/7	0.86	0.13	27,35,45,46	0
4	GOL	B	403	6/6	0.90	0.09	41,45,46,46	0

6.5 Other polymers (i)

There are no such residues in this entry.