Full wwPDB X-ray Structure Validation Report (i)

Nov 18, 2026 – 04:15 PM JST

PDB ID	:	44YZ / pdb_000044yz
Title	:	Human ADH6-NAD-1PO protein
Authors	:	Ding, J.; Zhong, Y.; Chen, Z.
Deposited on	:	2026-08-18
Resolution	:	1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	374	2% 94% 6%
1	B	374	5% 89% 10%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 6149 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Isoform 2 of Alcohol dehydrogenase 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	374	Total	C	N	O	S	0	0	0
			2773	1769	461	519	24			
1	B	374	Total	C	N	O	S	0	0	0
			2773	1769	461	519	24			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 4 is N-PROPANOL (CCD ID: POL) (formula: C_3H_8O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	3	1		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	306	Total O 306 306	0	0
6	B	193	Total O 193 193	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Isoform 2 of Alcohol dehydrogenase 6

Chain A:



• Molecule 1: Isoform 2 of Alcohol dehydrogenase 6

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	63.66Å 45.58Å 115.23Å 90.00° 91.90° 90.00°	Depositor
Resolution (Å)	42.38 – 1.90 42.38 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.9 (42.38-1.90) 96.9 (42.38-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.39 (at 1.91Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.174 , 0.210 0.174 , 0.210	Depositor DCC
R_{free} test set	1977 reflections (3.77%)	wwPDB-VP
Wilson B-factor (Å^2)	24.2	Xtriage
Anisotropy	0.357	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 39.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.024 for h,-k,-l	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	6149	wwPDB-VP
Average B, all atoms (Å^2)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.39% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, NAD, EDO, POL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.19	0/2817	0.41	0/3804
1	B	0.17	0/2817	0.43	0/3804
All	All	0.18	0/5634	0.42	0/7608

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2773	0	2862	13	0
1	B	2773	0	2862	31	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	44	0	26	1	0
3	B	44	0	26	4	0
4	A	4	0	7	0	0
5	A	4	0	6	0	0
5	B	4	0	6	0	0
6	A	306	0	0	4	0
6	B	193	0	0	0	0
All	All	6149	0	5795	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:19:GLY:HA3	1:B:57:LYS:HE3	1.74	0.68
1:B:166:PRO:HD2	1:B:337:MET:HE1	1.80	0.64
1:B:11:LYS:HE3	1:B:353:LEU:HD23	1.78	0.64
1:A:11:LYS:NZ	6:A:503:HOH:O	2.27	0.63
1:B:51:MET:HE1	1:B:360:VAL:HA	1.79	0.63
1:B:227:LYS:H	1:B:227:LYS:HZ3	1.44	0.63
1:B:51:MET:HE3	1:B:363:MET:HB2	1.80	0.62
1:A:118:GLN:NE2	6:A:501:HOH:O	2.21	0.59
1:B:56:SER:HB3	1:B:59:LEU:HB2	1.85	0.58
1:B:149:GLU:HG2	1:B:375:LEU:HD12	1.87	0.57
1:A:331:LYS:NZ	6:A:512:HOH:O	2.39	0.56
1:B:295:VAL:HG13	3:B:403:NAD:H1D	1.88	0.54
1:B:239:THR:C	1:B:240:GLU:HG3	2.33	0.54
1:A:9:ARG:NH1	6:A:506:HOH:O	2.35	0.53
1:A:175:CYS:SG	3:A:403:NAD:H5N	2.49	0.52
1:B:254:VAL:O	1:B:258:MET:HG3	2.11	0.50
1:B:51:MET:HE3	1:B:363:MET:CB	2.42	0.49
1:B:18:PRO:HG3	1:B:63:TYR:CE1	2.48	0.48
1:B:175:CYS:SG	3:B:403:NAD:H5N	2.53	0.48
1:B:295:VAL:H	3:B:403:NAD:H1D	1.78	0.47
1:A:124:MET:HE3	1:A:124:MET:HB3	1.75	0.47
1:B:52:LYS:O	1:B:56:SER:HB2	2.14	0.47
1:B:206:LEU:O	1:B:210:MET:HG3	2.15	0.46
1:B:300:VAL:HG12	1:B:301:GLN:H	1.81	0.46
1:A:164:VAL:HG23	1:A:337:MET:HE2	1.98	0.45
1:B:114:ILE:HD12	1:B:115:GLN:HG2	1.98	0.45
1:B:51:MET:HE3	1:B:363:MET:HG3	1.99	0.44
1:B:46:LEU:HG	1:B:51:MET:HE2	1.98	0.44
1:B:114:ILE:HD12	1:B:115:GLN:N	2.32	0.43
1:B:268:GLU:CD	1:B:271:GLY:H	2.26	0.43
1:B:303:LYS:HB2	1:B:303:LYS:NZ	2.33	0.43
1:A:219:ARG:HD2	1:A:240:GLU:OE1	2.19	0.43
1:B:355:LYS:HG2	1:B:358:GLU:OE2	2.19	0.43
1:A:33:LYS:HD2	1:A:130:ARG:CZ	2.49	0.42
1:B:166:PRO:CD	1:B:337:MET:HE1	2.48	0.42
1:B:363:MET:HE3	1:B:363:MET:HB3	1.96	0.41
1:B:200:GLY:O	1:B:205:GLY:HA3	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:188:ALA:HB2	1:A:291:VAL:HG21	2.03	0.41
1:A:200:GLY:O	1:A:205:GLY:HA3	2.20	0.41
1:B:227:LYS:H	1:B:227:LYS:NZ	2.17	0.41
1:B:179:THR:HG21	3:B:403:NAD:C4N	2.51	0.41
1:B:300:VAL:HG12	1:B:301:GLN:N	2.36	0.41
1:A:56:SER:HB2	1:A:59:LEU:HG	2.03	0.40
1:A:18:PRO:HG3	1:A:63:TYR:CE1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	372/374 (100%)	357 (96%)	14 (4%)	1 (0%)	36 29
1	B	372/374 (100%)	358 (96%)	14 (4%)	0	100 100
All	All	744/748 (100%)	715 (96%)	28 (4%)	1 (0%)	48 40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	260	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	306/306 (100%)	304 (99%)	2 (1%)	76 78
1	B	306/306 (100%)	302 (99%)	4 (1%)	61 61
All	All	612/612 (100%)	606 (99%)	6 (1%)	68 70

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	HIS
1	A	234	GLN
1	B	56	SER
1	B	58	HIS
1	B	295	VAL
1	B	304	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	107	ASN
1	A	307	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	POL	A	404	2	3,3,3	0.32	0	2,2,2	0.25	0
3	NAD	A	403	-	45,48,48	1.77	9 (20%)	63,73,73	1.86	10 (15%)
5	EDO	A	405	-	3,3,3	0.48	0	2,2,2	0.35	0
5	EDO	B	404	-	3,3,3	0.52	0	2,2,2	0.17	0
3	NAD	B	403	-	45,48,48	1.73	10 (22%)	63,73,73	1.89	9 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	POL	A	404	2	-	1/1/1/1	-
3	NAD	A	403	-	-	5/30/62/62	0/5/5/5
5	EDO	A	405	-	-	1/1/1/1	-
5	EDO	B	404	-	-	1/1/1/1	-
3	NAD	B	403	-	-	6/30/62/62	0/5/5/5

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	403	NAD	C3N-C7N	-6.28	1.41	1.50
3	B	403	NAD	C3N-C7N	-6.03	1.41	1.50
3	A	403	NAD	C8A-N7A	4.44	1.39	1.31
3	B	403	NAD	C8A-N7A	4.42	1.39	1.31
3	B	403	NAD	C5N-C4N	-3.48	1.31	1.38
3	A	403	NAD	C2N-C3N	-3.48	1.33	1.39
3	B	403	NAD	C2N-C3N	-3.46	1.33	1.39
3	A	403	NAD	C5N-C4N	-3.26	1.32	1.38
3	A	403	NAD	C2A-N1A	3.12	1.39	1.33
3	B	403	NAD	C2A-N1A	2.95	1.39	1.33
3	A	403	NAD	O4D-C1D	2.84	1.45	1.41
3	B	403	NAD	O4D-C1D	2.49	1.44	1.41
3	A	403	NAD	C2A-N3A	2.35	1.38	1.33
3	B	403	NAD	C2A-N3A	2.35	1.38	1.33
3	A	403	NAD	C2N-N1N	-2.33	1.32	1.35
3	B	403	NAD	C4A-N3A	2.18	1.38	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	403	NAD	C2N-N1N	-2.15	1.32	1.35
3	A	403	NAD	C4A-N3A	2.12	1.38	1.34
3	B	403	NAD	C2D-C1D	2.09	1.56	1.53

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	B	403	NAD	O7N-C7N-N7N	-6.36	113.55	122.58
3	A	403	NAD	O7N-C7N-N7N	-6.10	113.92	122.58
3	B	403	NAD	C3N-C7N-N7N	6.01	124.97	117.75
3	A	403	NAD	C3N-C7N-N7N	5.28	124.09	117.75
3	B	403	NAD	O4D-C1D-C2D	-5.15	99.41	106.93
3	A	403	NAD	O4D-C1D-C2D	-4.70	100.05	106.93
3	B	403	NAD	C6N-N1N-C2N	-4.15	118.19	121.97
3	A	403	NAD	N3A-C2A-N1A	-4.13	122.14	128.60
3	A	403	NAD	C6N-N1N-C2N	-4.01	118.32	121.97
3	B	403	NAD	N3A-C2A-N1A	-3.97	122.39	128.60
3	A	403	NAD	O4B-C4B-C3B	3.34	111.73	105.11
3	B	403	NAD	O4B-C4B-C3B	3.21	111.46	105.11
3	B	403	NAD	C5A-C4A-N9A	3.19	109.50	105.78
3	A	403	NAD	C5A-C4A-N9A	3.19	109.50	105.78
3	A	403	NAD	PN-O3-PA	-3.16	121.97	132.83
3	B	403	NAD	PN-O3-PA	-2.84	123.07	132.83
3	B	403	NAD	C6A-C5A-C4A	2.29	120.26	117.18
3	A	403	NAD	C4B-O4B-C1B	-2.23	104.55	109.47
3	A	403	NAD	C6A-C5A-C4A	2.17	120.11	117.18

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	403	NAD	O4D-C1D-N1N-C2N
3	A	403	NAD	O4D-C1D-N1N-C6N
3	A	403	NAD	C2D-C1D-N1N-C2N
3	A	403	NAD	C2D-C1D-N1N-C6N
3	B	403	NAD	O4D-C1D-N1N-C2N
3	B	403	NAD	O4D-C1D-N1N-C6N
3	B	403	NAD	C2D-C1D-N1N-C2N
3	B	403	NAD	C2D-C1D-N1N-C6N
5	A	405	EDO	O1-C1-C2-O2
3	B	403	NAD	O4B-C4B-C5B-O5B
5	B	404	EDO	O1-C1-C2-O2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	403	NAD	O4B-C4B-C5B-O5B
4	A	404	POL	O-C1-C2-C3
3	B	403	NAD	C3B-C4B-C5B-O5B

There are no ring outliers.


2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	403	NAD	1	0
3	B	403	NAD	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand NAD A 403




Bond lengths



Bond angles




Torsions



Rings

Ligand NAD B 403



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	374/374 (100%)	-0.17	6 (1%)	70	74	14, 23, 44, 62	0
1	B	374/374 (100%)	0.28	20 (5%)	32	34	18, 30, 53, 87	0
All	All	748/748 (100%)	0.05	26 (3%)	47	50	14, 27, 49, 87	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	299	SER	4.8
1	B	59	LEU	4.2
1	B	300	VAL	4.1
1	A	375	LEU	3.8
1	B	302	LEU	3.2
1	B	298	ALA	3.1
1	B	295	VAL	3.1
1	A	260	ASP	3.0
1	A	56	SER	2.9
1	B	260	ASP	2.7
1	B	261	ALA	2.6
1	A	247	LEU	2.6
1	B	273	LEU	2.6
1	B	297	PRO	2.5
1	B	375	LEU	2.5
1	B	347	ILE	2.5
1	B	58	HIS	2.4
1	B	56	SER	2.4
1	A	58	HIS	2.4
1	B	55	GLY	2.4
1	B	296	LEU	2.3
1	B	338	ALA	2.2
1	A	303	LYS	2.1
1	B	341	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	343	LEU	2.1
1	B	2	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	NAD	B	403	44/44	0.90	0.09	28,38,42,44	0
4	POL	A	404	4/4	0.92	0.10	20,21,23,31	0
5	EDO	B	404	4/4	0.96	0.07	22,22,26,27	0
5	EDO	A	405	4/4	0.97	0.09	17,19,22,24	0
3	NAD	A	403	44/44	0.97	0.06	16,22,29,32	0
2	ZN	B	402	1/1	0.98	0.03	26,26,26,26	1
2	ZN	A	402	1/1	0.99	0.02	19,19,19,19	0
2	ZN	B	401	1/1	0.99	0.03	23,23,23,23	0
2	ZN	A	401	1/1	1.00	0.01	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around NAD B 403:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around POL A 404:


$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around NAD A 403:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)









Electron density around ZN B 402:


$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around ZN A 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around ZN B 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)








Electron density around ZN A 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers

There are no such residues in this entry.