



# Full wwPDB X-ray Structure Validation Report i

Jun 24, 2025 – 05:46 pm BST

PDB ID : 4ASR / pdb\_00004asr  
Title : Crystal structure of ANCE in complex with Thr6-Bradykinin  
Authors : Akif, M.; Masuyer, G.; Sturrock, E.D.; Isaac, R.E.; Acharya, K.R.  
Deposited on : 2012-05-02  
Resolution : 1.90 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.44


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 5600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ANGIOTENSIN-CONVERTING ENZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	598	Total	C	N	O	S	0	2	0


- Molecule 2 is a protein called BRADYKININ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	P	4	Total	C	N	O		0	0	1

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
P	6	THR	SER	engineered mutation	UNP ?

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	B	6	Total	C	N	O		0	0	0

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N O 14 8 1 5	0	0
5	A	1	Total C N O 14 8 1 5	0	0

- Molecule 6 is CITRATE ANION (CCD ID: FLC) (formula: C<sub>6</sub>H<sub>5</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 13 6 7	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	572	Total O 572 572	0	0
7	P	1	Total O 1 1	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ANGIOTENSIN-CONVERTING ENZYME



- Molecule 2: BRADYKININ



- Molecule 3: beta-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics i

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	173.29 Å   173.29 Å   101.46 Å 90.00°   90.00°   120.00°	Depositor
Resolution (Å)	86.65 – 1.90 86.65 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.2 (86.65-1.90) 85.0 (86.65-1.90)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.54 (at 1.89 Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
$R$ , $R_{free}$	0.183, 0.200 0.183, 0.201	Depositor DCC
$R_{free}$ test set	3818 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.5	Xtriage
Anisotropy	0.852	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39, 40.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.009 for $-2/3^*h-1/3^*k-4/3^*l,-1/3^*h-2/3^*k+$ $4/3^*l,-1/3^*h+1/3^*k+1/3^*l$ 0.007 for $-h,1/3^*h-1/3^*k-4/3^*l,-1/3^*h-2/3^*k$ $+1/3^*l$ 0.007 for $-1/3^*h+1/3^*k+4/3^*l,-k,2/3^*h+1/$ $3^*k+1/3^*l$ 0.008 for $-h,2/3^*h+1/3^*k+4/3^*l,1/3^*h+2/3$ $*k-1/3^*l$ 0.013 for $-1/3^*h-2/3^*k+4/3^*l,-2/3^*h-1/3^*k-$ $4/3^*l,1/3^*h-1/3^*k-1/3^*l$ 0.011 for $1/3^*h+2/3^*k-4/3^*l,-k,-2/3^*h-1/3^*$ $k-1/3^*l$ 0.016 for $h,-h-k,-l$	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5600	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, NAG, FLC, MAN, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.43	0/5019	0.74	0/6800
2	P	1.49	1/27 (3.7%)	1.25	0/37
All	All	0.44	1/5046 (0.0%)	0.75	0/6837

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	P	3	PRO	C-N	-6.76	1.23	1.33

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4887	0	4688	31	0
2	P	26	0	29	0	0
3	B	72	0	61	0	0
4	A	1	0	0	0	0
5	A	28	0	26	0	0
6	A	13	0	5	0	0
7	A	572	0	0	1	0
7	P	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5600	0	4809	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:325:ILE:O	1:A:355:THR:HG23	1.76	0.85
1:A:277:PHE:H	1:A:429:ASN:HD21	1.29	0.78
1:A:377:GLN:HE22	1:A:553:HIS:HD2	1.31	0.75
1:A:285:VAL:H	1:A:359:GLN:HE21	1.46	0.63
1:A:313:THR:H	1:A:373:GLN:HE22	1.46	0.62
1:A:377:GLN:HE22	1:A:553:HIS:CD2	2.19	0.57
1:A:516:GLN:NE2	1:A:578:SER:H	2.05	0.55
1:A:347:ASP:H	1:A:379:GLN:HE22	1.53	0.55
1:A:300:PHE:HE1	1:A:355:THR:HG21	1.71	0.54
1:A:276:PRO:HB3	1:A:592:TRP:CH2	2.43	0.53
1:A:466:ASN:ND2	1:A:491:ASP:H	2.06	0.52
1:A:147:PRO:HB3	1:A:356:ARG:HD3	1.91	0.52
1:A:133:CYS:HA	1:A:141:CYS:HA	1.91	0.52
1:A:33:ASN:HD22	1:A:383:PHE:HB3	1.79	0.48
1:A:285:VAL:H	1:A:359:GLN:NE2	2.12	0.46
1:A:296:PRO:HG2	1:A:326:GLU:HG2	1.98	0.45
1:A:230:GLN:NE2	1:A:597:ASN:HD22	2.15	0.45
1:A:306:PHE:CD1	1:A:401:LEU:HD13	2.52	0.45
1:A:466:ASN:HD21	1:A:491:ASP:H	1.66	0.44
1:A:404:SER:HB2	1:A:541:ASP:HA	2.00	0.43
1:A:527:GLN:HE21	1:A:541:ASP:H	1.67	0.43
1:A:362:LEU:O	1:A:366:HIS:HD2	2.02	0.43
1:A:146:ASP:HA	1:A:147:PRO:HA	1.77	0.43
1:A:137:ASP:OD1	1:A:139:THR:HG22	2.19	0.42
1:A:285:VAL:HG22	1:A:359:GLN:HE21	1.84	0.42
1:A:527:GLN:NE2	1:A:541:ASP:H	2.18	0.42
1:A:229:GLN:HE22	1:A:606:TRP:HH2	1.68	0.41
1:A:355:THR:HA	1:A:361:GLN:OE1	2.20	0.41
1:A:113:TYR:CE2	1:A:117:LEU:HD11	2.54	0.41
1:A:603:HIS:HE1	7:A:2318:HOH:O	2.04	0.41
1:A:225:ARG:O	1:A:229:GLN:HG2	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	598/598 (100%)	592 (99%)	6 (1%)	0	100 100
2	P	2/9 (22%)	2 (100%)	0	0	100 100
All	All	600/607 (99%)	594 (99%)	6 (1%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	522/520 (100%)	516 (99%)	6 (1%)	70 71
2	P	3/8 (38%)	3 (100%)	0	100 100
All	All	525/528 (99%)	519 (99%)	6 (1%)	70 71

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	178	ARG
1	A	355	THR
1	A	378	TYR
1	A	394	HIS
1	A	401	LEU
1	A	446	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	ASN
1	A	84	GLN
1	A	126	ASN
1	A	229	GLN
1	A	230	GLN
1	A	359	GLN
1	A	366	HIS
1	A	373	GLN
1	A	377	GLN
1	A	379	GLN
1	A	380	HIS
1	A	381	GLN
1	A	429	ASN
1	A	466	ASN
1	A	516	GLN
1	A	527	GLN
1	A	553	HIS
1	A	601	ASN
1	A	603	HIS

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	B	1	1,3	14,14,15	0.49	0	17,19,21	0.59	0
3	NAG	B	2	3	14,14,15	0.49	0	17,19,21	1.25	2 (11%)
3	BMA	B	3	3	11,11,12	0.62	0	15,15,17	0.86	0
3	MAN	B	4	3	11,11,12	0.61	0	15,15,17	0.70	0
3	BMA	B	5	3	11,11,12	0.56	0	15,15,17	1.21	2 (13%)
3	MAN	B	6	3	11,11,12	0.62	0	15,15,17	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	B	2	3	-	3/6/23/26	0/1/1/1
3	BMA	B	3	3	-	1/2/19/22	0/1/1/1
3	MAN	B	4	3	-	2/2/19/22	0/1/1/1
3	BMA	B	5	3	-	2/2/19/22	0/1/1/1
3	MAN	B	6	3	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	B	5	BMA	C1-O5-C5	3.51	116.95	112.19
3	B	2	NAG	C2-N2-C7	3.19	127.45	122.90
3	B	2	NAG	C1-O5-C5	2.24	115.23	112.19
3	B	5	BMA	C1-C2-C3	2.03	112.16	109.67

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	4	MAN	C4-C5-C6-O6
3	B	4	MAN	O5-C5-C6-O6
3	B	5	BMA	O5-C5-C6-O6
3	B	2	NAG	C4-C5-C6-O6
3	B	5	BMA	C4-C5-C6-O6
3	B	2	NAG	O5-C5-C6-O6
3	B	2	NAG	C3-C2-N2-C7

*Continued on next page...*


Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	3	BMA	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
5	NAG	A	1622	-	14,14,15	0.57	0	17,19,21	0.62	0
5	NAG	A	1623	-	14,14,15	0.51	0	17,19,21	0.86	0
6	FLC	A	1624	-	12,12,12	1.00	0	17,17,17	1.30	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	1622	-	-	2/6/23/26	0/1/1/1
5	NAG	A	1623	-	-	0/6/23/26	0/1/1/1
6	FLC	A	1624	-	-	13/16/16/16	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
6	A	1624	FLC	OB2-CBC-CB	3.36	118.89	113.05
6	A	1624	FLC	OB1-CBC-CB	-2.06	119.34	122.25

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1624	FLC	CG-CB-CBC-OB1
6	A	1624	FLC	CG-CB-CBC-OB2
6	A	1624	FLC	OHB-CB-CBC-OB1
6	A	1624	FLC	OHB-CB-CBC-OB2
6	A	1624	FLC	CA-CB-CG-CGC
6	A	1624	FLC	CBC-CB-CG-CGC
6	A	1624	FLC	OHB-CB-CG-CGC
5	A	1622	NAG	O5-C5-C6-O6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
6	A	1624	FLC	CAC-CA-CB-CBC
6	A	1624	FLC	CAC-CA-CB-CG
5	A	1622	NAG	C4-C5-C6-O6
6	A	1624	FLC	CB-CA-CAC-OA2
6	A	1624	FLC	CB-CA-CAC-OA1
6	A	1624	FLC	CA-CB-CBC-OB2
6	A	1624	FLC	CAC-CA-CB-OHB

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [\(i\)](#)

### 6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	598/598 (100%)	0.31	34 (5%) 30 32	15, 25, 36, 43	9 (1%)
2	P	4/9 (44%)	0.46	0 100 100	24, 24, 25, 28	0
All	All	602/607 (99%)	0.31	34 (5%) 31 32	15, 25, 36, 43	9 (1%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	139	THR	6.7
1	A	614	SER	6.4
1	A	331	GLY	4.0
1	A	141	CYS	3.6
1	A	437	ASP	3.6
1	A	336	CYS	3.6
1	A	330	ASP	3.4
1	A	57	GLU	3.4
1	A	329	THR	3.3
1	A	17	ALA	3.3
1	A	136	LYS	3.2
1	A	415	GLY	3.2
1	A	531	ASP	3.1
1	A	360	ASP	2.8
1	A	30	GLU	2.8
1	A	467	CYS	2.8
1	A	64	GLU	2.4
1	A	534	GLU	2.4
1	A	20	LYS	2.3
1	A	461	ASP	2.3
1	A	404	SER	2.3
1	A	27	GLU	2.3
1	A	533	VAL	2.3
1	A	91	GLU	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	305	ASP	2.3
1	A	290	GLU	2.2
1	A	547	ALA	2.2
1	A	137	ASP	2.2
1	A	422	ARG	2.2
1	A	179	SER	2.1
1	A	61	LYS	2.1
1	A	140	LYS	2.1
1	A	544	GLY	2.1
1	A	59	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)


There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	B	1	14/15	-	-	20,22,23,26	0
3	NAG	B	2	14/15	-	-	29,32,34,36	0
3	BMA	B	3	11/12	-	-	40,43,45,48	0
3	MAN	B	4	11/12	-	-	45,46,48,49	0
3	BMA	B	5	11/12	-	-	50,51,51,52	0
3	MAN	B	6	11/12	-	-	50,51,52,52	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	NAG	A	1623	14/15	0.73	0.22	30,32,33,33	14
6	FLC	A	1624	13/13	0.73	0.26	54,55,56,56	0
5	NAG	A	1622	14/15	0.77	0.24	24,26,27,27	14
4	ZN	A	1615	1/1	0.99	0.03	21,21,21,21	0

## 6.5 Other polymers (i)

There are no such residues in this entry.