



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 4, 2025 – 08:25 AM EDT

PDB ID : 4AW7 / pdb_00004aw7
Title : BpGH86A: A beta-porphyrinase of glycoside hydrolase family 86 from the human gut bacterium *Bacteroides plebeius*
Authors : Hehemann, J.H.; Kelly, A.G.; Pudlo, N.A.; Martens, E.C.; Boraston, A.B.
Deposited on : 2012-06-01
Resolution : 1.33 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1904 (1.36-1.32)
Clashscore	180529	2038 (1.36-1.32)
Ramachandran outliers	177936	2016 (1.36-1.32)
Sidechain outliers	177891	2016 (1.36-1.32)
RSRZ outliers	164620	1903 (1.36-1.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	591	
2	B	6	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	EDO	A	1618	-	-	X	-

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 5264 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GH86A BETA-PORPHYRANASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	565	4563	2905	762	883	13	0	17	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	MET	-	expression tag	UNP B5CY96
A	8	GLY	-	expression tag	UNP B5CY96
A	9	SER	-	expression tag	UNP B5CY96
A	10	SER	-	expression tag	UNP B5CY96
A	11	HIS	-	expression tag	UNP B5CY96
A	12	HIS	-	expression tag	UNP B5CY96
A	13	HIS	-	expression tag	UNP B5CY96
A	14	HIS	-	expression tag	UNP B5CY96
A	15	HIS	-	expression tag	UNP B5CY96
A	16	HIS	-	expression tag	UNP B5CY96
A	17	SER	-	expression tag	UNP B5CY96
A	18	SER	-	expression tag	UNP B5CY96
A	19	GLY	-	expression tag	UNP B5CY96
A	20	LEU	-	expression tag	UNP B5CY96
A	21	VAL	-	expression tag	UNP B5CY96
A	22	PRO	-	expression tag	UNP B5CY96
A	23	ARG	-	expression tag	UNP B5CY96
A	24	GLY	-	expression tag	UNP B5CY96
A	25	SER	-	expression tag	UNP B5CY96
A	26	HIS	-	expression tag	UNP B5CY96
A	27	MET	-	expression tag	UNP B5CY96
A	28	ALA	-	expression tag	UNP B5CY96
A	29	SER	-	expression tag	UNP B5CY96
A	?	-	ASN	deletion	UNP B5CY96

- Molecule 2 is an oligosaccharide called 6-O-sulfo-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose-(1-4)-3,6-anhydro-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose-(1-4)-

6-O-sulfo-alpha-L-galactopyranose-(1-3)-alpha-D-galactopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	O	S			
2	B	6	74	36	36	2	0	1	0

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	A	1	1	1	0	0

- Molecule 4 is beta-D-galactopyranose (CCD ID: GAL) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	12	6	6	0	1


- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
5	A	1	1	1	0	0

- Molecule 6 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	2	Total K 2 2	0	0

- Molecule 7 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 4 2 2	0	0

- Molecule 8 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	13	Total I 13 13	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	594	Total O 594 594	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GH86A BETA-PORPHYRANASE



- Molecule 2: 6-O-sulfo-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose-(1-4)-3,6-anhydro-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose-(1-4)-6-O-sulfo-alpha-L-galactopyranose-(1-3)-alpha-D-galactopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.33Å 87.70Å 114.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	69.61 – 1.33 69.61 – 1.33	Depositor EDS
% Data completeness (in resolution range)	99.7 (69.61-1.33) 99.7 (69.61-1.33)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.07 (at 1.33Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.137 , 0.152 0.146 , 0.157	Depositor DCC
R_{free} test set	8189 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	11.9	Xtrriage
Anisotropy	0.151	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5264	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: L6S, GAL, K, CL, AAL, CA, GLA, EDO, IOD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.81	2/4728 (0.0%)	0.85	0/6413

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	382	ILE	CA-CB	6.94	1.64	1.54
1	A	203	TYR	N-CA	5.36	1.52	1.46

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4563	0	4354	35	1
2	B	74	0	50	2	0
3	A	1	0	0	0	0
4	A	12	0	6	0	0
5	A	1	0	0	0	0
6	A	2	0	0	0	0
7	A	4	0	6	5	0
8	A	13	0	0	4	0
9	A	594	0	0	15	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5264	0	4416	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:373:LYS:HE3	9:A:2093:HOH:O	1.23	1.28
1:A:170[B]:ARG:HD3	9:A:2196:HOH:O	1.45	1.13
1:A:279:GLU:OE1	2:B:1[A]:GLA:O1	2.03	0.76
8:A:1703:IOD:I	9:A:2014:HOH:O	2.73	0.75
1:A:507:SER:OG	9:A:2500:HOH:O	1.94	0.75
1:A:455:ILE:HD13	7:A:1618:EDO:C1	2.18	0.74
1:A:455:ILE:CD1	7:A:1618:EDO:H11	2.17	0.73
8:A:1719:IOD:I	9:A:2591:HOH:O	2.77	0.71
1:A:373:LYS:CG	9:A:2093:HOH:O	2.41	0.69
1:A:373:LYS:CE	9:A:2093:HOH:O	1.99	0.68
1:A:152[A]:GLU:CD	1:A:202:ALA:HB2	2.20	0.67
1:A:455:ILE:HD13	7:A:1618:EDO:H12	1.77	0.66
1:A:351:ASN:OD1	9:A:2367:HOH:O	2.12	0.66
1:A:293[B]:ASP:OD1	1:A:296:SER:OG	2.07	0.64
1:A:597:VAL:O	1:A:598:SER:CB	2.47	0.63
8:A:1701:IOD:I	9:A:2031:HOH:O	2.86	0.61
1:A:70:GLN:NE2	9:A:2093:HOH:O	2.36	0.58
1:A:30:VAL:HG22	1:A:382:ILE:HD12	1.85	0.57
1:A:152[A]:GLU:CG	1:A:202:ALA:HB2	2.36	0.56
1:A:195:LYS:NZ	9:A:2191:HOH:O	2.22	0.56
1:A:455:ILE:HD13	7:A:1618:EDO:H11	1.80	0.55
1:A:284:ASP:OD2	1:A:296:SER:HB2	2.08	0.53
1:A:455:ILE:CD1	7:A:1618:EDO:C1	2.82	0.51
1:A:522:VAL:HG11	1:A:551:ARG:O	2.12	0.49
1:A:70:GLN:CD	9:A:2093:HOH:O	2.55	0.48
1:A:240:ILE:HG23	1:A:437[A]:TYR:CE2	2.49	0.48
1:A:155:VAL:HG13	8:A:1700:IOD:I	2.85	0.47
1:A:373:LYS:HG3	9:A:2093:HOH:O	2.11	0.47
1:A:30:VAL:HG22	1:A:382:ILE:CD1	2.45	0.44
1:A:452:PRO:HG2	1:A:455:ILE:HD11	2.00	0.44
1:A:519:GLY:HA2	1:A:554:PHE:O	2.18	0.44
1:A:480:ARG:HD2	1:A:480:ARG:C	2.43	0.43
1:A:233:SER:HA	1:A:276:ALA:O	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:195:LYS:HE2	9:A:2121:HOH:O	2.20	0.42
1:A:170[B]:ARG:NH2	9:A:2217:HOH:O	2.42	0.42
1:A:433:LEU:C	1:A:433:LEU:HD23	2.45	0.41
1:A:152[A]:GLU:OE2	2:B:1[A]:GLA:C1	2.69	0.41
1:A:471:SER:C	1:A:473:ALA:N	2.79	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:170[A]:ARG:NH1	9:A:2418:HOH:O[4_565]	1.94	0.26

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	574/591 (97%)	561 (98%)	13 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	501/512 (98%)	496 (99%)	5 (1%)	73 43

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	93	LYS
1	A	149	PRO
1	A	325	VAL
1	A	488	TYR
1	A	546	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	86	ASN
1	A	89	HIS
1	A	271	HIS
1	A	552	ASN
1	A	584	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLA	B	1[A]	2	12,12,12	1.25	1 (8%)	17,17,17	1.78	5 (29%)
2	L6S	B	2	4,2	15,15,16	0.94	1 (6%)	21,22,24	1.25	2 (9%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GAL	B	3	2	11,11,12	0.52	0	15,15,17	1.15	1 (6%)
2	AAL	B	4	2	11,11,12	0.45	0	13,16,18	1.46	3 (23%)
2	GAL	B	5	2	11,11,12	0.66	0	15,15,17	0.53	0
2	L6S	B	6	2	15,15,16	1.50	2 (13%)	21,22,24	0.83	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLA	B	1[A]	2	-	2/2/22/22	0/1/1/1
2	L6S	B	2	4,2	-	2/6/23/26	0/1/1/1
2	GAL	B	3	2	-	0/2/19/22	0/1/1/1
2	AAL	B	4	2	-	-	0/3/2/2
2	GAL	B	5	2	-	0/2/19/22	0/1/1/1
2	L6S	B	6	2	-	0/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	6	L6S	O6-S1	-4.04	1.46	1.56
2	B	6	L6S	O5-C5	2.93	1.49	1.43
2	B	1[A]	GLA	O5-C5	2.51	1.50	1.44
2	B	2	L6S	C4-C3	2.19	1.58	1.52

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1[A]	GLA	O5-C5-C6	4.75	118.21	106.44
2	B	2	L6S	C1-O5-C5	3.22	116.50	112.19
2	B	4	AAL	O5-C5-C6	-2.78	109.26	113.33
2	B	1[A]	GLA	O1-C1-C2	2.62	116.57	108.98
2	B	1[A]	GLA	C1-O5-C5	-2.51	108.79	113.65
2	B	4	AAL	C2-C3-C4	-2.50	107.66	113.85
2	B	1[A]	GLA	O3-C3-C4	-2.49	104.50	110.38
2	B	1[A]	GLA	O5-C5-C4	-2.48	105.23	109.70
2	B	2	L6S	O5-C5-C6	-2.27	102.69	107.59
2	B	4	AAL	O5-C1-C2	2.25	116.15	110.79
2	B	6	L6S	O6-S1-O2S	2.23	113.73	106.92
2	B	3	GAL	C1-C2-C3	-2.01	106.72	109.64

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1[A]	GLA	O5-C5-C6-O6
2	B	1[A]	GLA	C4-C5-C6-O6
2	B	2	L6S	O5-C5-C6-O6
2	B	2	L6S	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1[A]	GLA	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 17 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	EDO	A	1618	-	3,3,3	0.28	0	2,2,2	0.77	0
4	GAL	A	1609[B]	2	12,12,12	1.49	2 (16%)	17,17,17	1.69	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	A	1618	-	-	1/1/1/1	-
4	GAL	A	1609[B]	2	-	2/2/22/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1609[B]	GAL	O1-C1	2.93	1.48	1.39
4	A	1609[B]	GAL	O5-C5	2.51	1.50	1.44

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1609[B]	GAL	O5-C5-C6	4.75	118.21	106.44
4	A	1609[B]	GAL	C1-O5-C5	-2.51	108.79	113.65
4	A	1609[B]	GAL	O3-C3-C4	-2.49	104.50	110.38
4	A	1609[B]	GAL	O5-C5-C4	-2.48	105.23	109.70

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1609[B]	GAL	O5-C5-C6-O6
4	A	1609[B]	GAL	C4-C5-C6-O6
7	A	1618	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	1618	EDO	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	565/591 (95%)	-0.14	18 (3%) 50 61	6, 13, 25, 38	17 (3%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	437[A]	TYR	6.4
1	A	473	ALA	5.7
1	A	598	SER	4.4
1	A	284	ASP	3.2
1	A	287	LYS	3.0
1	A	288	PRO	2.9
1	A	239	GLY	2.8
1	A	240	ILE	2.6
1	A	438[A]	ASN	2.6
1	A	285	ASN	2.6
1	A	455	ILE	2.5
1	A	597	VAL	2.5
1	A	357	ASN	2.2
1	A	353	GLN	2.1
1	A	96	GLN	2.1
1	A	471	SER	2.0
1	A	382	ILE	2.0
1	A	355	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	L6S	B	6	15/16	0.89	0.10	21,27,31,31	1
2	GAL	B	5	11/12	0.95	0.07	19,21,25,27	0
2	GLA	B	1[A]	12/12	0.95	0.07	11,17,23,24	12
2	AAL	B	4	10/11	0.96	0.07	15,18,20,20	0
2	GAL	B	3	11/12	0.97	0.05	12,14,16,21	0
2	L6S	B	2	15/16	0.97	0.06	9,13,16,17	1

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	EDO	A	1618	4/4	0.94	0.10	24,26,27,32	0
4	GAL	A	1609[B]	12/12	0.96	0.06	11,15,23,24	12
6	K	A	1616	1/1	0.98	0.05	23,23,23,23	1
5	CL	A	1615	1/1	0.98	0.07	18,18,18,18	0
8	IOD	A	1701	1/1	0.99	0.03	17,17,17,17	1
8	IOD	A	1702	1/1	0.99	0.11	23,23,23,23	1
8	IOD	A	1703	1/1	0.99	0.05	31,31,31,31	1
8	IOD	A	1704	1/1	0.99	0.04	21,21,21,21	1
8	IOD	A	1719	1/1	0.99	0.11	28,28,28,28	1
8	IOD	A	1722	1/1	0.99	0.05	24,24,24,24	1
6	K	A	1617	1/1	1.00	0.10	16,16,16,16	1
8	IOD	A	1705	1/1	1.00	0.03	7,7,7,7	1
8	IOD	A	1706	1/1	1.00	0.02	17,17,17,17	1
8	IOD	A	1707	1/1	1.00	0.02	18,18,18,18	1
8	IOD	A	1708	1/1	1.00	0.04	18,18,18,18	1
3	CA	A	1599	1/1	1.00	0.01	10,10,10,10	0
8	IOD	A	1720	1/1	1.00	0.05	35,35,35,35	1
8	IOD	A	1721	1/1	1.00	0.04	19,19,19,19	1
8	IOD	A	1700	1/1	1.00	0.01	10,10,10,10	1

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.