



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2025 – 03:54 pm BST

PDB ID : 4BSI / pdb_00004bsi
Title : H7N3 Avian Influenza Virus Haemagglutinin in Complex with Avian Receptor Analogue 3'-SLN
Authors : Xiong, X.; Haire, L.F.; Martin, S.R.; Wharton, S.A.; Daniels, R.S.; Bennett, M.S.; McCauley, J.W.; Collins, P.J.; Walker, P.A.; Skehel, J.J.; Gamblin, S.J.
Deposited on : 2013-06-10
Resolution : 2.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.44


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	4623 (2.64-2.60)
Clashscore	180529	5071 (2.64-2.60)
Ramachandran outliers	177936	5006 (2.64-2.60)
Sidechain outliers	177891	5006 (2.64-2.60)
RSRZ outliers	164620	4622 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	
2	B	177	
3	C	3	
4	D	2	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 4009 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called HEMAGGLUTININ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	317	2418	1497	436	469	16	0	0	1

- Molecule 2 is a protein called HEMAGGLUTININ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	170	1380	850	243	279	8	0	0	0

- Molecule 3 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	3	46	25	2	19	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	D	2	28	16	2	10	0	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
5	A	1	14	8	1	5	0	0

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total O S 5 4 1	0	0
6	A	1	Total O S 5 4 1	0	0
6	A	1	Total O S 5 4 1	0	0
6	A	1	Total O S 5 4 1	0	0
6	A	1	Total O S 5 4 1	0	0
6	A	1	Total O S 5 4 1	0	0
6	A	1	Total O S 5 4 1	0	0
6	A	1	Total O S 5 4 1	0	0
6	A	1	Total O S 5 4 1	0	0
6	A	1	Total O S 5 4 1	0	0
6	B	1	Total O S 5 4 1	0	0
6	B	1	Total O S 5 4 1	0	0
6	B	1	Total O S 5 4 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	30	Total O 30 30	0	0
7	B	19	Total O 19 19	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HEMAGGLUTININ



- Molecule 2: HEMAGGLUTININ



- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics i

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	116.36Å 116.36Å 297.21Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	59.88 – 2.62 59.88 – 2.62	Depositor EDS
% Data completeness (in resolution range)	99.5 (59.88-2.62) 99.5 (59.88-2.62)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.09 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.215 , 0.269 0.219 , 0.271	Depositor DCC
R_{free} test set	1216 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	70.2	Xtrriage
Anisotropy	0.004	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.002 for $-1/3^*h+1/3^*k+1/3^*l,-k,8/3^*h+4/3^*k+1/3^*l$ 0.019 for $-2/3^*h-1/3^*k-1/3^*l,-1/3^*h-2/3^*k+1/3^*l,-4/3^*h+4/3^*k+1/3^*l$ 0.016 for $-h,1/3^*h-1/3^*k-1/3^*l,-4/3^*h-8/3^*k+1/3^*l$	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4009	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, SO4, NAG, SIA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	1/2464 (0.0%)	0.74	1/3330 (0.0%)
2	B	0.50	0/1404	0.74	0/1891
All	All	0.52	1/3868 (0.0%)	0.74	1/5221 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	316	GLU	C-N	-7.04	1.23	1.33

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2	LYS	N-CA-C	6.03	118.02	108.79

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2418	0	2363	27	0
2	B	1380	0	1281	9	0
3	C	46	0	40	2	0
4	D	28	0	25	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	28	0	26	0	0
6	A	45	0	0	0	0
6	B	15	0	0	0	0
7	A	30	0	0	0	0
7	B	19	0	0	1	0
All	All	4009	0	3735	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:256:MET:HE1	1:A:294:GLY:HA2	1.62	0.81
1:A:38:VAL:HG21	1:A:280:ILE:HD12	1.70	0.71
1:A:131:ARG:NH2	1:A:136:PHE:O	2.23	0.71
2:B:129:ASN:HD22	2:B:129:ASN:N	1.94	0.66
7:B:2008:HOH:O	4:D:1:NAG:H83	1.98	0.64
1:A:170:ILE:HD13	1:A:225:TRP:HB3	1.80	0.63
2:B:75:LYS:HE3	4:D:1:NAG:H81	1.82	0.61
1:A:163:ARG:HD3	1:A:250:PHE:CE2	2.37	0.60
1:A:226:LEU:HD12	1:A:226:LEU:C	2.28	0.58
1:A:6:GLY:C	2:B:115:MET:HE1	2.30	0.57
1:A:228:LEU:HD22	1:A:234:VAL:HG23	1.87	0.56
1:A:152:PHE:HB3	1:A:239:ASN:O	2.07	0.55
1:A:144:LEU:HD11	3:C:3:SIA:H111	1.89	0.54
2:B:56:ILE:HG22	2:B:56:ILE:O	2.08	0.53
1:A:166:PRO:HB2	1:A:227:MET:HE3	1.92	0.52
1:A:106:GLY:HA2	1:A:255:SER:HB3	1.92	0.51
1:A:44:LYS:HE2	1:A:269:GLU:HB2	1.92	0.51
1:A:211:ARG:O	1:A:218:SER:HB2	2.12	0.50
1:A:163:ARG:HD3	1:A:250:PHE:CZ	2.48	0.49
1:A:179:THR:O	1:A:183:THR:HG23	2.14	0.48
1:A:268:CYS:SG	1:A:280:ILE:HD11	2.54	0.47
1:A:98:LEU:HD11	1:A:168:LEU:HD21	1.96	0.47
3:C:2:GAL:H4	3:C:3:SIA:O6	2.14	0.46
1:A:170:ILE:HD13	1:A:225:TRP:CB	2.47	0.45
1:A:120:ILE:HD13	1:A:155:MET:HE1	1.99	0.45
1:A:170:ILE:CD1	1:A:225:TRP:HB3	2.48	0.44
1:A:60:ILE:CG2	1:A:170:ILE:HG13	2.48	0.44
1:A:197:SER:OG	1:A:198:SER:N	2.50	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:9:PHE:CD1	2:B:10:ILE:HD12	2.53	0.43
2:B:129:ASN:N	2:B:129:ASN:ND2	2.64	0.43
1:A:228:LEU:HD22	1:A:234:VAL:CG2	2.49	0.42
2:B:131:GLU:HG3	2:B:141:PHE:CZ	2.55	0.42
1:A:242:PHE:CE2	1:A:244:ALA:HB2	2.55	0.41
2:B:169:ASN:O	2:B:170:ARG:C	2.62	0.41
1:A:186:TYR:CZ	1:A:241:ALA:HA	2.56	0.41
1:A:2:LYS:HE2	2:B:137:CYS:SG	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	315/321 (98%)	305 (97%)	10 (3%)	0	100	100
2	B	168/177 (95%)	161 (96%)	7 (4%)	0	100	100
All	All	483/498 (97%)	466 (96%)	17 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/271 (98%)	257 (96%)	10 (4%)	29	53

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	145/152 (95%)	132 (91%)	13 (9%)	8	15
All	All	412/423 (97%)	389 (94%)	23 (6%)	17	36

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	CYS
1	A	14	THR
1	A	91	LYS
1	A	94	ASN
1	A	104	GLU
1	A	123	ASN
1	A	197	SER
1	A	237	SER
1	A	256	MET
1	A	261	SER
2	B	19	ASP
2	B	24	PHE
2	B	32	GLU
2	B	49	THR
2	B	66	ILE
2	B	90	GLU
2	B	127	ARG
2	B	129	ASN
2	B	137	CYS
2	B	143	LYS
2	B	164	GLU
2	B	168	GLN
2	B	170	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	146	ASN
1	A	149	ASN
1	A	199	ASN
1	A	287	ASN
2	B	129	ASN
2	B	155	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	3	15,15,15	0.52	0	21,21,21	1.32	1 (4%)
3	GAL	C	2	3	11,11,12	0.48	0	15,15,17	1.05	2 (13%)
3	SIA	C	3	3	20,20,21	0.62	0	24,28,31	1.38	3 (12%)
4	NAG	D	1	4,2	14,14,15	0.48	0	17,19,21	0.89	1 (5%)
4	NAG	D	2	4	14,14,15	0.53	0	17,19,21	0.91	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3	-	1/6/26/26	0/1/1/1
3	GAL	C	2	3	-	0/2/19/22	0/1/1/1
3	SIA	C	3	3	-	1/18/34/38	0/1/1/1
4	NAG	D	1	4,2	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	NAG	C1-C2-N2	-4.86	105.09	110.73
3	C	3	SIA	C4-C5-N5	-2.85	104.73	110.38
3	C	3	SIA	C6-O6-C2	2.80	117.34	111.34
3	C	3	SIA	O1B-C1-C2	2.59	120.43	113.03
4	D	2	NAG	O5-C5-C6	2.23	110.69	107.20
3	C	2	GAL	O5-C5-C6	2.15	110.57	107.20
3	C	2	GAL	C1-C2-C3	2.09	112.23	109.67
4	D	1	NAG	C1-O5-C5	2.04	114.95	112.19

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	2	NAG	O5-C5-C6-O6
4	D	2	NAG	C4-C5-C6-O6
3	C	1	NAG	O5-C5-C6-O6
3	C	3	SIA	C6-C5-N5-C10

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	2	GAL	1	0
4	D	1	NAG	2	0
3	C	3	SIA	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry i

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	SO4	A	1327	-	4,4,4	0.39	0	6,6,6	0.19	0
6	SO4	B	1174	-	4,4,4	0.36	0	6,6,6	0.19	0
5	NAG	A	1318	1	14,14,15	0.47	0	17,19,21	1.29	2 (11%)
6	SO4	A	1325	-	4,4,4	0.35	0	6,6,6	0.16	0
5	NAG	A	1319	1	14,14,15	0.43	0	17,19,21	1.70	3 (17%)
6	SO4	A	1323	-	4,4,4	0.36	0	6,6,6	0.05	0
6	SO4	A	1326	-	4,4,4	0.38	0	6,6,6	0.13	0
6	SO4	A	1321	-	4,4,4	0.34	0	6,6,6	0.13	0
6	SO4	A	1322	-	4,4,4	0.35	0	6,6,6	0.07	0
6	SO4	B	1173	-	4,4,4	0.35	0	6,6,6	0.14	0
6	SO4	B	1175	-	4,4,4	0.35	0	6,6,6	0.16	0
6	SO4	A	1324	-	4,4,4	0.38	0	6,6,6	0.29	0
6	SO4	A	1328	-	4,4,4	0.32	0	6,6,6	0.13	0
6	SO4	A	1320	-	4,4,4	0.34	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	1318	1	-	2/6/23/26	0/1/1/1
5	NAG	A	1319	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1319	NAG	C1-O5-C5	4.30	118.02	112.19
5	A	1319	NAG	O5-C1-C2	-3.98	105.01	111.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1318	NAG	C1-O5-C5	3.07	116.36	112.19
5	A	1318	NAG	C4-C3-C2	-2.64	107.15	111.02
5	A	1319	NAG	C1-C2-N2	2.02	113.94	110.49

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1318	NAG	C4-C5-C6-O6
5	A	1318	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	317/321 (98%)	0.16	10 (3%) 50 45	47, 72, 118, 134	0
2	B	170/177 (96%)	0.31	8 (4%) 37 33	45, 98, 139, 155	0
All	All	487/498 (97%)	0.21	18 (3%) 45 40	45, 81, 132, 155	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	201	GLN	3.9
2	B	58	LYS	3.7
2	B	59	THR	3.1
1	A	268	CYS	2.8
1	A	192	LEU	2.8
2	B	54	ARG	2.6
2	B	141	PHE	2.5
1	A	150	ALA	2.4
1	A	1	ASP	2.3
2	B	144	CYS	2.3
1	A	149	ASN	2.2
2	B	29	ALA	2.2
1	A	194	THR	2.2
2	B	130	ALA	2.1
1	A	119	GLY	2.1
2	B	153	ARG	2.1
1	A	143	LEU	2.0
1	A	151	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	C	1	15/15	0.57	0.13	124,130,135,135	0
4	NAG	D	2	14/15	0.73	0.14	91,104,109,113	0
3	GAL	C	2	11/12	0.78	0.12	98,117,123,123	0
4	NAG	D	1	14/15	0.88	0.10	63,79,85,93	0
3	SIA	C	3	20/21	0.89	0.10	71,82,98,101	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q < 0.9
6	SO4	A	1320	5/5	0.39	0.10	151,151,156,157	0
6	SO4	A	1328	5/5	0.48	0.12	141,144,144,144	0
6	SO4	A	1325	5/5	0.64	0.09	136,136,139,141	0
6	SO4	B	1175	5/5	0.64	0.17	122,130,136,139	0
5	NAG	A	1319	14/15	0.68	0.12	112,120,127,128	0
5	NAG	A	1318	14/15	0.69	0.14	83,96,100,102	0
6	SO4	A	1323	5/5	0.72	0.11	132,136,140,142	0
6	SO4	A	1321	5/5	0.74	0.11	127,128,129,130	0
6	SO4	B	1174	5/5	0.79	0.10	110,114,115,118	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	SO4	A	1322	5/5	0.79	0.09	105,108,111,112	0
6	SO4	A	1326	5/5	0.80	0.10	104,108,112,114	0
6	SO4	A	1324	5/5	0.85	0.09	97,103,109,113	0
6	SO4	A	1327	5/5	0.88	0.18	86,91,95,102	0
6	SO4	B	1173	5/5	0.88	0.07	102,106,110,112	0

6.5 Other polymers [i](#)

There are no such residues in this entry.