



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 16, 2025 – 10:02 PM EDT

PDB ID : 4JJ2 / pdb_00004jj2
Title : High resolution structure of a C-terminal fragment of the T4 phage gp5 beta-helix
Authors : Buth, S.A.; Leiman, P.G.; Boudko, S.P.
Deposited on : 2013-03-07
Resolution : 1.28 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0rc1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.44


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2484 (1.30-1.26)
Clashscore	180529	2694 (1.30-1.26)
Ramachandran outliers	177936	2628 (1.30-1.26)
Sidechain outliers	177891	2627 (1.30-1.26)
RSRZ outliers	164620	2481 (1.30-1.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	95	
1	B	95	
1	C	95	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2647 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tail-associated lysozyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	92	Total 778	C 481	N 124	O 169	S 4	0	15	0
1	B	92	Total 724	C 443	N 119	O 159	S 3	0	9	0
1	C	92	Total 757	C 462	N 123	O 169	S 3	0	12	0


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	481	GLY	-	expression tag	UNP P16009
A	482	SER	-	expression tag	UNP P16009
B	481	GLY	-	expression tag	UNP P16009
B	482	SER	-	expression tag	UNP P16009
C	481	GLY	-	expression tag	UNP P16009
C	482	SER	-	expression tag	UNP P16009

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Mg 1	0	0

- Molecule 3 is Elaidic acid (CCD ID: ELA) (formula: C₁₈H₃₄O₂).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
3	A	1	Total	C	O	0	0
			20	18	2		

- Molecule 4 is PALMITIC ACID (CCD ID: PLM) (formula: $C_{16}H_{32}O_2$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
4	B	1	Total	C	O	0	0
			18	16	2		

- Molecule 5 is STEARIC ACID (CCD ID: STE) (formula: $C_{18}H_{36}O_2$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	1	Total	C O	0	0
			20	18 2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	111	Total	O	0	0
			111	111		
6	B	110	Total	O	0	0
			110	110		
6	C	108	Total	O	0	0
			108	108		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tail-associated lysozyme



- Molecule 1: Tail-associated lysozyme



- Molecule 1: Tail-associated lysozyme



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	57.61Å 72.76Å 130.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.00 – 1.28 33.00 – 1.28	Depositor EDS
% Data completeness (in resolution range)	98.6 (33.00-1.28) 94.3 (33.00-1.28)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.28 (at 1.28Å)	Xtrriage
Refinement program	SHELX 2013	Depositor
R, R_{free}	0.153 , 0.178 0.160 , 0.186	Depositor DCC
R_{free} test set	2437 reflections (3.48%)	wwPDB-VP
Wilson B-factor (Å ²)	15.2	Xtrriage
Anisotropy	0.139	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 60.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2647	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PLM, ELA, STE, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.95	0/805	1.59	17/1090 (1.6%)
1	B	1.10	0/742	1.74	19/1003 (1.9%)
1	C	1.03	0/773	1.55	11/1045 (1.1%)
All	All	1.03	0/2320	1.63	47/3138 (1.5%)

There are no bond length outliers.

All (47) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	571	ARG	CD-NE-CZ	13.59	143.42	124.40
1	A	574	ILE	O-C-N	-11.62	110.71	123.26
1	B	571	ARG	CA-CB-CG	9.37	132.84	114.10
1	B	564	GLN	OE1-CD-NE2	9.13	131.73	122.60
1	A	564	GLN	OE1-CD-NE2	-8.80	113.80	122.60
1	B	564	GLN	CA-CB-CG	8.32	130.75	114.10
1	B	574	ILE	O-C-N	-7.79	114.85	123.26
1	B	573	ASP	CA-CB-CG	7.73	120.33	112.60
1	A	528	ASN	OD1-CG-ND2	-7.71	114.89	122.60
1	B	572	ILE	CB-CG1-CD1	7.66	129.88	113.80
1	B	564	GLN	N-CA-CB	-7.36	99.16	109.97
1	A	573	ASP	CA-CB-CG	-7.34	105.26	112.60
1	A	565[A]	TYR	N-CA-CB	-7.19	99.43	110.65
1	A	565[B]	TYR	N-CA-CB	-7.19	99.43	110.65
1	A	526	GLN	OE1-CD-NE2	-7.13	115.47	122.60
1	A	574	ILE	CA-C-O	6.99	127.73	120.39
1	B	567	ILE	CA-C-N	-6.93	111.27	122.36
1	B	567	ILE	C-N-CA	-6.93	111.27	122.36
1	B	573	ASP	CA-C-O	-6.84	113.14	120.46
1	C	528	ASN	OD1-CG-ND2	-6.37	116.23	122.60
1	B	565	TYR	N-CA-CB	-6.33	99.97	110.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	562	SER	CA-C-O	6.20	126.99	119.31
1	B	533	ASN	OD1-CG-ND2	-5.97	116.62	122.60
1	A	533	ASN	OD1-CG-ND2	-5.87	116.73	122.60
1	B	564	GLN	CG-CD-NE2	-5.84	107.64	116.40
1	C	531	ASN	CA-CB-CG	-5.76	106.84	112.60
1	C	559	SER	CA-CB-OG	-5.69	99.72	111.10
1	B	485	ASP	CA-C-O	5.68	127.53	121.16
1	B	560	ILE	CB-CA-C	-5.66	102.43	110.12
1	C	562	SER	O-C-N	-5.63	114.69	122.46
1	C	568[A]	ASP	CA-CB-CG	-5.54	107.06	112.60
1	C	568[B]	ASP	CA-CB-CG	-5.54	107.06	112.60
1	C	526	GLN	OE1-CD-NE2	-5.45	117.15	122.60
1	C	519	THR	CA-CB-OG1	-5.45	101.42	109.60
1	B	528	ASN	OD1-CG-ND2	-5.44	117.16	122.60
1	A	571	ARG	CG-CD-NE	5.36	123.79	112.00
1	B	545	ASP	CA-CB-CG	5.35	117.95	112.60
1	B	565	TYR	CB-CA-C	5.23	119.31	110.79
1	A	509	ASN	OD1-CG-ND2	-5.22	117.38	122.60
1	A	570	SER	O-C-N	5.13	128.00	122.15
1	A	574	ILE	CA-C-N	5.13	130.94	121.70
1	A	574	ILE	C-N-CA	5.13	130.94	121.70
1	A	564	GLN	N-CA-CB	-5.07	102.51	109.97
1	B	493	ASN	OD1-CG-ND2	-5.04	117.56	122.60
1	A	568[A]	ASP	CA-C-O	5.02	126.82	121.45
1	A	568[B]	ASP	CA-C-O	5.02	126.82	121.45
1	C	564	GLN	CG-CD-NE2	5.02	123.93	116.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	778	0	760	32	0
1	B	724	0	711	17	0
1	C	757	0	722	19	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	1	0	0	0	0
3	A	20	0	33	5	0
4	B	18	0	31	2	0
5	C	20	0	35	2	0
6	A	111	0	0	5	0
6	B	110	0	0	10	0
6	C	108	0	0	4	0
All	All	2647	0	2292	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:565[B]:TYR:CE2	1:B:568:ASP:O	1.69	1.45
1:A:565[B]:TYR:HE2	1:B:568:ASP:O	0.85	1.18
1:B:573:ASP:HA	6:B:810:HOH:O	1.69	0.93
1:A:512:ILE:HD11	3:A:602:ELA:H42	1.61	0.81
1:C:554[A]:MET:SD	1:C:557:MET:HE2	2.21	0.81
6:B:786:HOH:O	1:C:553[A]:LYS:HD3	1.85	0.76
1:C:512:ILE:HD11	5:C:601:STE:H41	1.67	0.76
1:A:553[B]:LYS:HD3	6:A:717:HOH:O	1.86	0.74
1:B:554[B]:MET:HG3	1:B:556:SER:O	1.89	0.72
1:B:554[B]:MET:SD	1:B:557:MET:HE2	2.28	0.72
1:A:568[B]:ASP:O	1:C:560:ILE:HG23	1.91	0.70
1:A:486:GLU:OE1	3:A:602:ELA:H182	1.92	0.69
1:A:565[B]:TYR:CE2	1:B:568:ASP:C	2.68	0.69
1:A:560[A]:ILE:HA	6:B:784:HOH:O	1.94	0.68
3:A:602:ELA:H142	1:B:496[B]:ILE:HG13	1.76	0.67
1:A:554[A]:MET:SD	1:A:557[A]:MET:HE2	2.35	0.66
1:A:560[B]:ILE:HD11	6:A:795:HOH:O	1.95	0.66
1:A:561[B]:SER:HB3	1:A:565[B]:TYR:CE2	2.31	0.66
1:A:537[B]:LYS:HG3	6:C:794:HOH:O	1.98	0.63
6:B:786:HOH:O	1:C:553[B]:LYS:HD2	1.99	0.62
1:A:560[B]:ILE:HG23	6:B:784:HOH:O	2.00	0.61
1:B:568:ASP:HB2	6:B:785:HOH:O	2.00	0.60
1:B:572:ILE:HG12	1:C:574[B]:ILE:CG2	2.35	0.57
1:A:523[A]:GLU:OE1	1:C:515:LYS:HD2	2.04	0.57
1:A:496:ILE:HD13	4:B:601:PLM:HA1	1.87	0.56
1:B:556:SER:HA	1:C:561[B]:SER:OG	2.08	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:512:ILE:HD11	4:B:601:PLM:H42	1.89	0.54
1:A:512:ILE:HD11	3:A:602:ELA:C4	2.33	0.54
1:A:560[B]:ILE:HA	6:B:784:HOH:O	2.10	0.51
1:A:488:LYS:HE3	1:A:489:THR:O	2.12	0.49
1:A:559[B]:SER:C	1:A:565[B]:TYR:HH	2.21	0.49
1:A:519:THR:OG1	1:C:511:ASP:OD1	2.28	0.48
1:A:553[B]:LYS:HD2	6:A:737:HOH:O	2.15	0.47
1:B:568:ASP:HB3	6:B:784:HOH:O	2.15	0.47
3:A:602:ELA:H112	1:C:502:VAL:HG11	1.96	0.46
1:A:488:LYS:HE2	1:C:486:GLU:OE2	2.15	0.46
1:A:553[A]:LYS:HE2	1:C:545:ASP:OD1	2.15	0.46
1:A:484:GLY:N	6:A:808:HOH:O	2.49	0.45
1:B:572:ILE:HG22	1:B:574:ILE:HG13	1.99	0.45
1:C:484:GLY:N	6:C:803:HOH:O	2.49	0.44
1:A:559[B]:SER:O	1:A:560[B]:ILE:HD13	2.17	0.44
1:B:484:GLY:N	6:B:809:HOH:O	2.50	0.44
1:C:572:ILE:CG2	1:C:574[B]:ILE:HG12	2.48	0.44
1:C:554[A]:MET:HG3	1:C:556:SER:O	2.18	0.44
1:C:574[A]:ILE:HA	6:C:802:HOH:O	2.18	0.43
1:A:484:GLY:N	6:A:809:HOH:O	2.50	0.43
1:A:498:VAL:HG21	1:A:502:VAL:CG1	2.49	0.43
1:A:488:LYS:HE3	1:A:489:THR:C	2.43	0.43
1:A:488:LYS:HG3	1:A:489:THR:N	2.30	0.43
1:A:523[B]:GLU:OE1	1:B:531:ASN:ND2	2.46	0.42
1:C:575[A]:GLY:N	6:C:802:HOH:O	2.52	0.42
1:C:512:ILE:HD11	5:C:601:STE:C4	2.44	0.41
1:A:498:VAL:HG11	1:A:502:VAL:CG1	2.50	0.41
1:A:561[B]:SER:HB3	1:A:565[B]:TYR:CZ	2.55	0.41
1:B:553[B]:LYS:HE2	1:C:560:ILE:O	2.21	0.41
1:B:573:ASP:CG	6:B:810:HOH:O	2.63	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	105/95 (110%)	105 (100%)	0	0	100	100
1	B	99/95 (104%)	99 (100%)	0	0	100	100
1	C	101/95 (106%)	101 (100%)	0	0	100	100
All	All	305/285 (107%)	305 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	89/76 (117%)	85 (96%)	4 (4%)	23	1
1	B	83/76 (109%)	82 (99%)	1 (1%)	67	35
1	C	85/76 (112%)	81 (95%)	4 (5%)	22	1
All	All	257/228 (113%)	248 (96%)	9 (4%)	34	4

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	488	LYS
1	A	559[A]	SER
1	A	559[B]	SER
1	A	571	ARG
1	B	572	ILE
1	C	515	LYS
1	C	553[A]	LYS
1	C	553[B]	LYS
1	C	572	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	526	GLN
1	A	528	ASN
1	A	533	ASN
1	A	564	GLN
1	B	526	GLN
1	B	533	ASN
1	C	526	GLN
1	C	528	ASN
1	C	533	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	STE	C	601	-	19,19,19	0.62	0	19,19,19	1.33	3 (15%)
4	PLM	B	601	-	17,17,17	0.56	0	17,17,17	0.81	0
3	ELA	A	602	-	19,19,19	1.08	1 (5%)	19,19,19	1.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	STE	C	601	-	-	11/17/17/17	-
4	PLM	B	601	-	-	8/15/15/15	-
3	ELA	A	602	-	-	11/17/17/17	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	ELA	C10-C9	3.79	1.53	1.31

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	601	STE	C3-C2-C1	-2.79	107.23	114.51
5	C	601	STE	O1-C1-C2	-2.62	114.78	123.09
5	C	601	STE	O2-C1-O1	2.11	128.75	123.33

There are no chirality outliers.

All (30) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	601	STE	C1-C2-C3-C4
5	C	601	STE	C11-C10-C9-C8
5	C	601	STE	C12-C13-C14-C15
4	B	601	PLM	C8-C9-CA-CB
4	B	601	PLM	C1-C2-C3-C4
5	C	601	STE	C14-C15-C16-C17
3	A	602	ELA	C3-C4-C5-C6
3	A	602	ELA	C12-C13-C14-C15
3	A	602	ELA	C6-C7-C8-C9
3	A	602	ELA	C10-C11-C12-C13
3	A	602	ELA	C11-C10-C9-C8
3	A	602	ELA	C4-C5-C6-C7
4	B	601	PLM	CB-CC-CD-CE
5	C	601	STE	C2-C3-C4-C5
3	A	602	ELA	C15-C16-C17-C18
5	C	601	STE	C3-C4-C5-C6
4	B	601	PLM	C2-C3-C4-C5
5	C	601	STE	C10-C11-C12-C13

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	602	ELA	C5-C6-C7-C8
3	A	602	ELA	C14-C15-C16-C17
5	C	601	STE	C15-C16-C17-C18
4	B	601	PLM	C4-C5-C6-C7
4	B	601	PLM	O1-C1-C2-C3
5	C	601	STE	C4-C5-C6-C7
3	A	602	ELA	O-C-CA-C3
4	B	601	PLM	O2-C1-C2-C3
3	A	602	ELA	OXT-C-CA-C3
5	C	601	STE	C6-C7-C8-C9
5	C	601	STE	C9-C10-C11-C12
4	B	601	PLM	CC-CD-CE-CF


There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	601	STE	2	0
4	B	601	PLM	2	0
3	A	602	ELA	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	92/95 (96%)	0.28	3 (3%) 49 50	8, 20, 43, 58	15 (16%)
1	B	92/95 (96%)	0.34	5 (5%) 32 33	8, 21, 46, 69	9 (9%)
1	C	92/95 (96%)	0.29	3 (3%) 49 50	7, 20, 32, 38	12 (13%)
All	All	276/285 (96%)	0.30	11 (3%) 43 44	7, 20, 41, 69	36 (13%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	574	ILE	5.5
1	B	574	ILE	3.2
1	C	572	ILE	3.1
1	A	560[A]	ILE	3.0
1	B	568	ASP	2.5
1	B	484	GLY	2.4
1	C	484	GLY	2.4
1	C	560	ILE	2.4
1	A	484	GLY	2.3
1	B	487	THR	2.2
1	B	571	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ELA	A	602	20/20	0.74	0.20	45,59,80,86	0
5	STE	C	601	20/20	0.80	0.17	46,56,70,80	0
4	PLM	B	601	18/18	0.82	0.16	50,59,63,69	0
2	MG	A	601	1/1	0.99	0.03	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around STE C 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around PLM B 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.