



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 28, 2025 – 12:42 PM EDT

PDB ID : 4JTJ / pdb_00004jtj
Title : Crystal structure of R117K mutant of 3-deoxy-D-manno-octulosonate 8-phosphate synthase (KDO8PS) from *Neisseria meningitidis*
Authors : Allison, T.M.; Cochrane, F.C.; Jameson, G.B.; Parker, E.J.
Deposited on : 2013-03-23
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2888 (1.76-1.76)
Clashscore	180529	3097 (1.76-1.76)
Ramachandran outliers	177936	3072 (1.76-1.76)
Sidechain outliers	177891	3072 (1.76-1.76)
RSRZ outliers	164620	2887 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	<div> <div>8%</div> <div> <div></div> <div>81%</div> <div>9%</div> <div>10%</div> </div> </div>
1	B	280	<div> <div>4%</div> <div> <div></div> <div>77%</div> <div>12%</div> <div>10%</div> </div> </div>
1	C	280	<div> <div>7%</div> <div> <div></div> <div>83%</div> <div>9%</div> <div>8%</div> </div> </div>
1	D	280	<div> <div>9%</div> <div> <div></div> <div>83%</div> <div>9%</div> <div>8%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8274 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-dehydro-3-deoxyphosphooctonate aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	253	Total	C	N	O	S	0	2	0
			1963	1260	332	360	11			
1	B	253	Total	C	N	O	S	0	5	0
			1981	1274	333	362	12			
1	C	257	Total	C	N	O	S	0	3	0
			1991	1282	334	364	11			
1	D	259	Total	C	N	O	S	0	1	0
			1997	1280	338	368	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	117	LYS	ARG	engineered mutation	UNP Q9JZ55
B	117	LYS	ARG	engineered mutation	UNP Q9JZ55
C	117	LYS	ARG	engineered mutation	UNP Q9JZ55
D	117	LYS	ARG	engineered mutation	UNP Q9JZ55

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		
2	C	1	Total	Cl	0	0
			1	1		
2	D	1	Total	Cl	0	0
			1	1		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	Na	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	77	Total 77	O 77	0	0
4	B	96	Total 96	O 96	0	0
4	C	92	Total 92	O 92	0	0
4	D	73	Total 73	O 73	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase




- Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase




- Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase



- Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	81.47Å 85.37Å 163.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.55 – 1.75 39.55 – 1.75	Depositor EDS
% Data completeness (in resolution range)	98.5 (39.55-1.75) 98.7 (39.55-1.75)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 1.75Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.197 , 0.226 0.203 , 0.229	Depositor DCC
R_{free} test set	5686 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	31.5	Xtriage
Anisotropy	0.328	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 30.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.004 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8274	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.09	1/1998 (0.1%)	0.98	3/2698 (0.1%)
1	B	1.16	1/2025 (0.0%)	1.02	2/2735 (0.1%)
1	C	1.22	4/2034 (0.2%)	1.00	2/2749 (0.1%)
1	D	1.12	1/2034 (0.0%)	1.02	4/2748 (0.1%)
All	All	1.15	7/8091 (0.1%)	1.00	11/10930 (0.1%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	56	ASP	CA-C	-6.87	1.47	1.53
1	D	125	MET	SD-CE	-5.85	1.65	1.79
1	A	236	GLU	CA-C	5.54	1.59	1.52
1	C	173	ASN	CA-CB	-5.49	1.42	1.54
1	B	28	LEU	N-CA	-5.38	1.40	1.46
1	C	79	GLU	C-O	-5.34	1.17	1.24
1	C	30	SER	N-CA	-5.15	1.40	1.46

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	172	ASP	CB-CA-C	-6.46	101.52	111.66
1	B	201	LEU	N-CA-C	5.96	120.68	113.17
1	A	236	GLU	CB-CA-C	5.81	119.30	109.55
1	A	201	LEU	CB-CA-C	-5.56	100.77	110.17
1	B	201	LEU	CB-CA-C	-5.42	101.04	110.09
1	D	172	ASP	CB-CA-C	-5.19	103.30	111.51
1	A	172	ASP	CB-CA-C	-5.19	103.31	111.51
1	D	252	LEU	CA-C-N	-5.13	114.48	119.76
1	D	252	LEU	C-N-CA	-5.13	114.48	119.76
1	C	232	GLY	N-CA-C	5.06	118.08	110.18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	201	LEU	CB-CA-C	-5.06	101.62	110.17

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1963	0	2017	21	0
1	B	1981	0	2047	30	0
1	C	1991	0	2053	9	0
1	D	1997	0	2039	13	0
2	A	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	C	1	0	0	0	0
4	A	77	0	0	1	0
4	B	96	0	0	2	0
4	C	92	0	0	0	0
4	D	73	0	0	3	0
All	All	8274	0	8156	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:204:ARG:HH21	1:B:204:ARG:HG2	1.37	0.89
1:B:13:ASN:HD21	1:B:229:ARG:HH21	1.26	0.80
1:A:118:GLN:NE2	1:C:114:PHE:HE2	1.84	0.74
1:D:37:HIS:CD2	1:D:254:LEU:HG	2.25	0.71
1:A:118:GLN:NE2	1:C:114:PHE:CE2	2.61	0.69
1:B:214:ARG:HB3	1:B:217:GLN:HE21	1.60	0.67
1:B:204:ARG:HH21	1:B:204:ARG:CG	2.08	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:116:ALA:HB2	4:D:466:HOH:O	1.95	0.64
1:A:236:GLU:O	1:A:237:SER:CB	2.47	0.63
1:B:13:ASN:ND2	1:B:229:ARG:HH21	1.97	0.62
1:A:197[B]:VAL:CG1	1:A:233:LEU:HD11	2.29	0.61
1:A:197[B]:VAL:HG11	1:A:233:LEU:HD11	1.82	0.61
1:A:267:LEU:HD13	1:B:267:LEU:CD1	2.32	0.60
1:D:217:GLN:NE2	4:D:468:HOH:O	2.35	0.59
1:B:195:PHE:CE2	1:B:197[B]:VAL:HG22	2.38	0.59
1:A:267:LEU:CD1	1:B:267:LEU:HD13	2.32	0.59
1:D:45:LEU:O	1:D:265:LYS:HE3	2.03	0.59
1:B:279:ILE:HG22	1:B:279:ILE:O	2.02	0.58
1:C:201:LEU:HD11	1:C:218:ALA:HA	1.86	0.58
1:C:45:LEU:O	1:C:265:LYS:HE3	2.04	0.57
1:B:199:HIS:HB2	1:B:236:GLU:OE2	2.06	0.55
1:B:45:LEU:O	1:B:265:LYS:HE3	2.06	0.55
1:D:62:SER:HB3	1:D:65:SER:OG	2.05	0.55
1:C:39:VAL:O	1:C:43:ARG:HG3	2.08	0.54
1:A:62:SER:HB3	1:A:65:SER:OG	2.07	0.54
1:D:214:ARG:HH21	1:D:216:ALA:H	1.54	0.54
1:B:52:LYS:HE2	1:B:234:PHE:CZ	2.43	0.53
1:B:114:PHE:HE2	1:D:118:GLN:HG2	1.74	0.52
1:B:204:ARG:CG	1:B:204:ARG:NH2	2.73	0.52
1:B:180:GLY:HA3	4:B:324:HOH:O	2.11	0.50
1:A:254:LEU:HD23	1:A:255:HIS:NE2	2.27	0.50
1:A:267:LEU:HD13	1:B:267:LEU:HD13	1.95	0.48
1:A:236:GLU:O	1:A:237:SER:OG	2.28	0.48
1:A:235:LEU:HD13	1:A:237:SER:HA	1.96	0.47
1:D:267:LEU:C	1:D:267:LEU:HD23	2.39	0.47
1:A:254:LEU:HD23	1:A:255:HIS:CD2	2.48	0.47
1:C:100:GLN:O	1:C:104:GLU:HG3	2.14	0.47
1:A:267:LEU:CD1	1:B:267:LEU:CD1	2.92	0.46
1:D:180:GLY:HA3	4:D:440:HOH:O	2.15	0.46
1:B:93[B]:VAL:CG1	1:B:98:GLN:HB2	2.47	0.45
1:A:254:LEU:O	1:A:254:LEU:HG	2.17	0.45
1:B:63:ILE:HA	1:D:119:THR:HG21	1.99	0.45
1:A:20:GLY:HA2	1:A:235:LEU:O	2.18	0.44
1:A:45:LEU:O	1:A:265:LYS:HE3	2.16	0.44
1:B:141:SER:OG	1:B:144[A]:GLN:HG3	2.18	0.44
1:B:59[B]:ASN:O	1:B:59[B]:ASN:OD1	2.36	0.44
1:B:140:LEU:HG	1:B:144[A]:GLN:HB2	1.99	0.44
1:B:93[B]:VAL:HG13	1:B:98:GLN:HB2	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:22:ILE:HG23	1:B:34:THR:HG21	2.00	0.43
1:C:20:GLY:HA2	1:C:235:LEU:O	2.17	0.43
1:B:13:ASN:HD21	1:B:229:ARG:NH2	2.03	0.43
1:B:201:LEU:HD11	1:B:218:ALA:HA	2.00	0.43
1:C:141:SER:OG	1:C:144:GLN:HG3	2.18	0.43
1:B:110:GLN:O	1:B:125[B]:MET:HE1	2.18	0.43
1:A:180:GLY:HA3	4:A:427:HOH:O	2.18	0.43
1:D:37:HIS:HD2	1:D:254:LEU:HG	1.82	0.43
1:C:140:LEU:HG	1:C:144:GLN:HB2	2.01	0.41
1:B:63:ILE:HG22	1:D:119:THR:HG23	2.03	0.41
1:A:56:ASP:OD1	1:A:67:ARG:CZ	2.69	0.41
1:A:140:LEU:HG	1:A:144:GLN:HB2	2.03	0.40
1:A:197[B]:VAL:CG1	1:A:233:LEU:CD1	2.97	0.40
1:B:151:LYS:HD3	4:B:386:HOH:O	2.22	0.40
1:B:236:GLU:O	1:B:237:SER:C	2.65	0.40
1:D:57:LYS:HD2	1:D:57:LYS:HA	1.91	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/280 (89%)	244 (98%)	4 (2%)	1 (0%)	30	16
1	B	252/280 (90%)	245 (97%)	6 (2%)	1 (0%)	30	16
1	C	254/280 (91%)	249 (98%)	4 (2%)	1 (0%)	30	16
1	D	254/280 (91%)	250 (98%)	3 (1%)	1 (0%)	30	16
All	All	1009/1120 (90%)	988 (98%)	17 (2%)	4 (0%)	30	16

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	229	ARG
1	B	229	ARG
1	A	229	ARG
1	C	229	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/236 (92%)	213 (98%)	5 (2%)	45	25
1	B	222/236 (94%)	216 (97%)	6 (3%)	40	19
1	C	222/236 (94%)	218 (98%)	4 (2%)	54	37
1	D	220/236 (93%)	212 (96%)	8 (4%)	30	11
All	All	882/944 (93%)	859 (97%)	23 (3%)	42	21

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LYS
1	A	73	GLU
1	A	165	ARG
1	A	186	GLN
1	A	254	LEU
1	B	13	ASN
1	B	57	LYS
1	B	80	LYS
1	B	204	ARG
1	B	255	HIS
1	B	277	LEU
1	C	165	ARG
1	C	202	GLN
1	C	252	LEU
1	C	277	LEU
1	D	44	LYS
1	D	127	LYS
1	D	154[A]	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	154[B]	GLU
1	D	165	ARG
1	D	205	ASP
1	D	214	ARG
1	D	255	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (28) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	ASN
1	A	23	ASN
1	A	64	HIS
1	A	110	GLN
1	A	118	GLN
1	A	133	ASN
1	B	13	ASN
1	B	37	HIS
1	B	138	GLN
1	B	173	ASN
1	B	199	HIS
1	B	217	GLN
1	B	274	GLN
1	C	97	HIS
1	C	100	GLN
1	C	186	GLN
1	C	217	GLN
1	C	238	HIS
1	C	274	GLN
1	D	23	ASN
1	D	37	HIS
1	D	64	HIS
1	D	138	GLN
1	D	147	ASN
1	D	186	GLN
1	D	217	GLN
1	D	255	HIS
1	D	274	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	253/280 (90%)	0.47	22 (8%)	17 20	17, 35, 66, 80	3 (1%)
1	B	253/280 (90%)	0.17	10 (3%)	43 48	15, 29, 48, 75	5 (1%)
1	C	257/280 (91%)	0.24	19 (7%)	22 25	15, 29, 48, 85	3 (1%)
1	D	259/280 (92%)	0.59	26 (10%)	14 16	21, 36, 67, 75	1 (0%)
All	All	1022/1120 (91%)	0.37	77 (7%)	22 24	15, 31, 64, 85	12 (1%)

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	277	LEU	7.5
1	D	277	LEU	5.3
1	C	277	LEU	5.2
1	C	203	THR	5.2
1	A	235	LEU	5.1
1	A	254	LEU	5.1
1	B	279	ILE	4.9
1	C	278	THR	4.9
1	C	202	GLN	4.7
1	C	252	LEU	4.4
1	D	252	LEU	4.4
1	C	251	ALA	4.3
1	C	238	HIS	4.3
1	D	238	HIS	4.1
1	D	63	ILE	4.0
1	C	114	PHE	3.8
1	D	71	LEU	3.7
1	D	213	GLY	3.7
1	B	201	LEU	3.7
1	D	253	PRO	3.6
1	A	66	TYR	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	206	ALA	3.4
1	B	277	LEU	3.4
1	A	237	SER	3.4
1	C	254	LEU	3.3
1	C	201	LEU	3.3
1	D	254	LEU	3.2
1	C	255	HIS	3.0
1	B	203	THR	3.0
1	D	22	ILE	3.0
1	C	204	ARG	2.9
1	B	202	GLN	2.9
1	A	40	GLU	2.9
1	A	71	LEU	2.9
1	D	66	TYR	2.9
1	A	63	ILE	2.8
1	D	255	HIS	2.8
1	D	61	SER	2.8
1	B	114	PHE	2.8
1	C	214	ARG	2.8
1	C	253	PRO	2.7
1	A	59[A]	ASN	2.7
1	D	203	THR	2.7
1	A	61	SER	2.7
1	C	215	ARG	2.6
1	D	215	ARG	2.6
1	B	237	SER	2.6
1	A	202	GLN	2.6
1	C	63	ILE	2.5
1	D	251	ALA	2.5
1	D	202	GLN	2.5
1	A	203	THR	2.4
1	D	235	LEU	2.4
1	A	213	GLY	2.4
1	B	256	LEU	2.4
1	D	117	LYS	2.3
1	A	22	ILE	2.3
1	A	97	HIS	2.3
1	A	67	ARG	2.3
1	D	172	ASP	2.3
1	B	278	THR	2.2
1	B	276	ILE	2.2
1	D	94	HIS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	256	LEU	2.2
1	A	83	ALA	2.2
1	A	114	PHE	2.2
1	D	55	PHE	2.1
1	D	28	LEU	2.1
1	A	215	ARG	2.1
1	D	237	SER	2.1
1	C	172	ASP	2.1
1	A	64	HIS	2.1
1	C	94	HIS	2.1
1	A	140	LEU	2.1
1	D	29	ASP	2.0
1	D	59	ASN	2.0
1	C	235	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CL	D	301	1/1	0.93	0.12	40,40,40,40	0
2	CL	C	301	1/1	0.94	0.14	41,41,41,41	0
2	CL	A	301	1/1	0.97	0.08	34,34,34,34	0
3	NA	C	302	1/1	0.98	0.04	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.