



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2025 – 01:19 pm BST

PDB ID : 4ZB0 / pdb_00004zb0
Title : A dehydrated form of glucose isomerase collected at room temperature.
Authors : Sandy, J.
Deposited on : 2015-04-14
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.44


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	387	% 94% 6%
1	B	387	% 93% 6%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6618 atoms, of which 24 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Xylose isomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	387	Total	C	N	O	S	0	0	0
			3045	1913	550	574	8			
1	B	387	Total	C	N	O	S	0	0	0
			3045	1913	550	574	8			

- Molecule 2 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	5	Total	Mn	0	0
			5	5		
2	B	5	Total	Mn	0	0
			5	5		

- Molecule 3 is alpha-D-glucopyranose (CCD ID: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 12 6 6	0	0
3	B	1	Total C H O 24 6 12 6	0	0

- Molecule 4 is beta-D-fructofuranose (CCD ID: FRU) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C H O 24 6 12 6	0	0
4	B	1	Total C O 12 6 6	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	236	Total O 236 236	0	0
5	B	210	Total O 210 210	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Xylose isomerase



- Molecule 1: Xylose isomerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	83.48Å 95.02Å 98.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	68.38 – 2.00 68.38 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.1 (68.38-2.00) 99.1 (68.38-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.22 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.137 , 0.180 0.150 , 0.187	Depositor DCC
R_{free} test set	2622 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtriage
Anisotropy	0.746	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 43.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.016 for -h,l,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6618	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, FRU, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.31	5/3117 (0.2%)	1.13	1/4219 (0.0%)
1	B	1.28	5/3117 (0.2%)	1.11	8/4219 (0.2%)
All	All	1.29	10/6234 (0.2%)	1.12	9/8438 (0.1%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	175	ASP	CB-CG	7.27	1.70	1.52
1	B	175	ASP	CA-C	6.29	1.60	1.53
1	A	339	ALA	C-O	-6.16	1.16	1.24
1	A	216	PRO	C-O	5.40	1.29	1.23
1	B	174	TYR	C-O	-5.26	1.17	1.23
1	A	214	VAL	N-CA	5.19	1.52	1.46
1	A	368	ARG	C-O	-5.19	1.18	1.24
1	A	98	VAL	N-CA	-5.11	1.40	1.46
1	B	92	ASN	N-CA	-5.08	1.40	1.46
1	B	84	MET	C-O	-5.03	1.17	1.23

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	175	ASP	CB-CA-C	7.65	123.59	111.73
1	B	347	LEU	N-CA-C	5.93	117.42	111.07
1	B	24	ASP	CA-C-N	-5.72	113.78	119.56
1	B	24	ASP	C-N-CA	-5.72	113.78	119.56
1	B	266	ARG	NE-CZ-NH2	-5.52	114.23	119.20
1	B	109	ARG	N-CA-C	5.49	117.26	111.28
1	B	74	ARG	CB-CA-C	-5.29	102.01	110.79
1	A	266	ARG	NE-CZ-NH2	-5.20	114.52	119.20
1	B	175	ASP	CA-C-O	5.05	127.84	121.58

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3045	0	2914	11	0
1	B	3045	0	2914	12	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	12	0	10	0	0
3	B	12	12	9	2	0
4	A	12	12	12	1	0
4	B	12	0	12	0	0
5	A	236	0	0	7	0
5	B	210	0	0	8	2
All	All	6594	24	5871	25	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:175:ASP:CB	5:B:524:HOH:O	2.40	0.70
1:B:175:ASP:HB3	5:B:524:HOH:O	1.98	0.64
1:A:340:ARG:HD3	5:A:644:HOH:O	2.05	0.56
1:B:245:ASP:OD1	1:B:285:HIS:HD2	1.91	0.53
1:A:382:HIS:HE1	5:A:586:HOH:O	1.91	0.53
1:B:331:ARG:NH2	5:B:502:HOH:O	2.43	0.52
1:B:24:ASP:HB2	1:B:25:PRO:CD	2.41	0.51
4:A:407:FRU:H62	5:A:579:HOH:O	2.10	0.50
3:B:406:GLC:H61	5:B:611:HOH:O	2.12	0.49
3:B:406:GLC:C6	5:B:611:HOH:O	2.61	0.48
1:B:2:ASN:HD22	1:B:3:TYR:H	1.61	0.48
1:A:381:ASP:HB3	1:A:387:ARG:HG2	1.96	0.48
1:A:177:ARG:NH1	5:A:504:HOH:O	2.47	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:41:ARG:NE	5:B:503:HOH:O	2.45	0.47
1:A:245:ASP:OD1	1:A:285:HIS:HD2	1.97	0.47
1:A:382:HIS:CE1	5:A:586:HOH:O	2.66	0.47
1:B:76:ARG:NH2	5:B:504:HOH:O	2.48	0.46
1:B:24:ASP:HB2	1:B:25:PRO:HD3	1.98	0.45
1:B:175:ASP:HB2	5:B:524:HOH:O	2.14	0.44
1:A:218:VAL:HG22	1:A:228:PHE:CE2	2.53	0.44
1:A:97:PRO:HB2	1:B:366:ALA:HB1	2.01	0.42
1:A:340:ARG:CD	5:A:644:HOH:O	2.66	0.42
1:B:228:PHE:CZ	1:B:232:ILE:HD11	2.55	0.42
1:A:387:ARG:HB2	5:A:662:HOH:O	2.19	0.42
1:A:10:ARG:NH1	1:A:281:SER:O	2.50	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:512:HOH:O	5:B:583:HOH:O[2_565]	1.22	0.98
5:B:583:HOH:O	5:B:583:HOH:O[2_565]	1.65	0.55

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/387 (100%)	371 (96%)	13 (3%)	1 (0%)	37	35
1	B	385/387 (100%)	372 (97%)	11 (3%)	2 (0%)	25	21
All	All	770/774 (100%)	743 (96%)	24 (3%)	3 (0%)	30	27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	GLU
1	B	186	GLU
1	B	254	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/303 (100%)	299 (99%)	4 (1%)	65	71
1	B	303/303 (100%)	297 (98%)	6 (2%)	50	55
All	All	606/606 (100%)	596 (98%)	10 (2%)	56	61

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ASN
1	A	42	ARG
1	A	352	ASP
1	A	354	ARG
1	B	2	ASN
1	B	7	PRO
1	B	42	ARG
1	B	69	GLU
1	B	76	ARG
1	B	355	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	ASN
1	A	285	HIS
1	A	348	GLN
1	B	2	ASN
1	B	49	HIS
1	B	285	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	348	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 10 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	FRU	B	407	-	11,12,12	1.51	2 (18%)	10,18,18	4.06	7 (70%)
3	GLC	A	406	2	12,12,12	1.13	1 (8%)	17,17,17	2.30	8 (47%)
3	GLC	B	406	2	12,12,12	1.12	1 (8%)	17,17,17	1.82	2 (11%)
4	FRU	A	407	-	11,12,12	0.84	0	10,18,18	1.04	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FRU	B	407	-	-	0/5/24/24	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	A	406	2	-	1/2/22/22	0/1/1/1
3	GLC	B	406	2	-	1/2/22/22	0/1/1/1
4	FRU	A	407	-	-	2/5/24/24	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	407	FRU	C4-C5	3.13	1.61	1.53
4	B	407	FRU	C6-C5	-2.68	1.42	1.51
3	A	406	GLC	O5-C1	2.38	1.48	1.42
3	B	406	GLC	O5-C5	-2.37	1.38	1.44

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	407	FRU	O6-C6-C5	-7.98	83.91	111.29
3	B	406	GLC	O5-C5-C6	-5.74	92.17	106.44
4	B	407	FRU	O4-C4-C5	5.20	126.07	111.05
4	B	407	FRU	O5-C5-C4	4.81	117.47	105.49
3	A	406	GLC	C4-C3-C2	4.18	118.12	110.82
3	A	406	GLC	O3-C3-C2	4.09	119.80	110.35
3	A	406	GLC	O4-C4-C5	4.02	119.29	109.30
4	B	407	FRU	C5-C4-C3	-3.86	89.52	101.91
4	B	407	FRU	C6-C5-C4	-3.82	105.88	115.09
4	B	407	FRU	O5-C5-C6	-3.71	98.51	108.85
3	A	406	GLC	O2-C2-C1	2.85	115.76	109.16
3	A	406	GLC	O5-C5-C6	-2.82	99.42	106.44
3	A	406	GLC	O2-C2-C3	2.65	116.48	110.35
4	B	407	FRU	O2-C2-O5	-2.61	104.47	109.50
3	A	406	GLC	C1-O5-C5	2.51	118.40	113.66
3	B	406	GLC	O4-C4-C5	2.38	115.22	109.30
3	A	406	GLC	O5-C5-C4	2.31	113.89	109.69

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	407	FRU	O1-C1-C2-C3
4	A	407	FRU	C4-C5-C6-O6
3	B	406	GLC	O5-C5-C6-O6
3	A	406	GLC	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	406	GLC	2	0
4	A	407	FRU	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	387/387 (100%)	-0.61	3 (0%) 82 82	11, 17, 35, 86	0
1	B	387/387 (100%)	-0.61	3 (0%) 82 82	12, 17, 38, 76	0
All	All	774/774 (100%)	-0.61	6 (0%) 82 82	11, 17, 36, 86	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	388	GLY	3.9
1	B	175	ASP	3.6
1	B	388	GLY	3.1
1	B	2	ASN	2.8
1	A	387	ARG	2.3
1	A	175	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	FRU	A	407	12/12	0.65	0.12	20,20,20,20	0
3	GLC	B	406	12/12	0.87	0.10	20,20,20,20	0
2	MN	B	405	1/1	0.90	0.08	55,55,55,55	0
4	FRU	B	407	12/12	0.91	0.09	16,20,31,37	0
2	MN	B	401	1/1	0.94	0.05	17,17,17,17	0
3	GLC	A	406	12/12	0.95	0.07	17,19,24,42	0
2	MN	A	405	1/1	0.96	0.27	73,73,73,73	0
2	MN	A	404	1/1	0.97	0.05	33,33,33,33	0
2	MN	B	404	1/1	0.99	0.23	49,49,49,49	0
2	MN	B	402	1/1	0.99	0.02	16,16,16,16	0
2	MN	B	403	1/1	0.99	0.34	52,52,52,52	0
2	MN	A	403	1/1	1.00	0.01	15,15,15,15	0
2	MN	A	401	1/1	1.00	0.02	14,14,14,14	0
2	MN	A	402	1/1	1.00	0.02	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.