



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 13, 2025 – 01:14 PM EDT

PDB ID : 5BT9 / pdb_00005bt9
Title : Crystal Structure of FolM Alternative dihydrofolate reductase 1 from *Brucella canis* complexed with NADP
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2015-06-02
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3717 (1.50-1.50)
Clashscore	180529	4048 (1.50-1.50)
Ramachandran outliers	177936	3970 (1.50-1.50)
Sidechain outliers	177891	3967 (1.50-1.50)
RSRZ outliers	164620	3718 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	4% 86% 6% 7%
1	B	269	4% 87% 5% 8%
1	C	269	17% 83% 10% 7%
1	D	269	8% 84% 6% 9%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8662 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-oxoacyl-(Acyl-carrier-protein) reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	249	Total	C	N	O	S	0	6	0
			1936	1224	350	353	9			
1	B	247	Total	C	N	O	S	0	14	0
			1969	1251	355	354	9			
1	C	250	Total	C	N	O	S	0	6	0
			1926	1219	352	347	8			
1	D	244	Total	C	N	O	S	0	9	0
			1913	1216	348	343	6			

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	MET	-	initiating methionine	UNP A9MA73
A	-7	ALA	-	expression tag	UNP A9MA73
A	-6	HIS	-	expression tag	UNP A9MA73
A	-5	HIS	-	expression tag	UNP A9MA73
A	-4	HIS	-	expression tag	UNP A9MA73
A	-3	HIS	-	expression tag	UNP A9MA73
A	-2	HIS	-	expression tag	UNP A9MA73
A	-1	HIS	-	expression tag	UNP A9MA73
A	0	MET	-	expression tag	UNP A9MA73
A	1	VAL	-	expression tag	UNP A9MA73
A	2	LEU	-	expression tag	UNP A9MA73
A	3	ASN	-	expression tag	UNP A9MA73
A	4	ASP	-	expression tag	UNP A9MA73
A	5	PRO	-	expression tag	UNP A9MA73
A	6	GLU	-	expression tag	UNP A9MA73
A	7	ALA	-	expression tag	UNP A9MA73
A	8	ARG	-	expression tag	UNP A9MA73
A	256	GLU	-	expression tag	UNP A9MA73
A	257	LEU	-	expression tag	UNP A9MA73
A	258	PRO	-	expression tag	UNP A9MA73
A	259	ASN	-	expression tag	UNP A9MA73

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	260	LYS	-	expression tag	UNP A9MA73
B	-8	MET	-	initiating methionine	UNP A9MA73
B	-7	ALA	-	expression tag	UNP A9MA73
B	-6	HIS	-	expression tag	UNP A9MA73
B	-5	HIS	-	expression tag	UNP A9MA73
B	-4	HIS	-	expression tag	UNP A9MA73
B	-3	HIS	-	expression tag	UNP A9MA73
B	-2	HIS	-	expression tag	UNP A9MA73
B	-1	HIS	-	expression tag	UNP A9MA73
B	0	MET	-	expression tag	UNP A9MA73
B	1	VAL	-	expression tag	UNP A9MA73
B	2	LEU	-	expression tag	UNP A9MA73
B	3	ASN	-	expression tag	UNP A9MA73
B	4	ASP	-	expression tag	UNP A9MA73
B	5	PRO	-	expression tag	UNP A9MA73
B	6	GLU	-	expression tag	UNP A9MA73
B	7	ALA	-	expression tag	UNP A9MA73
B	8	ARG	-	expression tag	UNP A9MA73
B	256	GLU	-	expression tag	UNP A9MA73
B	257	LEU	-	expression tag	UNP A9MA73
B	258	PRO	-	expression tag	UNP A9MA73
B	259	ASN	-	expression tag	UNP A9MA73
B	260	LYS	-	expression tag	UNP A9MA73
C	-8	MET	-	initiating methionine	UNP A9MA73
C	-7	ALA	-	expression tag	UNP A9MA73
C	-6	HIS	-	expression tag	UNP A9MA73
C	-5	HIS	-	expression tag	UNP A9MA73
C	-4	HIS	-	expression tag	UNP A9MA73
C	-3	HIS	-	expression tag	UNP A9MA73
C	-2	HIS	-	expression tag	UNP A9MA73
C	-1	HIS	-	expression tag	UNP A9MA73
C	0	MET	-	expression tag	UNP A9MA73
C	1	VAL	-	expression tag	UNP A9MA73
C	2	LEU	-	expression tag	UNP A9MA73
C	3	ASN	-	expression tag	UNP A9MA73
C	4	ASP	-	expression tag	UNP A9MA73
C	5	PRO	-	expression tag	UNP A9MA73
C	6	GLU	-	expression tag	UNP A9MA73
C	7	ALA	-	expression tag	UNP A9MA73
C	8	ARG	-	expression tag	UNP A9MA73
C	256	GLU	-	expression tag	UNP A9MA73
C	257	LEU	-	expression tag	UNP A9MA73

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	258	PRO	-	expression tag	UNP A9MA73
C	259	ASN	-	expression tag	UNP A9MA73
C	260	LYS	-	expression tag	UNP A9MA73
D	-8	MET	-	initiating methionine	UNP A9MA73
D	-7	ALA	-	expression tag	UNP A9MA73
D	-6	HIS	-	expression tag	UNP A9MA73
D	-5	HIS	-	expression tag	UNP A9MA73
D	-4	HIS	-	expression tag	UNP A9MA73
D	-3	HIS	-	expression tag	UNP A9MA73
D	-2	HIS	-	expression tag	UNP A9MA73
D	-1	HIS	-	expression tag	UNP A9MA73
D	0	MET	-	expression tag	UNP A9MA73
D	1	VAL	-	expression tag	UNP A9MA73
D	2	LEU	-	expression tag	UNP A9MA73
D	3	ASN	-	expression tag	UNP A9MA73
D	4	ASP	-	expression tag	UNP A9MA73
D	5	PRO	-	expression tag	UNP A9MA73
D	6	GLU	-	expression tag	UNP A9MA73
D	7	ALA	-	expression tag	UNP A9MA73
D	8	ARG	-	expression tag	UNP A9MA73
D	256	GLU	-	expression tag	UNP A9MA73
D	257	LEU	-	expression tag	UNP A9MA73
D	258	PRO	-	expression tag	UNP A9MA73
D	259	ASN	-	expression tag	UNP A9MA73
D	260	LYS	-	expression tag	UNP A9MA73

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	B	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	C	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	D	1	Total 48	C 21	N 7	O 17	P 3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	190	Total O 191 191	0	1
3	B	207	Total O 208 208	0	1
3	C	128	Total O 128 128	0	0
3	D	199	Total O 199 199	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3-oxoacyl-(Acyl-carrier-protein) reductase




- Molecule 1: 3-oxoacyl-(Acyl-carrier-protein) reductase



- Molecule 1: 3-oxoacyl-(Acyl-carrier-protein) reductase



- Molecule 1: 3-oxoacyl-(Acyl-carrier-protein) reductase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	76.57Å 75.60Å 99.18Å 90.00° 109.23° 90.00°	Depositor
Resolution (Å)	36.15 – 1.50 36.15 – 1.50	Depositor EDS
% Data completeness (in resolution range)	96.2 (36.15-1.50) 96.2 (36.15-1.50)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.21 (at 1.50Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.169 , 0.188 0.169 , 0.189	Depositor DCC
R_{free} test set	8084 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å ²)	15.6	Xtriage
Anisotropy	0.745	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8662	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/1990	0.72	0/2711
1	B	0.41	0/2050	0.71	0/2787
1	C	0.39	0/1983	0.71	0/2701
1	D	0.42	0/1979	0.70	0/2692
All	All	0.41	0/8002	0.71	0/10891

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1936	0	1928	12	0
1	B	1969	0	2000	8	0
1	C	1926	0	1920	18	0
1	D	1913	0	1928	9	0
2	A	48	0	25	2	0
2	B	48	0	25	2	0
2	C	48	0	24	5	0
2	D	48	0	25	1	0
3	A	191	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	208	0	0	1	0
3	C	128	0	0	2	0
3	D	199	0	0	1	0
All	All	8662	0	7875	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:230[B]:GLU:OE1	3:B:401:HOH:O	2.11	0.68
1:C:30:GLU:OE2	1:C:56:ARG:NH1	2.29	0.65
1:A:195:SER:O	1:A:197:ARG:N	2.32	0.61
1:C:134:LEU:HD12	1:C:135:PRO:HD2	1.85	0.58
1:A:9:MET:HE2	1:A:12:ASN:HB2	1.87	0.55
1:A:169:ASN:HD22	1:A:172:ARG:HH21	1.55	0.54
1:A:131[B]:ARG:NH1	3:A:404:HOH:O	2.34	0.54
1:D:85:ARG:NH1	3:D:403:HOH:O	2.39	0.54
1:D:18:THR:HA	1:D:42:HIS:HB3	1.90	0.53
1:B:178:LEU:HB3	1:B:182[A]:ILE:HB	1.91	0.53
1:C:18:THR:HA	1:C:42:HIS:HB3	1.89	0.53
1:A:223:ARG:NH1	3:A:406:HOH:O	2.41	0.53
1:D:153:LEU:HD21	1:D:165[B]:THR:HG21	1.90	0.53
1:A:44:ASN:HB3	2:A:300:NAP:O1X	2.10	0.52
1:A:19:GLY:HA3	2:A:300:NAP:H4B	1.94	0.50
1:B:18:THR:HA	1:B:42:HIS:HB3	1.95	0.49
1:C:72:GLU:OE1	2:C:300:NAP:N6A	2.46	0.49
1:C:23:ARG:NE	2:C:300:NAP:O2A	2.44	0.49
1:D:191:PRO:HD3	1:D:245:GLN:OE1	2.15	0.47
1:A:236:GLY:HA3	1:B:247:LEU:HD11	1.95	0.47
1:D:80[A]:LYS:NZ	1:D:84:ASP:OD1	2.48	0.46
1:C:195:SER:H	1:C:198:GLN:HB3	1.79	0.46
1:D:169:ASN:HD22	1:D:172:ARG:HH21	1.63	0.46
1:B:24:ILE:HB	2:B:300:NAP:H51N	1.98	0.46
1:B:191:PRO:HD3	1:B:245:GLN:OE1	2.17	0.45
2:C:300:NAP:H2N	2:C:300:NAP:O5D	2.18	0.44
1:B:44:ASN:HB2	2:B:300:NAP:C2A	2.48	0.44
1:A:191:PRO:HD3	1:A:245:GLN:OE1	2.18	0.43
1:C:247:LEU:HD11	1:D:236:GLY:HA3	2.00	0.43
1:C:214[A]:ARG:NH2	1:C:217:GLU:OE2	2.52	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:236:GLY:HA3	1:D:247:LEU:HD11	2.00	0.43
1:A:125[B]:ILE:HD13	1:A:125[B]:ILE:HA	1.93	0.42
1:C:109[A]:MET:HA	1:C:112:TRP:HB3	2.01	0.42
1:C:23:ARG:NH2	2:C:300:NAP:O1N	2.45	0.42
1:C:24:ILE:HB	2:C:300:NAP:H51N	2.02	0.42
1:C:109[B]:MET:HA	1:C:112:TRP:HB3	2.01	0.42
1:A:247:LEU:HD11	1:B:236:GLY:HA3	2.02	0.41
1:A:18:THR:HA	1:A:42:HIS:HB3	2.02	0.41
1:C:178:LEU:HB3	1:C:182:ILE:HB	2.02	0.41
1:C:30:GLU:OE1	3:C:401:HOH:O	2.22	0.41
1:C:83:SER:HA	1:C:87:GLY:O	2.20	0.41
1:C:255:ALA:HB2	3:C:433:HOH:O	2.20	0.41
1:D:44:ASN:HB2	2:D:300:NAP:C2A	2.51	0.41
1:C:10:VAL:HG21	1:C:35:HIS:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	253/269 (94%)	238 (94%)	14 (6%)	1 (0%)	30	12
1	B	257/269 (96%)	245 (95%)	11 (4%)	1 (0%)	30	12
1	C	254/269 (94%)	237 (93%)	16 (6%)	1 (0%)	30	12
1	D	249/269 (93%)	238 (96%)	10 (4%)	1 (0%)	30	12
All	All	1013/1076 (94%)	958 (95%)	51 (5%)	4 (0%)	30	12

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	146	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	146	ILE
1	B	146	ILE
1	D	146	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/223 (91%)	200 (99%)	2 (1%)	73	53
1	B	210/223 (94%)	209 (100%)	1 (0%)	86	75
1	C	198/223 (89%)	197 (100%)	1 (0%)	86	75
1	D	200/223 (90%)	198 (99%)	2 (1%)	73	53
All	All	810/892 (91%)	804 (99%)	6 (1%)	84	66

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	131[A]	ARG
1	A	131[B]	ARG
1	B	254	ILE
1	C	181	ARG
1	D	8	ARG
1	D	254	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	94	ASN
1	C	81	GLN
1	C	148	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAP	B	300	-	46,52,52	3.96	13 (28%)	61,80,80	1.83	9 (14%)
2	NAP	D	300	-	46,52,52	3.83	13 (28%)	61,80,80	1.75	8 (13%)
2	NAP	C	300	-	46,52,52	4.23	14 (30%)	61,80,80	1.69	7 (11%)
2	NAP	A	300	-	46,52,52	4.25	15 (32%)	61,80,80	2.10	12 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	B	300	-	-	0/31/67/67	0/5/5/5
2	NAP	D	300	-	-	0/31/67/67	0/5/5/5
2	NAP	C	300	-	-	4/31/67/67	0/5/5/5
2	NAP	A	300	-	-	0/31/67/67	0/5/5/5

All (55) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	300	NAP	O4D-C1D	15.82	1.61	1.40
2	A	300	NAP	O4D-C1D	15.33	1.61	1.40
2	C	300	NAP	O4B-C1B	15.32	1.61	1.40
2	A	300	NAP	O4B-C1B	15.17	1.60	1.40
2	D	300	NAP	O4D-C1D	14.59	1.60	1.40
2	B	300	NAP	O4D-C1D	14.52	1.59	1.40
2	B	300	NAP	O4B-C1B	13.94	1.59	1.40
2	D	300	NAP	O4B-C1B	13.05	1.58	1.40
2	A	300	NAP	PN-O3	10.36	1.70	1.59
2	B	300	NAP	PN-O3	9.60	1.69	1.59
2	C	300	NAP	PN-O3	9.31	1.69	1.59
2	D	300	NAP	PN-O3	9.29	1.69	1.59
2	C	300	NAP	PA-O3	7.06	1.67	1.59
2	A	300	NAP	O4B-C4B	-7.02	1.29	1.45
2	A	300	NAP	PA-O3	6.78	1.66	1.59
2	C	300	NAP	O4B-C4B	-6.47	1.30	1.45
2	B	300	NAP	O4B-C4B	-6.38	1.30	1.45
2	D	300	NAP	O4B-C4B	-6.16	1.31	1.45
2	A	300	NAP	C7N-N7N	5.87	1.43	1.33
2	A	300	NAP	O4D-C4D	-5.83	1.32	1.45
2	B	300	NAP	PA-O3	5.68	1.65	1.59
2	C	300	NAP	O4D-C4D	-5.57	1.32	1.45
2	C	300	NAP	C7N-N7N	5.56	1.43	1.33
2	B	300	NAP	C7N-N7N	5.34	1.42	1.33
2	D	300	NAP	C7N-N7N	5.19	1.42	1.33
2	D	300	NAP	PA-O3	5.16	1.65	1.59
2	D	300	NAP	O4D-C4D	-5.06	1.33	1.45
2	B	300	NAP	O4D-C4D	-5.03	1.33	1.45
2	B	300	NAP	P2B-O2B	4.53	1.67	1.59
2	C	300	NAP	P2B-O2B	4.37	1.67	1.59
2	B	300	NAP	O7N-C7N	-3.66	1.17	1.24
2	A	300	NAP	P2B-O2B	3.64	1.65	1.59
2	C	300	NAP	O7N-C7N	-3.63	1.17	1.24
2	D	300	NAP	O7N-C7N	-3.61	1.17	1.24
2	A	300	NAP	O7N-C7N	-3.59	1.17	1.24
2	D	300	NAP	P2B-O2B	3.54	1.65	1.59
2	A	300	NAP	C6A-N6A	3.00	1.44	1.34
2	C	300	NAP	C6A-N6A	2.98	1.44	1.34
2	B	300	NAP	C2N-N1N	2.97	1.38	1.35
2	B	300	NAP	C6A-N6A	2.81	1.44	1.34
2	A	300	NAP	O2D-C2D	2.79	1.49	1.43
2	D	300	NAP	C6A-N6A	2.72	1.43	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	300	NAP	O3D-C3D	-2.69	1.36	1.43
2	C	300	NAP	O2D-C2D	2.68	1.49	1.43
2	D	300	NAP	C2N-N1N	2.61	1.37	1.35
2	B	300	NAP	O2D-C2D	2.55	1.49	1.43
2	D	300	NAP	O2D-C2D	2.54	1.49	1.43
2	C	300	NAP	C2N-N1N	2.43	1.37	1.35
2	A	300	NAP	O3D-C3D	-2.35	1.37	1.43
2	B	300	NAP	O3D-C3D	-2.27	1.37	1.43
2	A	300	NAP	O3B-C3B	-2.15	1.37	1.43
2	C	300	NAP	PN-O5D	2.15	1.67	1.59
2	D	300	NAP	O3D-C3D	-2.07	1.37	1.43
2	A	300	NAP	C2A-N3A	2.01	1.35	1.32
2	A	300	NAP	PN-O5D	2.00	1.67	1.59

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	300	NAP	C4B-O4B-C1B	-9.25	101.45	109.92
2	B	300	NAP	C4B-O4B-C1B	-7.31	103.23	109.92
2	D	300	NAP	C4B-O4B-C1B	-6.78	103.72	109.92
2	C	300	NAP	N3A-C2A-N1A	-6.58	119.73	128.67
2	B	300	NAP	N3A-C2A-N1A	-6.39	120.00	128.67
2	A	300	NAP	N3A-C2A-N1A	-6.30	120.12	128.67
2	D	300	NAP	N3A-C2A-N1A	-6.12	120.36	128.67
2	A	300	NAP	C5A-C6A-N6A	5.52	128.72	120.31
2	C	300	NAP	C1B-N9A-C4A	-5.52	116.94	126.64
2	A	300	NAP	C1B-N9A-C4A	-5.46	117.05	126.64
2	B	300	NAP	C5A-C6A-N6A	5.19	128.21	120.31
2	C	300	NAP	C5A-C6A-N6A	5.15	128.16	120.31
2	B	300	NAP	C1B-N9A-C4A	-5.06	117.75	126.64
2	D	300	NAP	C1B-N9A-C4A	-5.01	117.84	126.64
2	D	300	NAP	C5A-C6A-N6A	4.66	127.41	120.31
2	A	300	NAP	N6A-C6A-N1A	-3.46	110.95	118.33
2	B	300	NAP	N6A-C6A-N1A	-3.26	111.37	118.33
2	C	300	NAP	N6A-C6A-N1A	-3.08	111.74	118.33
2	D	300	NAP	N6A-C6A-N1A	-3.07	111.78	118.33
2	C	300	NAP	C4B-O4B-C1B	-2.96	107.21	109.92
2	B	300	NAP	O7N-C7N-N7N	-2.73	118.67	122.62
2	A	300	NAP	C4D-O4D-C1D	-2.72	107.43	109.92
2	C	300	NAP	C6N-N1N-C2N	-2.64	119.63	121.88
2	A	300	NAP	C6N-N1N-C2N	-2.58	119.69	121.88
2	D	300	NAP	C4D-O4D-C1D	-2.55	107.59	109.92

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	300	NAP	O2N-PN-O3	2.49	114.02	107.27
2	A	300	NAP	O4B-C4B-C3B	-2.49	100.21	105.15
2	A	300	NAP	O7N-C7N-N7N	-2.30	119.29	122.62
2	B	300	NAP	C3B-C2B-C1B	2.25	107.11	102.81
2	A	300	NAP	P2B-O2B-C2B	-2.14	117.71	123.43
2	C	300	NAP	C3B-C2B-C1B	2.14	106.90	102.81
2	D	300	NAP	C3B-C2B-C1B	2.12	106.87	102.81
2	B	300	NAP	O7N-C7N-C3N	2.11	122.17	119.60
2	A	300	NAP	O4B-C1B-N9A	2.03	111.44	108.75
2	D	300	NAP	O7N-C7N-N7N	-2.02	119.70	122.62
2	B	300	NAP	O2N-PN-O3	2.00	112.68	107.27

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	300	NAP	C5B-O5B-PA-O1A
2	C	300	NAP	O4B-C4B-C5B-O5B
2	C	300	NAP	O4D-C4D-C5D-O5D
2	C	300	NAP	PN-O3-PA-O2A


There are no ring outliers.


4 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	300	NAP	2	0
2	D	300	NAP	1	0
2	C	300	NAP	5	0
2	A	300	NAP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	249/269 (92%)	0.27	11 (4%)	39	42	10, 24, 46, 66	6 (2%)
1	B	247/269 (91%)	-0.00	11 (4%)	39	41	7, 18, 42, 71	14 (5%)
1	C	250/269 (92%)	0.97	45 (18%)	4	3	12, 35, 58, 77	6 (2%)
1	D	244/269 (90%)	0.09	22 (9%)	17	16	7, 18, 48, 74	9 (3%)
All	All	990/1076 (92%)	0.33	89 (8%)	17	16	7, 23, 52, 77	35 (3%)

All (89) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	255	ALA	6.0
1	C	7	ALA	5.9
1	B	7	ALA	5.5
1	D	200	PRO	5.4
1	D	202	ASP	4.7
1	C	67	VAL	4.5
1	D	10	VAL	4.5
1	C	43	CYS	4.5
1	D	254	ILE	4.4
1	C	196	GLU	4.3
1	C	255	ALA	4.3
1	B	254	ILE	4.0
1	D	8	ARG	3.9
1	C	195	SER	3.8
1	C	70	ASP	3.8
1	C	75	VAL	3.8
1	A	254	ILE	3.8
1	D	107	LEU	3.7
1	C	82	ALA	3.6
1	C	6	GLU	3.6
1	C	254	ILE	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	69	ALA	3.5
1	C	66	VAL	3.5
1	C	10	VAL	3.5
1	C	44	ASN	3.4
1	C	45	ARG	3.2
1	C	200	PRO	3.2
1	D	105	GLY	3.1
1	C	74	ASP	3.1
1	D	9	MET	3.1
1	C	194	PRO	3.1
1	C	65	CYS	3.0
1	C	11	ALA	3.0
1	D	11	ALA	3.0
1	D	193	LEU	3.0
1	D	203	PHE	3.0
1	C	62	GLY	3.0
1	C	73	GLY	3.0
1	C	110	ALA	3.0
1	D	191	PRO	2.9
1	C	77	GLY	2.8
1	B	256	GLU	2.8
1	C	134	LEU	2.8
1	C	71	LEU	2.7
1	D	194	PRO	2.7
1	C	193	LEU	2.6
1	D	192	THR	2.6
1	C	9	MET	2.6
1	B	10	VAL	2.5
1	C	47	LEU	2.5
1	C	46	SER	2.5
1	C	157	PHE	2.5
1	D	256	GLU	2.5
1	C	61	GLY	2.5
1	D	76	ARG	2.5
1	B	195	SER	2.5
1	B	106	ALA	2.5
1	A	196	GLU	2.5
1	A	111	LEU	2.4
1	B	200	PRO	2.4
1	B	255	ALA	2.4
1	C	54	ALA	2.3
1	A	108	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	77	GLY	2.3
1	C	79	VAL	2.3
1	C	76	ARG	2.3
1	A	157	PHE	2.3
1	C	64	ALA	2.3
1	B	99	PHE	2.2
1	A	195	SER	2.2
1	A	98	LEU	2.2
1	D	136	GLU	2.2
1	A	75	VAL	2.2
1	C	52	ALA	2.1
1	C	203	PHE	2.1
1	C	83	SER	2.1
1	C	107	LEU	2.1
1	B	9	MET	2.1
1	B	104	VAL	2.1
1	C	128	GLU	2.1
1	D	201	GLU	2.1
1	A	193	LEU	2.1
1	C	36	GLY	2.0
1	C	57	ILE	2.0
1	D	137	ASP	2.0
1	C	109[A]	MET	2.0
1	A	110	ALA	2.0
1	D	106	ALA	2.0
1	D	104	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAP	C	300	48/48	0.83	0.15	16,33,43,47	0
2	NAP	A	300	48/48	0.90	0.12	16,27,38,47	0
2	NAP	B	300	48/48	0.96	0.07	16,22,31,39	0
2	NAP	D	300	48/48	0.96	0.08	16,20,31,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around NAP A 300:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAP B 300:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.