



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:13 PM UTC

PDB ID : 5GL8 / pdb\_00005gl8  
Title : The crystal structure of Nattokinase fragments  
Authors : Liu, W.  
Deposited on : 2016-07-10  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	
1	B	275	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4747 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Subtilisin NAT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	275	1995	1244	341	406	4	0	8	0
1	B	275	1992	1239	341	408	4	0	7	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	130	SER	THR	conflict	UNP P35835
B	130	SER	THR	conflict	UNP P35835

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Ca	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	380	Total	O	0	0
			380	380		
3	B	378	Total	O	0	0
			378	378		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Subtilisin NAT



- Molecule 1: Subtilisin NAT



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.28Å 73.96Å 67.81Å 90.00° 97.47° 90.00°	Depositor
Resolution (Å)	50.00 – 1.80 50.00 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.6 (50.00-1.80) 99.6 (50.00-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.28 (at 1.79Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.231 , 0.291 0.244 , 0.305	Depositor DCC
$R_{free}$ test set	2342 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	9.0	Xtrriage
Anisotropy	0.211	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 41.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	4747	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	10.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.21 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.7535e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.36	9/2052 (0.4%)	1.21	3/2800 (0.1%)
1	B	1.36	8/2040 (0.4%)	1.16	1/2784 (0.0%)
All	All	1.36	17/4092 (0.4%)	1.19	4/5584 (0.1%)

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	60	ASP	C-O	-9.61	1.13	1.24
1	A	198	VAL	C-O	7.20	1.32	1.23
1	B	60	ASP	C-O	-6.99	1.16	1.24
1	A	200	ALA	C-O	-6.33	1.18	1.24
1	A	21	TYR	C-O	-6.04	1.17	1.24
1	A	107	ILE	C-O	-5.76	1.17	1.24
1	A	258	GLY	N-CA	5.47	1.51	1.44
1	B	186	ARG	C-O	5.45	1.30	1.23
1	B	241	TRP	N-CA	5.45	1.52	1.46
1	A	209	LEU	C-O	-5.43	1.17	1.24
1	B	68	VAL	CA-CB	5.23	1.61	1.54
1	B	209	LEU	C-O	-5.18	1.17	1.24
1	A	11	ILE	CA-CB	5.10	1.61	1.54
1	B	73	ALA	N-CA	5.09	1.51	1.46
1	B	143	VAL	C-O	5.09	1.29	1.24
1	B	51	VAL	CA-C	5.06	1.57	1.53
1	A	186	ARG	C-O	5.01	1.29	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	209	LEU	CA-C-N	-6.91	112.87	119.85
1	A	209	LEU	C-N-CA	-6.91	112.87	119.85

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	45	ARG	CG-CD-NE	5.78	124.70	112.00
1	B	45	ARG	NE-CZ-NH1	-5.43	116.07	121.50

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1995	0	1973	64	0
1	B	1992	0	1954	37	0
2	B	2	0	0	0	0
3	A	380	0	0	34	3
3	B	378	0	0	22	3
All	All	4747	0	3927	101	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:SER:O	1:A:162[B]:THR:HG22	1.53	1.07
1:B:34:GLY:C	1:B:35[B]:ILE:HD12	1.86	1.00
1:A:34:GLY:C	1:A:35[A]:ILE:HD12	1.94	0.93
1:B:27[B]:LYS:HE3	3:B:466:HOH:O	1.68	0.92
1:A:25:ASN:ND2	3:A:301:HOH:O	2.04	0.90
1:A:57:PRO:HA	1:A:94[A]:LYS:HZ3	1.40	0.87
1:A:45:ARG:NE	1:A:89:SER:OG	2.12	0.81
1:B:262:TYR:HB3	3:B:665:HOH:O	1.79	0.81
1:A:33:SER:HB3	3:A:394:HOH:O	1.80	0.81
1:A:113:TRP:CZ2	3:A:464:HOH:O	2.34	0.81
1:A:206:GLN:NE2	3:A:303:HOH:O	2.14	0.81
1:A:56:ASN:ND2	1:A:59[A]:GLN:HG3	1.96	0.80
1:B:116:SER:HB3	3:B:598:HOH:O	1.81	0.79

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:148:VAL:HG23	3:A:497:HOH:O	1.81	0.79
1:B:150:ALA:HB2	3:B:656:HOH:O	1.85	0.77
1:B:31:ILE:HD13	3:B:622:HOH:O	1.85	0.76
1:A:210:PRO:HB2	3:A:524:HOH:O	1.85	0.76
1:A:106:TRP:CH2	3:A:549:HOH:O	2.38	0.75
1:B:181:ASN:HD21	1:B:185:GLN:HE21	1.36	0.74
1:A:106:TRP:CZ3	3:A:549:HOH:O	2.40	0.74
1:A:57:PRO:HA	1:A:94[A]:LYS:NZ	2.02	0.74
1:A:37:SER:HB2	1:A:44[B]:VAL:CG2	2.19	0.73
1:A:243:ASN:HB3	3:A:497:HOH:O	1.91	0.70
1:A:246:VAL:HG12	3:A:365:HOH:O	1.91	0.70
1:B:35[B]:ILE:HD12	1:B:35[B]:ILE:N	2.08	0.69
1:B:34:GLY:O	1:B:35[B]:ILE:HD12	1.95	0.67
1:B:31:ILE:HG23	3:B:622:HOH:O	1.94	0.67
1:B:186:ARG:HD3	3:B:665:HOH:O	1.96	0.66
1:A:35[A]:ILE:HD13	1:A:92:ALA:HA	1.77	0.65
1:B:230:ALA:O	1:B:234:ILE:HG13	1.95	0.65
1:A:45:ARG:HE	1:A:89:SER:CB	2.10	0.64
1:A:32:ASP:OD1	1:A:64:HIS:HD2	1.82	0.63
1:B:186:ARG:CD	3:B:665:HOH:O	2.47	0.62
1:A:37:SER:HB2	1:A:44[B]:VAL:HG21	1.81	0.62
1:A:37:SER:CB	1:A:44[B]:VAL:HG21	2.29	0.62
1:B:178:GLY:C	3:B:443:HOH:O	2.45	0.60
1:A:159:SER:O	1:A:162[B]:THR:CG2	2.42	0.60
1:A:106:TRP:CZ2	3:A:549:HOH:O	2.54	0.59
1:A:51:VAL:HA	3:A:549:HOH:O	2.03	0.58
1:A:185:GLN:HB3	3:A:543:HOH:O	2.04	0.58
1:A:113:TRP:CH2	3:A:464:HOH:O	2.55	0.57
1:A:206:GLN:HE21	1:A:216:ALA:HB2	1.69	0.56
1:A:106:TRP:CE3	3:A:549:HOH:O	2.56	0.56
1:A:64:HIS:HB3	3:A:394:HOH:O	2.06	0.56
1:A:59[B]:GLN:NE2	1:A:94[B]:LYS:NZ	2.53	0.56
1:B:249:ARG:CZ	3:B:423:HOH:O	2.53	0.55
1:B:115[B]:ILE:HD11	1:B:142:ALA:HB2	1.89	0.55
1:A:230:ALA:O	1:A:234:ILE:HG13	2.07	0.55
1:B:56:ASN:ND2	1:B:59:GLN:OE1	2.39	0.55
1:A:77:ASN:HD22	1:A:77:ASN:C	2.15	0.54
1:B:223:ALA:HB3	3:B:443:HOH:O	2.07	0.54
1:B:27[B]:LYS:CE	3:B:466:HOH:O	2.41	0.54
1:B:35[B]:ILE:N	1:B:35[B]:ILE:CD1	2.69	0.54
1:B:27[B]:LYS:HD2	1:B:89:SER:HB2	1.90	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:255:THR:HG22	3:A:305:HOH:O	2.07	0.53
1:B:35[B]:ILE:HD13	1:B:92:ALA:CB	2.38	0.53
1:B:93:VAL:HG11	3:B:622:HOH:O	2.09	0.53
1:B:110:GLY:HA3	3:B:622:HOH:O	2.08	0.53
1:B:16:LEU:HD11	3:B:600:HOH:O	2.08	0.53
1:A:265[A]:LYS:NZ	3:A:312:HOH:O	2.41	0.52
1:B:234:ILE:HA	3:B:600:HOH:O	2.09	0.52
1:A:248:ASP:HB2	3:A:571:HOH:O	2.09	0.51
1:A:35[A]:ILE:HD13	1:A:92:ALA:CB	2.40	0.51
1:B:111:ILE:O	1:B:115[B]:ILE:HD13	2.11	0.51
1:B:181:ASN:HD21	1:B:185:GLN:NE2	2.06	0.51
1:A:259:ASN:ND2	3:A:311:HOH:O	2.38	0.50
1:A:106:TRP:CE2	3:A:549:HOH:O	2.65	0.50
1:A:77:ASN:ND2	3:A:307:HOH:O	2.44	0.50
1:B:35[B]:ILE:HD13	1:B:92:ALA:HA	1.94	0.49
1:A:162[B]:THR:HG23	1:A:162[B]:THR:O	2.13	0.49
1:A:35[A]:ILE:HD13	1:A:92:ALA:CA	2.42	0.49
1:A:157:GLY:HA3	3:A:341:HOH:O	2.13	0.48
1:A:243:ASN:HA	3:A:497:HOH:O	2.12	0.48
1:B:274:ALA:HB1	3:B:600:HOH:O	2.13	0.47
1:A:265[A]:LYS:HD3	3:A:552:HOH:O	2.13	0.47
1:A:106:TRP:CD2	3:A:549:HOH:O	2.68	0.46
1:A:243:ASN:CA	3:A:497:HOH:O	2.63	0.46
1:A:52:PRO:HD2	3:A:549:HOH:O	2.16	0.45
1:A:257:LEU:CD2	3:A:305:HOH:O	2.64	0.45
1:B:93:VAL:CG1	3:B:622:HOH:O	2.64	0.45
1:B:221:SER:O	1:B:225:PRO:HD2	2.16	0.45
1:A:64:HIS:HE1	1:A:221:SER:OG	1.98	0.45
1:A:159:SER:HB3	1:A:162[B]:THR:CG2	2.47	0.45
1:B:274:ALA:CB	3:B:600:HOH:O	2.64	0.45
1:A:255:THR:CG2	3:A:305:HOH:O	2.64	0.45
1:B:31:ILE:CD1	3:B:622:HOH:O	2.57	0.44
1:A:59[A]:GLN:NE2	3:A:319:HOH:O	2.51	0.44
1:B:220:THR:HG22	3:B:443:HOH:O	2.17	0.43
1:A:35[A]:ILE:CD1	3:A:433:HOH:O	2.67	0.43
1:A:206:GLN:NE2	1:A:216:ALA:HB2	2.34	0.42
1:A:34:GLY:O	1:A:35[A]:ILE:HD12	2.17	0.42
1:A:249:ARG:HD3	1:A:273:ALA:O	2.19	0.42
1:A:59[A]:GLN:HE21	1:A:59[A]:GLN:HB3	1.56	0.41
1:B:155:ASN:HA	1:B:189:PHE:O	2.21	0.41
1:A:45:ARG:CZ	1:A:89:SER:OG	2.67	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:243:ASN:CB	3:A:497:HOH:O	2.60	0.41
1:A:255:THR:HB	3:A:305:HOH:O	2.21	0.41
1:A:224:THR:N	1:A:225:PRO:HD2	2.36	0.40
1:B:257:LEU:HD23	3:B:574:HOH:O	2.21	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:371:HOH:O	3:B:532:HOH:O[2_656]	1.59	0.61
3:A:541:HOH:O	3:B:745:HOH:O[2_656]	2.11	0.09
3:A:621:HOH:O	3:B:408:HOH:O[2_656]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	281/275 (102%)	269 (96%)	12 (4%)	0	100	100
1	B	280/275 (102%)	267 (95%)	13 (5%)	0	100	100
All	All	561/550 (102%)	536 (96%)	25 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/208 (104%)	211 (98%)	5 (2%)	44	33
1	B	215/208 (103%)	215 (100%)	0	100	100
All	All	431/416 (104%)	426 (99%)	5 (1%)	68	57

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35[A]	ILE
1	A	35[B]	ILE
1	A	77	ASN
1	A	101	SER
1	A	275	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	HIS
1	A	77	ASN
1	A	206	GLN
1	A	271	GLN
1	B	185	GLN
1	B	271	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	275/275 (100%)	0.60	8 (2%) 53 53	3, 8, 15, 28	8 (2%)
1	B	275/275 (100%)	0.59	10 (3%) 46 46	3, 8, 16, 25	7 (2%)
All	All	550/550 (100%)	0.59	18 (3%) 49 49	3, 8, 16, 28	15 (2%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	35[A]	ILE	3.5
1	A	104	TYR	3.3
1	B	275	GLN	3.2
1	A	275	GLN	2.8
1	A	196	LEU	2.8
1	B	196	LEU	2.8
1	A	263	TYR	2.4
1	A	51	VAL	2.4
1	B	263	TYR	2.3
1	B	11	ILE	2.2
1	B	96	LEU	2.2
1	B	104	TYR	2.1
1	A	99	THR	2.1
1	B	257	LEU	2.1
1	B	273	ALA	2.1
1	A	45	ARG	2.1
1	A	162[A]	THR	2.0
1	B	194	SER	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CA	B	301	1/1	0.97	0.09	22,22,22,22	0
2	CA	B	302	1/1	1.00	0.01	6,6,6,6	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.