



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 4, 2025 – 08:47 AM EDT

PDB ID : 5I4D / pdb\_00005i4d  
Title : 1.75 Angstrom Crystal Structure of Superantigen-like Protein, Exotoxin from Staphylococcus aureus, in Complex with Sialyl-LewisX.  
Authors : Minasov, G.; Nocadello, S.; Shuvalova, L.; Filippova, E.V.; Halavaty, A.; Dubrovskaya, I.; Bagnoli, F.; Falugi, F.; Bottomley, M.; Grandi, G.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases (CSGID)  
Deposited on : 2016-02-11  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.45.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2888 (1.76-1.76)
Clashscore	180529	3097 (1.76-1.76)
Ramachandran outliers	177936	3072 (1.76-1.76)
Sidechain outliers	177891	3072 (1.76-1.76)
RSRZ outliers	164620	2887 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	
1	B	343	
2	C	4	
2	D	4	

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 4379 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Superantigen-like protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	192	1787	1154	298	326	9	0	25	0
1	B	193	1746	1122	294	319	11	0	19	0

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	14	MET	-	initiating methionine	UNP A0A0D6GBT0
A	15	GLY	-	expression tag	UNP A0A0D6GBT0
A	16	SER	-	expression tag	UNP A0A0D6GBT0
A	17	SER	-	expression tag	UNP A0A0D6GBT0
A	18	HIS	-	expression tag	UNP A0A0D6GBT0
A	19	HIS	-	expression tag	UNP A0A0D6GBT0
A	20	HIS	-	expression tag	UNP A0A0D6GBT0
A	21	HIS	-	expression tag	UNP A0A0D6GBT0
A	22	HIS	-	expression tag	UNP A0A0D6GBT0
A	23	HIS	-	expression tag	UNP A0A0D6GBT0
A	24	GLU	-	expression tag	UNP A0A0D6GBT0
A	25	ASN	-	expression tag	UNP A0A0D6GBT0
A	26	LEU	-	expression tag	UNP A0A0D6GBT0
A	27	TYR	-	expression tag	UNP A0A0D6GBT0
A	28	PHE	-	expression tag	UNP A0A0D6GBT0
A	29	GLN	-	expression tag	UNP A0A0D6GBT0
A	30	GLY	-	expression tag	UNP A0A0D6GBT0
B	14	MET	-	initiating methionine	UNP A0A0D6GBT0
B	15	GLY	-	expression tag	UNP A0A0D6GBT0
B	16	SER	-	expression tag	UNP A0A0D6GBT0
B	17	SER	-	expression tag	UNP A0A0D6GBT0
B	18	HIS	-	expression tag	UNP A0A0D6GBT0
B	19	HIS	-	expression tag	UNP A0A0D6GBT0
B	20	HIS	-	expression tag	UNP A0A0D6GBT0
B	21	HIS	-	expression tag	UNP A0A0D6GBT0

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	22	HIS	-	expression tag	UNP A0A0D6GBT0
B	23	HIS	-	expression tag	UNP A0A0D6GBT0
B	24	GLU	-	expression tag	UNP A0A0D6GBT0
B	25	ASN	-	expression tag	UNP A0A0D6GBT0
B	26	LEU	-	expression tag	UNP A0A0D6GBT0
B	27	TYR	-	expression tag	UNP A0A0D6GBT0
B	28	PHE	-	expression tag	UNP A0A0D6GBT0
B	29	GLN	-	expression tag	UNP A0A0D6GBT0
B	30	GLY	-	expression tag	UNP A0A0D6GBT0

- Molecule 2 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-alpha-D-glucopyranose.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
2	C	4	Total	C	N	O	0	0	0
			56	31	2	23			
2	D	4	Total	C	N	O	0	3	0
			66	37	2	27			

- Molecule 3 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>4</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 4 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C<sub>8</sub>H<sub>18</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	8	5		
4	B	1	Total	C	O	0	0
			13	8	5		

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	1
			15	8	1	6		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	326	Total 349	O 349	0	33
6	B	315	Total 324	O 324	0	22





3W51
GAL2
SIA3
FUC4

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.97Å 166.88Å 34.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.94 – 1.75 29.94 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.6 (29.94-1.75) 99.6 (29.94-1.75)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.11 (at 1.75Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.161 , 0.202 0.170 , 0.210	Depositor DCC
$R_{free}$ test set	2563 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.6	Xtrriage
Anisotropy	0.170	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 46.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4379	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.15% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, FUC, GAL, SIA, NDG, PG4, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.67	0/1838	0.83	1/2455 (0.0%)
1	B	0.68	0/1785	0.82	1/2379 (0.0%)
All	All	0.68	0/3623	0.83	2/4834 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	235	GLU	N-CA-C	5.39	117.88	109.52
1	A	235	GLU	CB-CA-C	-5.19	99.87	109.37

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1787	0	1837	15	0
1	B	1746	0	1805	5	0
2	C	56	0	46	0	0
2	D	66	0	47	0	0
3	A	10	0	14	0	0
4	A	13	0	18	0	0
4	B	13	0	18	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	15	0	10	0	0
6	A	349	0	0	3	0
6	B	324	0	0	2	0
All	All	4379	0	3795	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:260[B]:LYS:HA	1:A:260[B]:LYS:HE3	1.67	0.74
1:A:210:ILE:CD1	1:A:230[B]:VAL:HG11	2.22	0.69
1:A:260[B]:LYS:NZ	1:A:282[B]:MET:HG3	2.10	0.66
1:A:260[B]:LYS:HZ1	1:A:282[B]:MET:HG3	1.62	0.64
1:A:210:ILE:HD11	1:A:230[B]:VAL:HG11	1.80	0.62
1:A:260[B]:LYS:HZ3	1:A:282[B]:MET:CG	2.18	0.57
1:A:240[B]:GLN:HA	1:A:240[B]:GLN:OE1	2.05	0.56
1:B:246:VAL:HG11	1:B:336[A]:GLU:HG3	1.89	0.54
1:A:260[B]:LYS:NZ	1:A:282[B]:MET:CG	2.70	0.54
1:A:210:ILE:HD13	1:A:230[B]:VAL:HG11	1.88	0.52
1:A:269:ASP:HA	1:A:311:MET:SD	2.54	0.48
1:B:347[A]:ASN:ND2	6:B:501:HOH:O	2.46	0.47
1:A:167:LYS:NZ	6:A:503:HOH:O	2.44	0.47
1:A:255:LYS:NZ	6:A:506:HOH:O	2.49	0.45
1:B:189:MET:HE1	1:B:212:LEU:HD11	1.99	0.45
1:B:273:MET:HA	1:B:273:MET:HE2	1.99	0.44
1:B:184[A]:LYS:HB3	6:B:650[A]:HOH:O	2.17	0.43
1:A:276[B]:ARG:C	1:A:277[B]:ASP:OD1	2.63	0.42
1:A:193:TRP:CZ2	1:A:198[B]:PHE:CE1	3.09	0.40
1:A:276[B]:ARG:NH2	6:A:510:HOH:O	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	215/343 (63%)	208 (97%)	7 (3%)	0	100	100
1	B	210/343 (61%)	205 (98%)	5 (2%)	0	100	100
All	All	425/686 (62%)	413 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/313 (64%)	202 (100%)	0	100	100
1	B	197/313 (63%)	194 (98%)	3 (2%)	60	45
All	All	399/626 (64%)	396 (99%)	3 (1%)	84	71

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	242[A]	LYS
1	B	242[B]	LYS
1	B	356	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	258	ASN
1	A	347	ASN
1	A	350	ASN
1	B	271	GLN
1	B	354	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NDG	C	1	2	15,15,15	0.52	0	21,21,21	0.99	1 (4%)
2	GAL	C	2	2	11,11,12	0.43	0	15,15,17	0.95	0
2	SIA	C	3	2	20,20,21	0.64	1 (5%)	21,28,31	1.05	4 (19%)
2	FUC	C	4	2	10,10,11	0.48	0	14,14,16	0.71	0
2	NDG	D	1[A]	2	15,15,15	0.44	0	21,21,21	0.81	0
2	GAL	D	2[A]	2	11,11,12	0.49	0	15,15,17	0.94	1 (6%)
2	GAL	D	2[B]	5	11,11,12	0.49	0	15,15,17	0.93	0
2	SIA	D	3	2	20,20,21	0.60	0	21,28,31	1.12	2 (9%)
2	FUC	D	4[A]	2	10,10,11	0.43	0	14,14,16	0.72	0
2	FUC	D	4[B]	5	10,10,11	0.44	0	14,14,16	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDG	C	1	2	-	0/6/26/26	0/1/1/1
2	GAL	C	2	2	-	0/2/19/22	0/1/1/1
2	SIA	C	3	2	-	0/18/34/38	0/1/1/1
2	FUC	C	4	2	-	-	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDG	D	1[A]	2	-	0/6/26/26	0/1/1/1
2	GAL	D	2[A]	2	-	0/2/19/22	0/1/1/1
2	GAL	D	2[B]	5	-	0/2/19/22	0/1/1/1
2	SIA	D	3	2	-	0/18/34/38	0/1/1/1
2	FUC	D	4[A]	2	-	-	0/1/1/1
2	FUC	D	4[B]	5	-	-	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	3	SIA	O1B-C1	-2.11	1.23	1.30

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	3	SIA	O6-C2-C3	-2.98	106.54	110.56
2	D	3	SIA	O1B-C1-C2	2.45	119.08	112.71
2	C	3	SIA	O6-C2-C1	2.36	112.18	107.72
2	C	3	SIA	O1B-C1-C2	2.29	118.67	112.71
2	C	1	NDG	O1-C1-C2	2.23	113.84	109.22
2	D	2[A]	GAL	C1-O5-C5	2.19	115.12	112.19
2	C	3	SIA	O6-C2-C3	-2.16	107.65	110.56
2	C	3	SIA	O1A-C1-C2	-2.01	118.51	122.85


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PGE	A	405	-	9,9,9	0.57	0	8,8,8	0.45	0
4	PG4	B	406	-	12,12,12	0.51	0	11,11,11	0.32	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NAG	B	402[B]	2	15,15,15	0.42	0	21,21,21	1.05	2 (9%)
4	PG4	A	406	-	12,12,12	0.49	0	11,11,11	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGE	A	405	-	-	3/7/7/7	-
4	PG4	B	406	-	-	2/10/10/10	-
5	NAG	B	402[B]	2	-	0/6/26/26	0/1/1/1
4	PG4	A	406	-	-	5/10/10/10	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	402[B]	NAG	O5-C1-C2	2.75	112.28	109.52
5	B	402[B]	NAG	C1-C2-N2	-2.17	108.22	110.73

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	406	PG4	O1-C1-C2-O2
3	A	405	PGE	O3-C5-C6-O4
4	A	406	PG4	O1-C1-C2-O2
4	A	406	PG4	O4-C7-C8-O5
3	A	405	PGE	O2-C3-C4-O3
4	B	406	PG4	C8-C7-O4-C6
4	A	406	PG4	C3-C4-O3-C5
4	A	406	PG4	O2-C3-C4-O3
3	A	405	PGE	C3-C4-O3-C5
4	A	406	PG4	C5-C6-O4-C7

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	192/343 (55%)	-0.25	3 (1%) 70 76	8, 18, 37, 74	25 (13%)
1	B	193/343 (56%)	-0.17	3 (1%) 70 76	8, 19, 43, 62	19 (9%)
All	All	385/686 (56%)	-0.21	6 (1%) 70 76	8, 18, 39, 74	44 (11%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	165	THR	5.3
1	B	164	MET	3.5
1	A	166	PRO	3.4
1	B	273	MET	3.3
1	A	167	LYS	2.3
1	B	165	THR	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NDG	C	1	15/15	0.94	0.07	15,18,22,22	0
2	NDG	D	1[A]	15/15	0.95	0.07	15,17,17,18	15
2	FUC	D	4[A]	10/11	0.96	0.06	18,19,20,22	1


*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FUC	D	4[B]	10/11	0.96	0.06	18,19,20,22	1
2	SIA	D	3	20/21	0.97	0.05	13,13,17,19	0
2	FUC	C	4	10/11	0.97	0.05	15,17,19,19	0
2	SIA	C	3	20/21	0.97	0.04	12,14,16,17	0
2	GAL	C	2	11/12	0.98	0.04	13,14,15,15	0
2	GAL	D	2[A]	11/12	0.98	0.05	13,14,15,15	9
2	GAL	D	2[B]	11/12	0.98	0.05	14,14,15,15	9

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	PGE	A	405	10/10	0.69	0.18	41,54,68,71	0
4	PG4	A	406	13/13	0.84	0.13	48,50,60,60	0
4	PG4	B	406	13/13	0.89	0.10	37,42,46,49	0
5	NAG	B	402[B]	15/15	0.94	0.08	15,18,22,24	15

## 6.5 Other polymers [i](#)

There are no such residues in this entry.