



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 03:50 AM EDT

PDB ID : 5LOY
Title : Helical Assembly of a Designed Anbu Protein
Authors : Fuchs, A.C.D.; Albrecht, R.; Martin, J.; Hartmann, M.D.
Deposited on : 2016-08-10
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	240	 3% 92% 5% 5%
1	B	240	 2% 92% 5% 5%
1	C	240	 3% 90% 5% 5%
1	D	240	 3% 92% 5% 5%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7504 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Designed Anbu Protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	229	1784	1135	306	337	6	0	0	0
1	B	229	1780	1133	306	335	6	0	0	0
1	C	229	1788	1137	306	339	6	0	0	0
1	D	229	1781	1134	306	335	6	0	0	0

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	89	89	89	0	0
2	B	82	82	82	0	0
2	C	99	99	99	0	0
2	D	101	101	101	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Designed Anbu Protein

Chain A: 




- Molecule 1: Designed Anbu Protein

Chain B: 





- Molecule 1: Designed Anbu Protein

Chain C: 



- Molecule 1: Designed Anbu Protein

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	170.74Å 170.74Å 92.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.70 – 2.50 38.73 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (38.70-2.50) 99.8 (38.73-2.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.181 , 0.222 0.185 , 0.220	Depositor DCC
R_{free} test set	2381 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	45.3	Xtrriage
Anisotropy	0.040	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7504	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	0/1821	0.75	0/2474
1	B	0.61	0/1817	0.74	0/2469
1	C	0.63	0/1825	0.78	0/2479
1	D	0.63	0/1818	0.75	0/2470
All	All	0.62	0/7281	0.76	0/9892

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1784	0	1761	3	0
1	B	1780	0	1757	4	0
1	C	1788	0	1765	5	0
1	D	1781	0	1759	8	0
2	A	89	0	0	0	0
2	B	82	0	0	0	0
2	C	99	0	0	3	0
2	D	101	0	0	3	0
All	All	7504	0	7042	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:198:ARG:HD2	2:D:377:HOH:O	1.92	0.68
1:C:31:ARG:CD	2:C:333:HOH:O	2.56	0.53
1:D:206:ARG:HD2	2:D:309:HOH:O	2.10	0.51
1:B:239:TRP:CZ3	1:D:205:HIS:HB3	2.47	0.50
1:B:239:TRP:CE3	1:D:205:HIS:HB3	2.51	0.45
1:A:205:HIS:HB3	1:C:239:TRP:CZ3	2.51	0.45
1:A:228:ARG:HD2	1:A:228:ARG:HA	1.85	0.45
1:A:238:PRO:O	1:A:239:TRP:C	2.55	0.44
1:B:205:HIS:CE1	1:D:239:TRP:CD1	3.07	0.43
1:C:163:ASP:HB2	2:C:382:HOH:O	2.18	0.43
2:C:349:HOH:O	1:D:28:SER:HB3	2.17	0.42
1:D:161:THR:HB	1:D:162:PRO:HD2	2.02	0.42
1:D:198:ARG:CD	2:D:377:HOH:O	2.61	0.42
1:C:138:THR:HB	1:C:139:PRO:HD2	2.02	0.42
1:C:153:LYS:N	1:C:154:PRO:CD	2.83	0.42
1:B:161:THR:HB	1:B:162:PRO:HD2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	225/240 (94%)	223 (99%)	2 (1%)	0	100	100
1	B	225/240 (94%)	223 (99%)	2 (1%)	0	100	100
1	C	225/240 (94%)	223 (99%)	2 (1%)	0	100	100
1	D	225/240 (94%)	221 (98%)	4 (2%)	0	100	100
All	All	900/960 (94%)	890 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/203 (94%)	187 (98%)	4 (2%)	53	78
1	B	190/203 (94%)	185 (97%)	5 (3%)	46	72
1	C	192/203 (95%)	187 (97%)	5 (3%)	46	72
1	D	190/203 (94%)	188 (99%)	2 (1%)	73	89
All	All	763/812 (94%)	747 (98%)	16 (2%)	53	78

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	ASN
1	A	41	GLU
1	A	206	ARG
1	A	218	MET
1	B	21	ASN
1	B	41	GLU
1	B	206	ARG
1	B	218	MET
1	B	228	ARG
1	C	21	ASN
1	C	41	GLU
1	C	206	ARG
1	C	228	ARG
1	C	229	GLN
1	D	21	ASN
1	D	206	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	222	GLN
1	C	222	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	229/240 (95%)	-0.14	7 (3%) 49 52	29, 43, 82, 130	0
1	B	229/240 (95%)	-0.14	5 (2%) 62 65	29, 44, 79, 134	0
1	C	229/240 (95%)	-0.22	8 (3%) 44 47	28, 41, 83, 141	0
1	D	229/240 (95%)	-0.16	7 (3%) 49 52	26, 42, 84, 132	0
All	All	916/960 (95%)	-0.16	27 (2%) 51 55	26, 43, 83, 141	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	66	LEU	5.6
1	D	239	TRP	5.5
1	C	107	ASP	5.1
1	C	69	PRO	5.0
1	A	69	PRO	5.0
1	A	70	GLU	4.8
1	C	70	GLU	4.8
1	D	70	GLU	4.8
1	B	107	ASP	4.8
1	B	69	PRO	4.6
1	D	69	PRO	4.3
1	C	108	PHE	3.7
1	A	108	PHE	3.5
1	A	107	ASP	3.4
1	A	68	ASP	3.1
1	B	70	GLU	3.1
1	C	239	TRP	3.1
1	D	107	ASP	3.1
1	A	71	GLU	3.0
1	D	66	LEU	2.9
1	A	66	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	66	LEU	2.5
1	B	68	ASP	2.5
1	C	71	GLU	2.4
1	C	24	VAL	2.2
1	D	71	GLU	2.2
1	D	65	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.