



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 23, 2025 – 06:44 pm BST

PDB ID : 5NF0 / pdb\_00005nf0  
Title : Discovery, crystal structures and atomic force microscopy study of thioether ligated D,L-cyclic antimicrobial peptides against multidrug resistant *Pseudomonas aeruginosa*  
Authors : Reymond, J.-L.; Darbre, T.; Stocker, A.; Hong, W.; van Delden, C.; Koehler, T.; Luscher, A.; Visini, R.; Fu, Y.; Di Bonaventura, I.; He, R.  
Deposited on : 2017-03-13  
Resolution : 1.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.46


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2484 (1.30-1.26)
Clashscore	180529	2694 (1.30-1.26)
Ramachandran outliers	177936	2628 (1.30-1.26)
Sidechain outliers	177891	2627 (1.30-1.26)
RSRZ outliers	164620	2481 (1.30-1.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	114	
1	B	114	
1	C	114	
1	D	114	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	E	12	 8% 83% 17%
2	G	12	 8% 83% 17%
2	H	12	 100%
3	F	5	 20% 20% 80%

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4303 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fucose-binding lectin II (PA-IIL).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	A	114	818	504	140	174	0	0	0
1	B	114	823	506	143	174	0	0	0
1	C	114	823	506	143	174	0	0	0
1	D	114	818	504	140	174	0	0	0

- Molecule 2 is a protein called CYD-TRP-TRD-LYS-LYD-LYS-LYD-LYS-TRD-TRP-CYD-GLY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	12	114	80	21	11	2	0	0	1
2	G	12	114	80	21	11	2	0	0	1
2	H	12	114	80	21	11	2	0	0	1

- Molecule 3 is a protein called Fragment of ligand.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	F	1	4	2	1	1	0	0	0

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Ca	0	0
			2	2		

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	2	Total Ca 2 2	0	0
4	C	2	Total Ca 2 2	0	0
4	D	2	Total Ca 2 2	0	0

- Molecule 5 is 3,7-anhydro-2,8-dideoxy-L-glycero-D-gluco-octonic acid (CCD ID: ZDC) (formula: C<sub>8</sub>H<sub>14</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	E	1	Total C O 13 8 5	0	0
5	F	1	Total C O 13 8 5	0	0
5	G	1	Total C O 13 8 5	0	0
5	H	1	Total C O 13 8 5	0	0

- Molecule 6 is ORTHO-XYLENE (CCD ID: OXE) (formula: C<sub>8</sub>H<sub>10</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	E	1	Total C 8 8	0	0
6	G	1	Total C 8 8	0	0
6	H	1	Total C 8 8	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	134	Total O 134 134	0	0
7	B	135	Total O 135 135	0	0
7	C	132	Total O 132 132	0	0
7	D	137	Total O 137 137	0	0
7	E	14	Total O 14 14	0	0
7	F	1	Total O 1 1	0	0
7	G	19	Total O 19 19	0	0
7	H	19	Total O 19 19	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fucose-binding lectin II (PA-IIL)




- Molecule 1: Fucose-binding lectin II (PA-IIL)





- Molecule 1: Fucose-binding lectin II (PA-IIL)




- Molecule 1: Fucose-binding lectin II (PA-IIL)



- Molecule 2: CYD-TRP-TRD-LYS-LYD-LYS-LYD-LYS-TRD-TRP-CYD-GLY



- Molecule 2: CYD-TRP-TRD-LYS-LYD-LYS-LYD-LYS-TRD-TRP-CYD-GLY



- Molecule 2: CYD-TRP-TRD-LYS-LYD-LYS-LYD-LYS-TRD-TRP-CYD-GLY



There are no outlier residues recorded for this chain.

- Molecule 3: Fragment of ligand



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.28Å 99.97Å 106.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.03 – 1.27 47.03 – 1.27	Depositor EDS
% Data completeness (in resolution range)	92.1 (47.03-1.27) 91.6 (47.03-1.27)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.57 (at 1.27Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, $R_{free}$	0.163 , 0.184 0.163 , 0.185	Depositor DCC
$R_{free}$ test set	6023 reflections (4.71%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	11.5	Xtrriage
Anisotropy	0.492	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 34.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4303	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DLY, NH2, OXE, ZDC, DTR, DCY, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.53	0/828	0.69	0/1135
1	B	0.52	0/833	0.71	0/1141
1	C	0.57	0/833	0.72	0/1141
1	D	0.53	0/828	0.69	0/1135
2	E	0.33	0/54	0.44	0/64
2	G	0.44	0/54	0.63	0/64
2	H	0.50	0/54	0.44	0/64
3	F	0.60	0/3	0.55	0/2
All	All	0.53	0/3487	0.69	0/4746

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	818	0	787	2	0
1	B	823	0	796	6	0
1	C	823	0	796	5	0
1	D	818	0	787	7	0
2	E	114	0	106	1	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	G	114	0	106	2	0
2	H	114	0	108	0	0
3	F	4	0	3	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
4	C	2	0	0	0	0
4	D	2	0	0	0	0
5	E	13	0	10	0	0
5	F	13	0	10	0	0
5	G	13	0	10	0	0
5	H	13	0	10	0	0
6	E	8	0	7	0	0
6	G	8	0	8	0	0
6	H	8	0	8	0	0
7	A	134	0	0	1	1
7	B	135	0	0	3	0
7	C	132	0	0	2	1
7	D	137	0	0	2	0
7	E	14	0	0	0	0
7	F	1	0	0	0	0
7	G	19	0	0	0	0
7	H	19	0	0	0	0
All	All	4303	0	3552	20	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:81:VAL:HG21	1:D:81:VAL:HG21	1.49	0.94
1:D:13:ARG:NH1	7:D:401:HOH:O	1.99	0.94
1:A:81:VAL:HG21	1:C:81:VAL:HG21	1.51	0.93
1:A:33:ASN:OD1	7:A:401:HOH:O	2.02	0.77
1:B:13:ARG:NH1	1:D:46:ASN:OD1	2.20	0.75
7:B:454:HOH:O	2:E:7:DLY:HD2	2.06	0.55
1:B:90:ALA:HB3	1:B:109:ILE:HB	1.96	0.48
2:G:117:LYS:HD3	2:G:122:DTR:HZ3	1.95	0.47
1:D:46:ASN:ND2	7:D:402:HOH:O	2.47	0.47
1:C:13:ARG:NH1	7:C:402:HOH:O	2.32	0.47
1:B:11:ASN:CB	7:B:482:HOH:O	2.63	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:90:ALA:HB3	1:C:109:ILE:HB	1.98	0.46
1:B:11:ASN:HB2	7:B:482:HOH:O	2.16	0.45
1:C:66:GLN:NE2	7:C:401:HOH:O	2.17	0.44
2:G:117:LYS:HD3	2:G:122:DTR:CZ3	2.47	0.44
1:C:15:GLY:O	1:C:109:ILE:HA	2.19	0.43
1:D:15:GLY:O	1:D:109:ILE:HA	2.20	0.42
1:D:90:ALA:HB3	1:D:109:ILE:HB	2.02	0.42
1:B:26:GLN:O	1:B:41:SER:HA	2.20	0.41
1:D:103:ASN:OD1	1:D:103:ASN:C	2.63	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:491:HOH:O	7:C:428:HOH:O[4_555]	2.06	0.14

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	112/114 (98%)	109 (97%)	3 (3%)	0	100	100
1	B	112/114 (98%)	109 (97%)	3 (3%)	0	100	100
1	C	112/114 (98%)	109 (97%)	3 (3%)	0	100	100
1	D	112/114 (98%)	110 (98%)	2 (2%)	0	100	100
2	E	5/12 (42%)	5 (100%)	0	0	100	100
2	G	5/12 (42%)	5 (100%)	0	0	100	100
2	H	5/12 (42%)	4 (80%)	1 (20%)	0	100	100
All	All	463/492 (94%)	451 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	91/93 (98%)	90 (99%)	1 (1%)	70	39
1	B	92/93 (99%)	92 (100%)	0	100	100
1	C	92/93 (99%)	91 (99%)	1 (1%)	70	39
1	D	91/93 (98%)	91 (100%)	0	100	100
2	E	5/5 (100%)	4 (80%)	1 (20%)	1	0
2	G	5/5 (100%)	5 (100%)	0	100	100
2	H	5/5 (100%)	5 (100%)	0	100	100
All	All	381/387 (98%)	378 (99%)	3 (1%)	79	54

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	ASN
1	C	33	ASN
2	E	6	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	ASN
1	A	47	ASN
1	B	21	ASN
1	B	47	ASN
1	B	110	ASN
1	C	110	ASN
1	D	21	ASN
1	D	47	ASN
1	D	66	GLN
1	D	110	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

18 non-standard protein/DNA/RNA residues are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 8 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	ZDC	H	201	2,4	13,13,14	1.13	0	17,18,20	1.06	1 (5%)
6	OXE	E	102	2	8,8,8	4.25	5 (62%)	10,10,10	0.64	0
6	OXE	H	202	2	8,8,8	3.84	6 (75%)	10,10,10	1.10	1 (10%)
5	ZDC	E	101	2,4	13,13,14	1.14	0	17,18,20	1.41	3 (17%)
5	ZDC	F	101	4	13,13,14	1.09	1 (7%)	17,18,20	1.76	5 (29%)
5	ZDC	G	201	2,4	13,13,14	1.17	1 (7%)	17,18,20	1.32	2 (11%)
6	OXE	G	202	2	8,8,8	4.00	5 (62%)	10,10,10	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ZDC	H	201	2,4	-	0/3/23/24	0/1/1/1
6	OXE	E	102	2	-	-	0/1/1/1
6	OXE	H	202	2	-	-	0/1/1/1
5	ZDC	E	101	2,4	-	1/3/23/24	0/1/1/1
5	ZDC	F	101	4	-	1/3/23/24	0/1/1/1
5	ZDC	G	201	2,4	-	0/3/23/24	0/1/1/1
6	OXE	G	202	2	-	-	0/1/1/1

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	E	102	OXE	C5-C6	6.89	1.53	1.38
6	E	102	OXE	C4-C3	6.68	1.53	1.38
6	G	202	OXE	C5-C6	6.52	1.52	1.38
6	H	202	OXE	C4-C3	6.31	1.52	1.38
6	H	202	OXE	C5-C6	6.22	1.52	1.38
6	G	202	OXE	C4-C3	5.82	1.51	1.38
6	E	102	OXE	C2-C1	4.86	1.53	1.40
6	G	202	OXE	C2-C1	4.51	1.52	1.40
6	H	202	OXE	C6-C1	-3.47	1.32	1.39
6	H	202	OXE	C2-C1	3.44	1.49	1.40
6	G	202	OXE	C6-C1	-3.37	1.32	1.39
6	E	102	OXE	C6-C1	-3.29	1.32	1.39
6	G	202	OXE	C3-C2	-3.29	1.32	1.39
6	E	102	OXE	C3-C2	-3.08	1.33	1.39
6	H	202	OXE	C3-C2	-2.79	1.33	1.39
6	H	202	OXE	C5-C4	-2.37	1.31	1.38
5	G	201	ZDC	C2-C1	2.34	1.58	1.52
5	F	101	ZDC	C6-C7	2.11	1.55	1.49

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	101	ZDC	O7A-C7-C6	-3.60	114.93	125.43
5	F	101	ZDC	C3-C4-C5	-3.10	104.70	110.24
5	E	101	ZDC	O7A-C7-C6	-2.93	116.88	125.43
5	G	201	ZDC	O7A-C7-C6	-2.84	117.15	125.43
5	E	101	ZDC	C3-C4-C5	-2.74	105.35	110.24

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	H	202	OXE	C1'-C1-C2	-2.45	115.72	120.74
5	G	201	ZDC	O5-C5-C6	2.17	117.84	110.56
5	F	101	ZDC	O3-C3-C2	-2.10	105.49	110.35
5	F	101	ZDC	O5-C5-C6	2.08	117.53	110.56
5	F	101	ZDC	O4-C4-C5	2.07	114.44	109.30
5	H	201	ZDC	O5-C5-C6	2.06	117.48	110.56
5	E	101	ZDC	O5-C5-C6	2.04	117.39	110.56

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	F	101	ZDC	C5-C6-C7-O7A
5	E	101	ZDC	C5-C6-C7-O7A

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	114/114 (100%)	-0.02	4 (3%) 47 48	9, 12, 22, 32	0
1	B	114/114 (100%)	-0.10	2 (1%) 67 69	9, 12, 18, 27	0
1	C	114/114 (100%)	-0.13	2 (1%) 67 69	9, 12, 20, 29	0
1	D	114/114 (100%)	-0.07	3 (2%) 57 59	9, 12, 21, 34	0
2	E	5/12 (41%)	1.52	1 (20%) 3 3	20, 24, 32, 33	0
2	G	5/12 (41%)	0.72	1 (20%) 3 3	15, 18, 19, 29	0
2	H	5/12 (41%)	0.55	0 100 100	15, 15, 22, 30	0
3	F	1/5 (20%)	4.31	1 (100%) 0 0	31, 31, 31, 31	0
All	All	472/497 (94%)	-0.04	14 (2%) 52 54	9, 12, 22, 34	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	11	ASN	5.4
3	F	1	GLY	4.3
1	A	70	ASN	3.6
2	E	2	TRP	3.3
2	G	115	TRP	3.1
1	D	46	ASN	2.7
1	A	33	ASN	2.6
1	D	70	ASN	2.5
1	C	33	ASN	2.4
1	A	71	GLY	2.2
1	C	13	ARG	2.2
1	B	41	SER	2.1
1	A	13	ARG	2.1
1	D	33	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	DLY	E	5	9/10	0.85	0.15	26,37,41,43	0
2	DCY	E	11	6/7	0.88	0.15	17,17,18,30	0
2	DLY	E	7	9/10	0.89	0.15	19,21,43,50	0
2	DTR	E	9	14/15	0.90	0.11	20,26,32,34	0
2	DLY	H	118	9/10	0.92	0.11	14,17,36,38	0
2	DTR	H	116	14/15	0.93	0.08	11,13,17,19	0
2	DTR	G	122	14/15	0.93	0.09	13,19,22,25	0
2	DCY	E	1	6/7	0.93	0.11	15,18,19,23	0
2	DCY	G	114	6/7	0.94	0.09	12,15,17,19	0
2	DTR	E	3	14/15	0.94	0.09	19,21,23,24	0
2	DTR	H	122	14/15	0.94	0.09	17,22,36,36	0
2	DLY	H	120	9/10	0.94	0.08	16,18,21,23	0
2	DLY	G	118	9/10	0.95	0.11	16,18,38,40	0
2	DLY	G	120	9/10	0.96	0.10	11,13,32,37	0
2	DTR	G	116	14/15	0.96	0.07	13,14,15,18	0
2	DCY	G	124	6/7	0.96	0.10	20,22,26,30	0
2	DCY	H	124	6/7	0.97	0.08	13,15,17,17	0
2	DCY	H	114	6/7	0.98	0.05	11,12,14,15	0

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	OXE	E	102	8/8	0.74	0.14	22,24,27,30	0
6	OXE	G	202	8/8	0.89	0.11	19,21,25,26	0
6	OXE	H	202	8/8	0.94	0.08	14,16,17,17	0
5	ZDC	H	201	13/14	0.97	0.06	9,11,12,12	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	ZDC	E	101	13/14	0.97	0.06	10,12,16,17	0
5	ZDC	F	101	13/14	0.97	0.07	11,12,22,23	0
5	ZDC	G	201	13/14	0.97	0.06	9,11,13,14	0
4	CA	D	202	1/1	0.99	0.03	10,10,10,10	0
4	CA	B	301	1/1	0.99	0.03	11,11,11,11	0
4	CA	C	201	1/1	0.99	0.02	9,9,9,9	0
4	CA	C	202	1/1	0.99	0.02	10,10,10,10	0
4	CA	B	302	1/1	1.00	0.02	10,10,10,10	0
4	CA	A	302	1/1	1.00	0.02	10,10,10,10	0
4	CA	A	301	1/1	1.00	0.02	11,11,11,11	0
4	CA	D	201	1/1	1.00	0.02	9,9,9,9	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.