



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:00 AM UTC

PDB ID : 5NVK / pdb_00005nvk
Title : Crystal structure of the human 4EHP-GIGYF1 complex
Authors : Peter, D.; Valkov, E.
Deposited on : 2017-05-04
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	189	
1	C	189	
1	E	189	
1	G	189	
2	B	75	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	75	 <p>12% 92% 7%</p>
2	F	75	 <p>11% 85% 11%</p>
2	H	75	 <p>% 84% 12%</p>

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 15534 atoms, of which 7701 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Eukaryotic translation initiation factor 4E type 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	172	2805	899	1392	255	251	8	0	0	0
1	C	168	2722	876	1341	247	251	7	0	0	0
1	E	168	2721	876	1340	247	251	7	0	0	0
1	G	172	2807	900	1395	254	250	8	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	46	GLY	-	expression tag	UNP O60573
A	47	PRO	-	expression tag	UNP O60573
A	48	HIS	-	expression tag	UNP O60573
A	49	MET	-	expression tag	UNP O60573
A	50	LEU	-	expression tag	UNP O60573
A	51	GLU	-	expression tag	UNP O60573
C	46	GLY	-	expression tag	UNP O60573
C	47	PRO	-	expression tag	UNP O60573
C	48	HIS	-	expression tag	UNP O60573
C	49	MET	-	expression tag	UNP O60573
C	50	LEU	-	expression tag	UNP O60573
C	51	GLU	-	expression tag	UNP O60573
E	46	GLY	-	expression tag	UNP O60573
E	47	PRO	-	expression tag	UNP O60573
E	48	HIS	-	expression tag	UNP O60573
E	49	MET	-	expression tag	UNP O60573
E	50	LEU	-	expression tag	UNP O60573
E	51	GLU	-	expression tag	UNP O60573
G	46	GLY	-	expression tag	UNP O60573
G	47	PRO	-	expression tag	UNP O60573
G	48	HIS	-	expression tag	UNP O60573

Continued on next page...

Continued from previous page...



Chain	Residue	Modelled	Actual	Comment	Reference
G	49	MET	-	expression tag	UNP O60573
G	50	LEU	-	expression tag	UNP O60573
G	51	GLU	-	expression tag	UNP O60573

- Molecule 2 is a protein called GRB10-interacting GYF protein 1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	B	68	Total 1103	C 351	H 550	N 90	O 110	S 2	0	0	0
2	D	74	Total 1217	C 388	H 609	N 100	O 117	S 3	0	0	0
2	F	67	Total 1085	C 345	H 540	N 89	O 109	S 2	0	0	0
2	H	66	Total 1074	C 342	H 534	N 88	O 108	S 2	0	0	0

There are 16 discrepancies between the modelled and reference sequences:



Chain	Residue	Modelled	Actual	Comment	Reference
B	29	GLY	-	expression tag	UNP O75420
B	30	PRO	-	expression tag	UNP O75420
B	31	HIS	-	expression tag	UNP O75420
B	32	MET	-	expression tag	UNP O75420
D	29	GLY	-	expression tag	UNP O75420
D	30	PRO	-	expression tag	UNP O75420
D	31	HIS	-	expression tag	UNP O75420
D	32	MET	-	expression tag	UNP O75420
F	29	GLY	-	expression tag	UNP O75420
F	30	PRO	-	expression tag	UNP O75420
F	31	HIS	-	expression tag	UNP O75420
F	32	MET	-	expression tag	UNP O75420
H	29	GLY	-	expression tag	UNP O75420
H	30	PRO	-	expression tag	UNP O75420
H	31	HIS	-	expression tag	UNP O75420
H	32	MET	-	expression tag	UNP O75420



- Molecule 2: GRB10-interacting GYF protein 1



- Molecule 2: GRB10-interacting GYF protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 42	Depositor
Cell constants a, b, c, α , β , γ	135.32Å 135.32Å 60.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.84 – 2.90 47.84 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.84-2.90) 99.6 (47.84-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 2.91Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.204 , 0.254 0.207 , 0.258	Depositor DCC
R_{free} test set	1218 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	51.9	Xtrriage
Anisotropy	0.165	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 40.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.031 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	15534	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.14	0/1449	0.31	0/1957
1	C	0.13	0/1418	0.32	0/1919
1	E	0.12	0/1418	0.28	0/1919
1	G	0.13	0/1448	0.31	0/1956
2	B	0.12	0/561	0.30	0/759
2	D	0.11	0/619	0.27	0/835
2	F	0.13	0/553	0.30	0/748
2	H	0.11	0/548	0.30	0/741
All	All	0.13	0/8014	0.30	0/10834

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1413	1392	1391	5	0
1	C	1381	1341	1340	5	0
1	E	1381	1340	1340	3	0
1	G	1412	1395	1394	3	0
2	B	553	550	550	5	0
2	D	608	609	609	4	0
2	F	545	540	539	2	0
2	H	540	534	534	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	7833	7701	7697	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:40:ARG:NE	2:D:45:GLU:OE2	2.29	0.62
2:F:62:ASP:OD1	2:F:63:LYS:N	2.35	0.60
1:C:103:ARG:NH1	2:D:75:GLN:O	2.40	0.54
1:A:103:ARG:NH2	2:B:75:GLN:O	2.41	0.54
1:C:100:HIS:NE2	2:D:55:LYS:O	2.42	0.53
1:G:121:LYS:O	1:G:123:MET:N	2.40	0.52
2:D:62:ASP:OD1	2:D:63:LYS:N	2.42	0.51
1:G:103:ARG:NH2	2:H:75:GLN:O	2.42	0.49
1:E:140:ARG:NH2	2:F:93:SER:O	2.39	0.47
1:E:121:LYS:O	1:E:123:MET:N	2.46	0.46
1:C:63:TRP:HB2	1:C:114:HIS:HB2	2.00	0.44
1:C:92:GLU:OE1	1:C:92:GLU:N	2.47	0.43
1:A:161:MET:HE1	2:B:36:LEU:HG	2.01	0.42
1:A:63:TRP:HB2	1:A:114:HIS:HB2	2.01	0.42
1:C:121:LYS:O	1:C:123:MET:N	2.49	0.42
1:A:92:GLU:OE1	1:A:92:GLU:N	2.50	0.42
2:B:62:ASP:OD1	2:B:63:LYS:N	2.52	0.42
2:H:62:ASP:OD1	2:H:63:LYS:N	2.53	0.41
1:E:63:TRP:CE3	1:E:84:GLN:HB3	2.56	0.41
1:A:146:ARG:NH1	2:B:77:LEU:O	2.54	0.41
1:G:92:GLU:OE1	1:G:92:GLU:N	2.53	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	168/189 (89%)	160 (95%)	8 (5%)	0	100	100
1	C	166/189 (88%)	159 (96%)	7 (4%)	0	100	100
1	E	166/189 (88%)	161 (97%)	5 (3%)	0	100	100
1	G	168/189 (89%)	159 (95%)	9 (5%)	0	100	100
2	B	66/75 (88%)	64 (97%)	2 (3%)	0	100	100
2	D	72/75 (96%)	68 (94%)	4 (6%)	0	100	100
2	F	65/75 (87%)	64 (98%)	1 (2%)	0	100	100
2	H	64/75 (85%)	62 (97%)	2 (3%)	0	100	100
All	All	935/1056 (88%)	897 (96%)	38 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	151/166 (91%)	151 (100%)	0	100	100
1	C	148/166 (89%)	148 (100%)	0	100	100
1	E	148/166 (89%)	148 (100%)	0	100	100
1	G	151/166 (91%)	151 (100%)	0	100	100
2	B	61/67 (91%)	61 (100%)	0	100	100
2	D	67/67 (100%)	67 (100%)	0	100	100
2	F	60/67 (90%)	60 (100%)	0	100	100
2	H	60/67 (90%)	60 (100%)	0	100	100
All	All	846/932 (91%)	846 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such

sidechains are listed below:

Mol	Chain	Res	Type
2	B	54	ASN
1	C	205	ASN
1	E	81	ASN
1	G	205	ASN
2	H	54	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	172/189 (91%)	-0.08	5 (2%) 53 45	28, 37, 76, 154	0
1	C	168/189 (88%)	0.27	11 (6%) 25 20	28, 50, 110, 168	0
1	E	168/189 (88%)	0.08	4 (2%) 59 50	31, 53, 88, 119	0
1	G	172/189 (91%)	-0.01	5 (2%) 53 45	28, 39, 78, 153	0
2	B	68/75 (90%)	0.11	0 100 100	31, 50, 83, 91	0
2	D	74/75 (98%)	0.66	9 (12%) 8 7	34, 69, 116, 211	0
2	F	67/75 (89%)	0.58	8 (11%) 9 7	31, 57, 116, 164	0
2	H	66/75 (88%)	0.12	1 (1%) 72 64	32, 55, 90, 115	0
All	All	955/1056 (90%)	0.15	43 (4%) 38 30	28, 47, 98, 211	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	90	PHE	8.0
2	F	92	LEU	5.8
2	F	90	PHE	5.4
1	C	69	PRO	5.3
2	F	91	SER	5.0
1	G	68	THR	4.7
2	D	92	LEU	4.0
2	D	91	SER	3.6
1	C	70	GLY	3.2
2	D	30	PRO	3.2
1	C	77	SER	3.0
1	C	68	THR	3.0
1	C	73	THR	3.0
1	C	176	GLN	2.9
2	D	103	MET	2.9
1	A	68	THR	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	233	LEU	2.9
1	C	52	ALA	2.8
1	A	110	HIS	2.8
1	E	175	PHE	2.6
1	C	124	TRP	2.6
1	G	110	HIS	2.5
2	F	89	ASN	2.5
2	D	89	ASN	2.4
1	A	69	PRO	2.4
1	E	76	GLN	2.3
2	F	94	VAL	2.3
1	A	109	GLY	2.3
1	A	108	THR	2.3
1	G	109	GLY	2.3
2	F	55	LYS	2.3
1	E	67	ARG	2.3
2	D	85	GLU	2.3
1	G	108	THR	2.3
2	H	85	GLU	2.2
2	F	103	MET	2.2
2	D	93	SER	2.1
2	D	83	THR	2.1
1	C	74	SER	2.0
1	C	75	SER	2.0
1	C	125	GLU	2.0
1	E	124	TRP	2.0
2	F	64	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.