



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 9, 2025 – 01:34 AM JST

PDB ID : 5YX2 / pdb_00005yx2
Title : Crystal structure of DNMT3A-DNMT3L in complex with DNA containing two CpG sites
Authors : Zhang, Z.M.; Song, J.
Deposited on : 2017-12-01
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 2.0rc1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1003 (2.66-2.66)
Clashscore	180529	1063 (2.66-2.66)
Ramachandran outliers	177936	1052 (2.66-2.66)
Sidechain outliers	177891	1052 (2.66-2.66)
RSRZ outliers	164620	1003 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	285	
1	D	285	
2	B	208	
2	C	208	
3	E	25	
3	F	25	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8380 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA (cytosine-5)-methyltransferase 3A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	285	2267	1450	402	402	13	0	0	0
1	D	285	2269	1449	402	405	13	0	0	0


- Molecule 2 is a protein called DNA (cytosine-5)-methyltransferase 3-like.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	188	1398	917	233	245	3	0	0	0
2	C	174	1263	821	211	229	2	0	0	0

- Molecule 3 is a DNA chain called DNA (25-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	E	25	510	245	93	148	24	0	0	0
3	F	25	510	245	93	148	24	0	0	0

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is S-ADENOSYL-L-HOMOCYSTEINE (CCD ID: SAH) (formula: C₁₄H₂₀N₆O₅S) (labeled as "Ligand of Interest" by depositor).


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA (cytosine-5)-methyltransferase 3A




- Molecule 1: DNA (cytosine-5)-methyltransferase 3A



- Molecule 2: DNA (cytosine-5)-methyltransferase 3-like



- Molecule 2: DNA (cytosine-5)-methyltransferase 3-like






- Molecule 3: DNA (25-MER)



- Molecule 3: DNA (25-MER)



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	205.42Å 205.42Å 89.34Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.11 – 2.65 33.11 – 2.65	Depositor EDS
% Data completeness (in resolution range)	98.4 (33.11-2.65) 98.4 (33.11-2.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 2.65Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.191 , 0.222 0.194 , 0.224	Depositor DCC
R_{free} test set	2006 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	65.0	Xtrriage
Anisotropy	0.205	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 49.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.010 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8380	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAH, GOL, PYO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.29	0/2321	0.72	2/3138 (0.1%)
1	D	0.29	0/2323	0.71	0/3141
2	B	0.32	0/1441	0.77	2/1977 (0.1%)
2	C	0.33	0/1299	0.80	1/1780 (0.1%)
3	E	0.20	0/550	0.39	0/846
3	F	0.19	0/550	0.41	0/846
All	All	0.29	0/8484	0.70	5/11728 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	367	CYS	N-CA-C	-5.34	106.27	112.89
2	B	327	ILE	CA-C-N	5.28	124.46	118.97
2	B	327	ILE	C-N-CA	5.28	124.46	118.97
1	A	903	ALA	CA-C-N	-5.18	115.54	120.83
1	A	903	ALA	C-N-CA	-5.18	115.54	120.83

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2267	0	2212	31	0
1	D	2269	0	2217	21	0
2	B	1398	0	1235	15	0
2	C	1263	0	1086	17	0
3	E	510	0	279	9	0
3	F	510	0	279	2	0
4	A	24	0	32	3	0
4	D	6	0	8	1	0
4	E	6	0	8	1	0
5	A	26	0	19	1	0
5	D	26	0	19	0	0
6	A	22	0	0	1	0
6	B	5	0	0	0	0
6	C	2	0	0	0	0
6	D	38	0	0	0	0
6	E	1	0	0	0	0
6	F	7	0	0	0	0
All	All	8380	0	7394	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:733:GLU:OE1	1:A:736:ARG:NH1	2.23	0.71
1:A:866:ARG:HE	4:A:1003:GOL:H2	1.56	0.70
1:D:790:ARG:NH1	1:D:892:SER:O	2.28	0.66
1:D:820:GLU:OE1	1:D:866:ARG:NH2	2.29	0.66
1:A:840:ILE:HD11	1:A:859:LEU:HD12	1.79	0.64
2:C:322:ARG:NH2	2:C:340:GLU:OE1	2.22	0.63
2:C:247:PRO:HA	2:C:287:ASN:HD22	1.63	0.63
1:D:733:GLU:OE1	1:D:736:ARG:NH1	2.31	0.63
1:D:690:VAL:O	1:D:736:ARG:NH2	2.31	0.62
1:A:690:VAL:O	1:A:736:ARG:NH2	2.31	0.62
1:D:882:ARG:NH1	3:F:438:DG:OP2	2.33	0.62
1:D:835:THR:HG21	4:E:501:GOL:H31	1.81	0.61
1:A:882:ARG:NH1	3:E:438:DG:OP2	2.34	0.60
1:D:840:ILE:HD11	1:D:859:LEU:HD12	1.83	0.60
1:A:709:PRO:HB3	4:A:1002:GOL:H31	1.82	0.60
1:A:885:ARG:NH2	1:D:876:ASP:OD1	2.34	0.59
1:A:876:ASP:OD1	1:D:885:ARG:NH2	2.35	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:882:ARG:NH2	1:D:886:GLN:OE1	2.35	0.58
2:C:342:GLU:OE1	2:C:342:GLU:N	2.38	0.56
2:C:256:PRO:HB2	2:C:290:LEU:HD23	1.90	0.54
1:A:790:ARG:NH1	1:A:892:SER:O	2.41	0.54
1:D:792:ARG:NH1	3:E:427:PYO:O4'	2.41	0.54
3:E:432:DT:H2''	3:E:433:DA:C8	2.44	0.53
2:C:339:SER:OG	2:C:340:GLU:N	2.42	0.52
2:C:306:PRO:HB3	2:C:324:TRP:CE2	2.45	0.52
2:B:334:HIS:ND1	2:B:334:HIS:O	2.43	0.51
1:A:878:SER:HA	1:D:860:TRP:CE2	2.45	0.51
3:E:424:DA:H2''	3:E:425:DT:H72	1.93	0.49
1:D:835:THR:HG22	1:D:887:ARG:HA	1.94	0.49
2:C:275:GLY:O	2:C:277:PRO:HD3	2.12	0.49
1:D:875:THR:O	1:D:885:ARG:HD3	2.14	0.48
2:C:327:ILE:HG22	2:C:329:ALA:H	1.78	0.48
1:D:710:CYS:N	3:E:427:PYO:HC5	2.28	0.48
2:B:191:ARG:HD2	2:B:237:PRO:O	2.14	0.48
2:C:224:VAL:HB	2:C:266:LEU:HD11	1.96	0.48
4:A:1002:GOL:H11	5:A:1005:SAH:H8	1.97	0.47
1:A:715:ILE:HG12	1:A:760:ALA:HB1	1.98	0.46
2:C:195:LEU:HD22	2:C:224:VAL:HG11	1.97	0.46
1:A:875:THR:O	1:A:885:ARG:HD3	2.16	0.46
3:E:445:DT:H2''	3:E:446:DG:C8	2.51	0.46
1:D:745:GLU:HB2	2:C:274:PRO:HG3	1.97	0.46
1:A:639:LEU:HD12	1:A:734:PHE:CD2	2.51	0.46
2:B:182:VAL:O	2:B:187:ARG:NH1	2.46	0.45
2:C:190:VAL:HB	2:C:375:PHE:CG	2.51	0.45
1:A:808:THR:HG23	1:A:810:ASN:H	1.82	0.45
2:B:229:ARG:O	2:B:233:GLU:HG3	2.17	0.45
2:B:264:HIS:O	2:B:268:GLN:HG2	2.17	0.44
1:D:831:ARG:NH2	3:E:425:DT:H5''	2.33	0.44
2:B:182:VAL:HG22	2:B:183:PRO:HD2	1.98	0.44
1:A:906:LYS:HB2	1:A:911:CYS:SG	2.57	0.44
2:B:308:THR:O	2:B:310:PRO:HD3	2.16	0.44
1:A:743:PRO:HB2	1:A:747:ASP:HB3	1.99	0.44
2:C:264:HIS:O	2:C:268:GLN:HG2	2.17	0.44
3:E:425:DT:H2''	3:E:426:DG:C8	2.53	0.44
1:A:826:LYS:HD3	1:A:851:PHE:CZ	2.53	0.43
2:B:268:GLN:HE22	2:B:271:ARG:HH21	1.66	0.43
2:B:353:SER:O	2:B:353:SER:OG	2.36	0.43
1:D:798:LEU:O	1:D:801:MET:HG3	2.17	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:199:ILE:HB	2:C:203:LEU:HD13	2.00	0.43
1:A:777:PRO:HB3	1:A:795:TRP:CE2	2.53	0.43
1:A:826:LYS:HD3	1:A:851:PHE:CE2	2.54	0.43
2:B:240:LEU:HD23	2:B:281:PHE:HB2	2.00	0.43
1:A:631:ARG:NH1	1:A:906:LYS:O	2.52	0.42
2:B:348:GLN:O	2:B:352:SER:OG	2.34	0.42
1:A:865:GLU:OE1	1:A:875:THR:OG1	2.33	0.42
1:A:674:MET:HB3	6:A:1117:HOH:O	2.19	0.42
1:A:752:PHE:HB3	1:A:798:LEU:HD23	2.01	0.42
2:B:202:GLU:OE1	2:B:202:GLU:N	2.48	0.42
3:E:428:DG:OP2	3:E:428:DG:H8	2.02	0.42
1:A:813:LEU:O	1:A:829:LYS:NZ	2.52	0.42
1:A:764:SER:OG	1:A:767:ARG:NH2	2.53	0.42
1:A:772:PHE:CE2	2:B:265:ARG:HB2	2.55	0.42
3:F:422:DG:H1'	3:F:423:DC:H5'	2.01	0.42
2:C:190:VAL:HB	2:C:375:PHE:CD2	2.55	0.41
2:B:194:SER:OG	2:B:197:GLU:O	2.37	0.41
1:D:865:GLU:OE1	1:D:875:THR:OG1	2.38	0.41
1:A:835:THR:HG22	1:A:887:ARG:HA	2.02	0.41
1:D:752:PHE:HB3	1:D:798:LEU:HD23	2.03	0.41
1:A:724:TYR:CZ	2:B:258:TRP:HD1	2.38	0.41
1:D:751:PHE:H	4:D:1001:GOL:H12	1.84	0.41
1:A:808:THR:HG22	1:A:811:ASP:CG	2.45	0.41
1:A:649:VAL:HG21	1:A:898:ILE:HB	2.03	0.41
1:A:808:THR:HG22	1:A:811:ASP:OD2	2.21	0.41
2:C:290:LEU:HD22	2:C:294:ASP:HB3	2.02	0.40
2:C:322:ARG:HH12	2:C:340:GLU:CD	2.23	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/285 (99%)	276 (98%)	7 (2%)	0	100	100
1	D	283/285 (99%)	274 (97%)	9 (3%)	0	100	100
2	B	180/208 (86%)	171 (95%)	9 (5%)	0	100	100
2	C	164/208 (79%)	148 (90%)	15 (9%)	1 (1%)	22	35
All	All	910/986 (92%)	869 (96%)	40 (4%)	1 (0%)	48	67

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	278	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/250 (94%)	232 (99%)	3 (1%)	65	80
1	D	238/250 (95%)	235 (99%)	3 (1%)	65	80
2	B	129/190 (68%)	127 (98%)	2 (2%)	58	76
2	C	114/190 (60%)	111 (97%)	3 (3%)	41	63
All	All	716/880 (81%)	705 (98%)	11 (2%)	60	77

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	661	ILE
1	A	877	VAL
1	A	885	ARG
2	B	182	VAL
2	B	222	VAL
1	D	696	GLN
1	D	877	VAL
1	D	882	ARG
2	C	188	GLN
2	C	223	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	C	321	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	287	ASN
1	D	692	GLN
2	C	268	GLN
2	C	287	ASN
2	C	334	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PYO	F	427	3	16,20,21	1.19	2 (12%)	22,28,31	1.22	2 (9%)
3	PYO	E	427	3	16,20,21	1.20	2 (12%)	22,28,31	1.24	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PYO	F	427	3	-	4/7/25/26	0/2/2/2
3	PYO	E	427	3	-	4/7/25/26	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	427	PYO	C2-N1	-3.33	1.32	1.40
3	F	427	PYO	C2-N1	-3.29	1.32	1.40
3	F	427	PYO	C6-C5	2.50	1.41	1.35
3	E	427	PYO	C6-C5	2.32	1.40	1.35

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	427	PYO	C5-C4-N3	-4.00	119.39	124.29
3	E	427	PYO	C5-C4-N3	-3.76	119.68	124.29
3	E	427	PYO	C5-C6-N1	-2.87	118.28	121.19
3	F	427	PYO	C5-C6-N1	-2.45	118.71	121.19

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	427	PYO	C2'-C1'-N1-C6
3	F	427	PYO	C2'-C1'-N1-C6
3	E	427	PYO	C2'-C1'-N1-C2
3	F	427	PYO	C2'-C1'-N1-C2
3	F	427	PYO	O4'-C1'-N1-C6
3	F	427	PYO	O4'-C1'-N1-C2
3	E	427	PYO	O4'-C1'-N1-C6
3	E	427	PYO	O4'-C1'-N1-C2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	427	PYO	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	E	501	-	5,5,5	0.36	0	5,5,5	0.37	0
5	SAH	D	1002	-	24,28,28	1.21	3 (12%)	25,40,40	1.65	4 (16%)
4	GOL	A	1002	-	5,5,5	0.34	0	5,5,5	0.29	0
4	GOL	A	1004	-	5,5,5	0.34	0	5,5,5	0.31	0
4	GOL	A	1003	-	5,5,5	0.36	0	5,5,5	0.26	0
4	GOL	D	1001	-	5,5,5	0.34	0	5,5,5	0.34	0
5	SAH	A	1005	-	24,28,28	1.21	3 (12%)	25,40,40	1.65	4 (16%)
4	GOL	A	1001	-	5,5,5	0.35	0	5,5,5	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	E	501	-	-	0/4/4/4	-
5	SAH	D	1002	-	-	5/11/31/31	0/3/3/3
4	GOL	A	1002	-	-	4/4/4/4	-
4	GOL	A	1004	-	-	3/4/4/4	-
4	GOL	A	1003	-	-	1/4/4/4	-
4	GOL	D	1001	-	-	2/4/4/4	-
5	SAH	A	1005	-	-	3/11/31/31	0/3/3/3
4	GOL	A	1001	-	-	2/4/4/4	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1005	SAH	C2-N3	4.03	1.38	1.32
5	D	1002	SAH	C2-N3	3.95	1.38	1.32
5	D	1002	SAH	C2-N1	2.45	1.38	1.33
5	A	1005	SAH	C2-N1	2.39	1.38	1.33
5	D	1002	SAH	OXT-C	-2.15	1.23	1.30
5	A	1005	SAH	OXT-C	-2.11	1.23	1.30

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	1002	SAH	N3-C2-N1	-5.64	119.87	128.68
5	A	1005	SAH	N3-C2-N1	-5.53	120.03	128.68
5	A	1005	SAH	C5'-SD-CG	-3.10	92.96	102.27
5	D	1002	SAH	OXT-C-O	-2.95	117.39	124.09
5	A	1005	SAH	OXT-C-O	-2.83	117.67	124.09
5	D	1002	SAH	C5'-SD-CG	-2.80	93.88	102.27
5	A	1005	SAH	OXT-C-CA	2.30	121.23	113.38
5	D	1002	SAH	OXT-C-CA	2.27	121.10	113.38

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1002	GOL	O1-C1-C2-C3
4	A	1002	GOL	C1-C2-C3-O3
4	A	1004	GOL	O1-C1-C2-C3
4	D	1001	GOL	C1-C2-C3-O3
5	A	1005	SAH	N-CA-CB-CG
5	D	1002	SAH	N-CA-CB-CG
4	A	1004	GOL	O1-C1-C2-O2
4	A	1001	GOL	C1-C2-C3-O3
4	A	1001	GOL	O2-C2-C3-O3
4	A	1002	GOL	O2-C2-C3-O3
4	D	1001	GOL	O2-C2-C3-O3
5	A	1005	SAH	C-CA-CB-CG
4	A	1002	GOL	O1-C1-C2-O2
5	D	1002	SAH	OXT-C-CA-CB
5	D	1002	SAH	CB-CG-SD-C5'
5	D	1002	SAH	O-C-CA-CB
5	D	1002	SAH	C-CA-CB-CG
5	A	1005	SAH	CB-CG-SD-C5'
4	A	1003	GOL	C1-C2-C3-O3
4	A	1004	GOL	C1-C2-C3-O3

There are no ring outliers.

5 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	501	GOL	1	0
4	A	1002	GOL	2	0


Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1003	GOL	1	0
4	D	1001	GOL	1	0
5	A	1005	SAH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	285/285 (100%)	-0.45	3 (1%) 77 75	41, 62, 95, 142	0
1	D	285/285 (100%)	-0.52	2 (0%) 84 82	40, 56, 88, 118	0
2	B	188/208 (90%)	0.26	7 (3%) 45 43	64, 100, 144, 156	0
2	C	174/208 (83%)	0.47	12 (6%) 24 22	72, 122, 161, 172	0
3	E	24/25 (96%)	-0.55	0 100 100	69, 87, 108, 114	0
3	F	24/25 (96%)	-0.45	0 100 100	66, 91, 103, 108	0
All	All	980/1036 (94%)	-0.17	24 (2%) 59 58	40, 75, 145, 172	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	215	PRO	4.3
2	B	367	CYS	4.0
2	B	251	THR	3.6
2	B	192	VAL	3.3
2	C	253	ASP	2.9
1	A	805	LEU	2.8
2	C	364	VAL	2.7
2	C	320	ALA	2.7
2	C	378	PHE	2.6
1	A	628	ALA	2.6
2	C	361	THR	2.4
2	B	222	VAL	2.3
1	D	851	PHE	2.3
2	B	179	PHE	2.3
2	B	336	ALA	2.3
2	B	312	VAL	2.2
2	C	363	LEU	2.2
2	C	372	ARG	2.2
1	A	844	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	628	ALA	2.2
2	C	313	HIS	2.1
2	C	367	CYS	2.0
2	C	250	HIS	2.0
2	C	336	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PYO	E	427	19/20	0.95	0.08	59,61,68,68	0
3	PYO	F	427	19/20	0.96	0.07	51,57,62,63	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	A	1003	6/6	0.73	0.14	76,82,87,89	0
4	GOL	A	1004	6/6	0.78	0.14	69,79,87,90	0
4	GOL	A	1001	6/6	0.81	0.12	89,95,100,102	0
4	GOL	E	501	6/6	0.84	0.26	65,83,88,91	0
4	GOL	D	1001	6/6	0.87	0.10	83,86,89,90	0
4	GOL	A	1002	6/6	0.89	0.13	65,70,73,82	0
5	SAH	A	1005	26/26	0.94	0.08	39,56,68,71	0
5	SAH	D	1002	26/26	0.96	0.07	45,53,63,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different

orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.