



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 10, 2025 – 10:15 PM JST

PDB ID : 6JU4 / pdb_00006ju4
Title : Aspergillus oryzae pro-tyrosinase F513Y mutant
Authors : Fujieda, N.; Umakoshi, K.; Nishikawa, Y.; Kurisu, G.; Itoh, S.
Deposited on : 2019-04-13
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1089 (1.36-1.36)
Clashscore	180529	1157 (1.36-1.36)
Ramachandran outliers	177936	1146 (1.36-1.36)
Sidechain outliers	177891	1146 (1.36-1.36)
RSRZ outliers	164620	1088 (1.36-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	621	3% 80% 7% 13%
1	B	621	3% 84% 6% 10%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9596 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	541	Total	C	N	O	S	0	4	0
			4298	2767	730	788	13			
1	B	560	Total	C	N	O	S	0	4	0
			4446	2860	761	812	13			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP A0A1S9DK56
A	-2	PRO	-	expression tag	UNP A0A1S9DK56
A	-1	GLY	-	expression tag	UNP A0A1S9DK56
A	0	GLY	-	expression tag	UNP A0A1S9DK56
A	1	SER	-	expression tag	UNP A0A1S9DK56
A	513	TYR	PHE	engineered mutation	UNP A0A1S9DK56
A	513	TYR	PHE	microheterogeneity	UNP A0A1S9DK56
A	513	DAH	PHE	microheterogeneity	UNP A0A1S9DK56
B	-3	GLY	-	expression tag	UNP A0A1S9DK56
B	-2	PRO	-	expression tag	UNP A0A1S9DK56
B	-1	GLY	-	expression tag	UNP A0A1S9DK56
B	0	GLY	-	expression tag	UNP A0A1S9DK56
B	1	SER	-	expression tag	UNP A0A1S9DK56
B	513	TYR	PHE	engineered mutation	UNP A0A1S9DK56
B	513	TYR	PHE	microheterogeneity	UNP A0A1S9DK56
B	513	DAH	PHE	microheterogeneity	UNP A0A1S9DK56

- Molecule 2 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cu	0	2
			4	4		
2	B	2	Total	Cu	0	2
			4	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	401	Total 409	O 409	0	10
3	B	429	Total 435	O 435	0	7


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tyrosinase



• Molecule 1: Tyrosinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.22Å 118.14Å 79.35Å 90.00° 91.11° 90.00°	Depositor
Resolution (Å)	30.00 – 1.35 30.00 – 1.35	Depositor EDS
% Data completeness (in resolution range)	99.9 (30.00-1.35) 99.7 (30.00-1.35)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 1.35Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.155 , 0.201 0.142 , 0.186	Depositor DCC
R_{free} test set	10896 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	11.8	Xtriage
Anisotropy	0.081	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 53.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.023 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9596	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CU, DAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/4427	1.03	4/6040 (0.1%)
1	B	0.42	0/4581	1.04	7/6245 (0.1%)
All	All	0.42	0/9008	1.03	11/12285 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	372	HIS	CA-CB-CG	-7.68	106.12	113.80
1	B	28	GLU	CA-C-N	-7.21	110.75	121.99
1	B	28	GLU	C-N-CA	-7.21	110.75	121.99
1	A	303	ASP	CA-CB-CG	6.33	118.92	112.60
1	A	372	HIS	CA-CB-CG	-6.24	107.56	113.80
1	B	281	ASP	CA-CB-CG	5.97	118.57	112.60
1	A	359	VAL	CB-CA-C	-5.92	108.08	113.70
1	B	28	GLU	CA-C-O	5.72	126.43	119.11
1	B	224	ASP	CA-CB-CG	5.25	117.84	112.60
1	A	600	ARG	CA-C-O	-5.15	115.14	120.70
1	B	405	ARG	CD-NE-CZ	5.12	131.57	124.40

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	513[B]	DAH	Mainchain
1	B	513[B]	DAH	Mainchain
1	B	513[A]	TYR	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4298	0	3958	26	0
1	B	4446	0	4095	29	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
3	A	409	0	0	5	0
3	B	435	0	0	7	0
All	All	9596	0	8053	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:359:VAL:HG22	1:B:513[A]:TYR:CE2	2.00	0.94
1:B:359:VAL:HG22	1:B:513[A]:TYR:CZ	2.08	0.88
1:B:359:VAL:CG2	1:B:513[A]:TYR:CE2	2.70	0.75
1:A:240:GLU:O	1:A:241:ASN:HB2	1.92	0.69
1:A:453:GLY:O	1:A:454:GLN:C	2.34	0.67
1:B:359:VAL:CG2	1:B:513[A]:TYR:CZ	2.79	0.66
1:B:284:PHE:CE1	1:B:288:THR:HG21	2.31	0.65
1:A:267:TYR:OH	1:A:543:LEU:HB3	1.99	0.62
1:A:24:ASP:O	1:A:28:GLU:HG3	2.00	0.62
1:B:284:PHE:O	1:B:288:THR:HG23	2.02	0.59
1:A:207:ARG:HA	3:A:1128:HOH:O	2.02	0.58
1:B:240:GLU:O	1:B:241:ASN:HB2	2.04	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:126:VAL:O	1:B:131:LYS:HE3	2.05	0.56
1:A:264:HIS:NE2	1:B:347:ASP:OD2	2.34	0.55
1:B:347:ASP:HB2	3:B:814:HOH:O	2.07	0.54
1:A:359:VAL:HG22	1:A:513[A]:TYR:CE2	2.43	0.53
1:B:542:ARG:HD3	1:B:542:ARG:C	2.34	0.52
1:A:317:ALA:N	1:A:318:PRO:CD	2.73	0.52
1:B:131:LYS:CE	3:B:851:HOH:O	2.59	0.51
1:A:233:THR:HA	1:A:355:HIS:CE1	2.46	0.50
1:A:547:TYR:CZ	1:A:551:GLN:HG3	2.47	0.50
1:A:1:SER:HA	1:A:2:PRO:C	2.35	0.50
1:A:302:TYR:CE2	3:A:1032:HOH:O	2.53	0.50
1:A:515:GLY:HA3	3:A:991[B]:HOH:O	2.12	0.50
1:A:359:VAL:HG22	1:A:513[A]:TYR:CZ	2.47	0.48
1:B:346:HIS:O	1:B:349:LYS:NZ	2.46	0.48
1:A:74:TRP:CE3	1:A:75:ASP:HB2	2.48	0.48
1:B:586:THR:OG1	1:B:589:ASP:OD2	2.28	0.47
1:A:302:TYR:CE2	1:A:473:ARG:NH1	2.83	0.47
1:B:349:LYS:HD3	3:B:1108:HOH:O	2.15	0.46
1:B:88:PRO:C	1:B:89:THR:CG2	2.89	0.45
1:A:333:ASN:HB2	1:A:511:PHE:CD2	2.51	0.45
1:B:233:THR:HA	1:B:355:HIS:CE1	2.51	0.45
1:B:349:LYS:HE3	3:B:1108:HOH:O	2.17	0.45
1:B:333:ASN:HB2	1:B:511:PHE:CD2	2.52	0.44
1:B:88:PRO:C	1:B:89:THR:HG23	2.43	0.44
1:A:43:VAL:HG21	1:A:175:VAL:HG11	2.00	0.44
1:B:363:ALA:HA	1:B:368:PHE:CG	2.54	0.43
1:A:339:MET:HE2	3:A:830:HOH:O	2.18	0.43
1:B:267:TYR:CD1	1:B:267:TYR:C	2.96	0.43
1:A:304:ASP:OD1	1:A:304:ASP:C	2.60	0.42
1:A:363:ALA:HA	1:A:368:PHE:CG	2.54	0.42
1:B:550:LYS:HD3	1:B:550:LYS:HA	1.85	0.42
1:A:267:TYR:CD1	1:A:267:TYR:C	2.96	0.42
1:A:302:TYR:CD2	3:A:1032:HOH:O	2.73	0.42
1:B:457:LYS:HD3	1:B:588:TYR:CD1	2.55	0.42
1:B:131:LYS:O	1:B:135:GLU:HG3	2.20	0.41
1:B:131:LYS:HE2	3:B:851:HOH:O	2.21	0.41
1:B:118:MET:HE1	1:B:141:TRP:CD2	2.56	0.41
1:A:178:LEU:HD23	1:A:186:GLN:HG2	2.03	0.41
1:B:7:GLY:HA2	3:B:945:HOH:O	2.21	0.41
1:A:333:ASN:HB2	1:A:511:PHE:CG	2.56	0.40
1:B:54:LEU:HD23	3:B:873:HOH:O	2.20	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:385:VAL:HG23	1:A:386:ASN:ND2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	527/621 (85%)	516 (98%)	11 (2%)	0	100	100
1	B	548/621 (88%)	542 (99%)	6 (1%)	0	100	100
All	All	1075/1242 (87%)	1058 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	433/537 (81%)	429 (99%)	4 (1%)	75	51
1	B	447/537 (83%)	445 (100%)	2 (0%)	89	78
All	All	880/1074 (82%)	874 (99%)	6 (1%)	81	62

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	328	HIS
1	A	406	PRO
1	A	439	LYS
1	A	504	GLN
1	B	328	HIS
1	B	504	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	371	HIS
1	A	546	HIS
1	B	50	GLN
1	B	197	GLN
1	B	280	GLN
1	B	333	ASN
1	B	371	HIS
1	B	512	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	DAH	B	513[B]	2,1	12,13,14	0.35	0	14,17,19	0.92	0
1	DAH	A	513[B]	2,1	12,13,14	0.35	0	14,17,19	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	DAH	B	513[B]	2,1	-	0/5/6/8	0/1/1/1
1	DAH	A	513[B]	2,1	-	1/5/6/8	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	513[B]	DAH	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	540/621 (86%)	-0.15	16 (2%)	52	63	9, 16, 37, 71	3 (0%)
1	B	559/621 (90%)	-0.14	16 (2%)	54	64	8, 15, 35, 52	3 (0%)
All	All	1099/1242 (88%)	-0.15	32 (2%)	54	64	8, 16, 37, 71	6 (0%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	526	ALA	5.4
1	A	32	ALA	4.7
1	B	84	SER	3.9
1	A	183	PRO	3.8
1	B	317	ALA	3.6
1	B	30	THR	3.4
1	A	184	GLY	3.4
1	A	317	ALA	3.4
1	B	272	GLY	2.9
1	B	222	SER	2.9
1	B	223	PHE	2.9
1	B	212	THR	2.9
1	A	410	PHE	2.8
1	A	33	GLY	2.8
1	B	184	GLY	2.7
1	B	-2	PRO	2.7
1	A	272	GLY	2.6
1	A	212	THR	2.6
1	A	31	SER	2.6
1	A	89	THR	2.5
1	B	-1	GLY	2.5
1	B	589	ASP	2.5
1	B	153	ARG	2.4
1	A	153	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	514	SER	2.3
1	A	224	ASP	2.3
1	B	89	THR	2.2
1	B	213	SER	2.2
1	A	304	ASP	2.2
1	B	0	GLY	2.1
1	A	187	LEU	2.1
1	A	181	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	DAH	A	513[B]	13/14	-	-	12,14,21,23	13
1	DAH	B	513[B]	13/14	-	-	13,15,19,25	13

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CU	A	701[A]	1/1	1.00	0.01	15,15,15,15	1
2	CU	A	701[B]	1/1	1.00	0.01	20,20,20,20	1
2	CU	A	702[A]	1/1	1.00	0.03	12,12,12,12	1
2	CU	A	702[B]	1/1	1.00	0.03	11,11,11,11	1
2	CU	B	701[A]	1/1	1.00	0.02	17,17,17,17	1
2	CU	B	701[B]	1/1	1.00	0.02	15,15,15,15	1
2	CU	B	702[A]	1/1	1.00	0.01	13,13,13,13	1
2	CU	B	702[B]	1/1	1.00	0.01	12,12,12,12	1

6.5 Other polymers [i](#)

There are no such residues in this entry.