



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:59 AM UTC

PDB ID : 6KVL / pdb_00006kvl
Title : Crystal structure of UDP-RebB-SrUGT76G1
Authors : Li, J.X.; Liu, Z.F.; Wang, Y.; Zhang, P.
Deposited on : 2019-09-04
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	461	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4057 atoms, of which 70 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called UDP-glycosyltransferase 76G1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	448	3578	2295	601	671	11	0	0	0

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q6VAB4
A	460	THR	-	expression tag	UNP Q6VAB4
A	461	SER	-	expression tag	UNP Q6VAB4

- Molecule 2 is URIDINE-5'-DIPHOSPHATE (CCD ID: UDP) (formula: $C_9H_{14}N_2O_{12}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	A	1	36	9	11	2	12	0	0

- Molecule 3 is (8alpha,9beta,10alpha,13alpha)-13-{[beta-D-glucopyranosyl-(1->2)-[beta-D-glucopyranosyl-(1->3)]-beta-D-glucopyranosyl]oxy}kaur-16-en-18-oic acid (CCD ID: AUO) (formula: C₃₈H₆₀O₁₈) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			115	38	59	18		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	328	Total	O	0	0
			328	328		

- Molecule 1: UDP-glycosyltransferase 76G1



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	97.46Å 97.46Å 89.90Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.67 – 2.00 39.67 – 2.00	Depositor EDS
% Data completeness (in resolution range)	82.3 (39.67-2.00) 64.9 (39.67-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.10_2155	Depositor
R, R_{free}	0.168 , 0.204 0.170 , 0.199	Depositor DCC
R_{free} test set	1146 reflections (3.39%)	wwPDB-VP
Wilson B-factor (Å ²)	20.2	Xtriage
Anisotropy	0.117	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.029 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4057	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AUO, UDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	4/3667 (0.1%)	0.74	12/4976 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	330	LEU	N-CA	-6.38	1.38	1.46
1	A	323	GLU	C-O	-5.48	1.19	1.24
1	A	321	TRP	C-O	-5.17	1.17	1.24
1	A	180	ALA	C-O	-5.04	1.17	1.23

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	264	THR	N-CA-C	-17.65	87.81	114.64
1	A	263	ARG	CB-CA-C	-10.02	96.66	112.09
1	A	322	VAL	N-CA-C	9.79	129.69	109.34
1	A	181	SER	N-CA-C	9.77	127.40	111.37
1	A	264	THR	N-CA-CB	7.83	123.50	111.06
1	A	322	VAL	N-CA-CB	-7.08	99.55	111.23
1	A	321	TRP	N-CA-C	-6.85	96.21	110.80
1	A	332	GLU	CA-C-N	-5.58	111.67	122.06
1	A	332	GLU	C-N-CA	-5.58	111.67	122.06

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	181	SER	CB-CA-C	-5.56	101.61	110.84
1	A	263	ARG	N-CA-C	-5.50	103.64	111.24
1	A	164	GLU	N-CA-C	5.15	117.56	111.33

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	329	PHE	Peptide
1	A	330	LEU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3578	0	3527	75	0
2	A	25	11	11	0	0
3	A	56	59	0	1	0
4	A	328	0	0	13	0
All	All	3987	70	3538	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:298:ARG:NH2	1:A:405:ARG:HD2	1.62	1.13
1:A:298:ARG:HH22	1:A:405:ARG:HD2	0.93	1.04
1:A:298:ARG:HH22	1:A:405:ARG:CD	1.72	1.01
1:A:20:PRO:HB3	1:A:30:ILE:HG21	1.49	0.94
1:A:298:ARG:NH2	1:A:405:ARG:CD	2.28	0.93
1:A:317:LYS:HA	1:A:317:LYS:HE2	1.51	0.93
1:A:224:LYS:HG3	1:A:231:LEU:HD11	1.52	0.92
1:A:263:ARG:NH2	4:A:602:HOH:O	2.00	0.92

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:96:GLU:HG3	4:A:609:HOH:O	1.72	0.88
1:A:262:ASP:O	4:A:601:HOH:O	1.94	0.85
1:A:298:ARG:NH2	1:A:405:ARG:NE	2.32	0.78
1:A:74:ASP:OD2	1:A:76:ARG:HG3	1.89	0.72
1:A:333:ARG:CZ	1:A:333:ARG:HA	2.21	0.71
1:A:312:ARG:HH21	1:A:313:PRO:HD2	1.58	0.69
1:A:86:LEU:O	1:A:89:MET:HG3	1.93	0.68
1:A:146:MET:CE	1:A:212:THR:HG21	2.24	0.66
1:A:236:ARG:NH1	1:A:237:GLU:OE2	2.30	0.65
1:A:317:LYS:HA	1:A:317:LYS:CE	2.28	0.61
1:A:73:GLN:HB3	1:A:77:ILE:CD1	2.31	0.60
1:A:297:ALA:O	1:A:301:VAL:HG23	2.02	0.60
1:A:298:ARG:NH2	1:A:405:ARG:HE	2.00	0.60
1:A:249:LYS:HD2	1:A:368:CYS:HB3	1.84	0.59
1:A:286:SER:O	4:A:603:HOH:O	2.17	0.59
1:A:282:PHE:O	1:A:286:SER:OG	2.18	0.59
1:A:109:LEU:HD23	1:A:109:LEU:C	2.29	0.58
1:A:80:LEU:HD11	1:A:93:ILE:HD12	1.84	0.58
1:A:409:ALA:O	1:A:413:ARG:HG3	2.06	0.56
1:A:91:ILE:HB	1:A:92:PRO:HD3	1.87	0.56
1:A:317:LYS:O	4:A:605:HOH:O	2.18	0.55
1:A:148:SER:OG	4:A:604:HOH:O	2.18	0.54
1:A:388:TYR:CE2	1:A:393:LEU:HD21	2.43	0.54
1:A:20:PRO:HD2	1:A:47:PHE:O	2.09	0.52
1:A:75:GLU:HB3	4:A:671:HOH:O	2.10	0.52
1:A:172:ASP:O	1:A:192:LYS:HE3	2.08	0.52
1:A:388:TYR:HE2	1:A:393:LEU:HD21	1.75	0.52
1:A:404:GLU:HG3	1:A:407:GLU:H	1.76	0.51
1:A:330:LEU:HD22	1:A:330:LEU:N	2.26	0.50
1:A:105:GLU:OE2	4:A:606:HOH:O	2.20	0.49
1:A:77:ILE:HD12	1:A:78:SER:N	2.27	0.49
1:A:86:LEU:HD13	1:A:89:MET:SD	2.53	0.49
1:A:96:GLU:CG	4:A:609:HOH:O	2.46	0.48
1:A:399:LEU:HD22	1:A:407:GLU:HG2	1.94	0.48
1:A:240:ALA:HB1	1:A:241:PRO:HD2	1.96	0.48
1:A:301:VAL:CG2	1:A:330:LEU:HB2	2.43	0.48
1:A:40:LYS:HG3	1:A:453:VAL:HG21	1.95	0.47
1:A:48:HIS:O	1:A:66:PHE:HA	2.14	0.47
1:A:301:VAL:CG1	1:A:330:LEU:HB2	2.44	0.47
1:A:306:SER:HA	1:A:333:ARG:HB3	1.97	0.47
1:A:173:LYS:HB3	1:A:173:LYS:HE2	1.51	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:317:LYS:HE2	1:A:317:LYS:CA	2.34	0.47
1:A:393:LEU:HD22	1:A:393:LEU:N	2.30	0.46
1:A:290:GLU:O	1:A:294:LEU:HD23	2.15	0.46
1:A:312:ARG:HD2	1:A:313:PRO:HD2	1.99	0.45
1:A:317:LYS:CE	1:A:317:LYS:CA	2.95	0.45
1:A:263:ARG:HB2	4:A:601:HOH:O	2.16	0.45
1:A:77:ILE:HG22	1:A:80:LEU:HD12	1.99	0.45
1:A:262:ASP:O	1:A:263:ARG:HB2	2.17	0.44
1:A:317:LYS:HA	1:A:318:GLY:HA2	1.55	0.44
1:A:107:GLU:HA	1:A:110:MET:HE2	2.00	0.43
1:A:55:LYS:CE	4:A:630:HOH:O	2.66	0.43
3:A:502:AUO:C16	3:A:502:AUO:C55	2.95	0.43
1:A:367:VAL:HG13	1:A:431:LEU:HG	2.01	0.43
1:A:146:MET:HE2	1:A:212:THR:HG21	1.98	0.42
1:A:118:GLU:OE2	1:A:460:THR:CB	2.67	0.42
1:A:208:MET:HE2	4:A:856:HOH:O	2.19	0.42
1:A:388:TYR:CE2	1:A:393:LEU:CD2	3.03	0.42
1:A:161:GLN:O	1:A:165:LEU:HD12	2.21	0.41
1:A:169:ASP:HA	1:A:170:PRO:HD3	1.68	0.41
1:A:249:LYS:HE2	1:A:436:ASP:OD2	2.21	0.41
1:A:89:MET:O	1:A:92:PRO:HD2	2.21	0.41
1:A:213:LYS:NZ	1:A:237:GLU:O	2.53	0.41
1:A:126:ALA:HA	1:A:144:VAL:HG13	2.03	0.41
1:A:20:PRO:HB3	1:A:30:ILE:CG2	2.36	0.40
1:A:301:VAL:HG13	1:A:330:LEU:CB	2.51	0.40
1:A:148:SER:HB2	4:A:642:HOH:O	2.21	0.40
1:A:75:GLU:HG2	1:A:76:ARG:N	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	446/461 (97%)	430 (96%)	15 (3%)	1 (0%)	43	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	322	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	397/412 (96%)	383 (96%)	14 (4%)	32	32

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	75	GLU
1	A	77	ILE
1	A	83	HIS
1	A	118	GLU
1	A	173	LYS
1	A	175	ARG
1	A	176	LEU
1	A	181	SER
1	A	258	LEU
1	A	291	LYS
1	A	298	ARG
1	A	320	THR
1	A	333	ARG
1	A	404	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	97	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	179	GLN
1	A	342	GLN
1	A	401	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	AUO	A	502	-	60,62,62	1.61	9 (15%)	90,99,99	1.41	17 (18%)
2	UDP	A	501	-	25,26,26	2.02	7 (28%)	38,40,40	2.36	9 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AUO	A	502	-	-	5/25/146/146	0/8/7/7

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UDP	A	501	-	-	2/16/32/32	0/2/2/2

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	AUO	C17-C14	-7.19	1.41	1.51
2	A	501	UDP	PB-O2B	-4.23	1.39	1.54
2	A	501	UDP	PB-O3B	-3.78	1.40	1.54
2	A	501	UDP	PA-O1A	-3.48	1.38	1.50
2	A	501	UDP	PB-O1B	-3.27	1.40	1.50
2	A	501	UDP	PA-O2A	-2.96	1.41	1.55
3	A	502	AUO	O5-C5	-2.93	1.37	1.44
3	A	502	AUO	C22-C32	2.52	1.57	1.52
2	A	501	UDP	PA-O3A	-2.50	1.56	1.59
3	A	502	AUO	C3-C4	2.47	1.58	1.52
2	A	501	UDP	C6-N1	-2.46	1.32	1.38
3	A	502	AUO	O51-C11	-2.31	1.35	1.41
3	A	502	AUO	O5-C1	-2.30	1.35	1.41
3	A	502	AUO	O4-C4	-2.19	1.37	1.43
3	A	502	AUO	C62-C22	2.11	1.57	1.52
3	A	502	AUO	C13-C8	2.11	1.56	1.53

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	UDP	O2-C2-N1	-7.46	113.09	122.80
2	A	501	UDP	C4-N3-C2	-6.82	118.14	126.61
2	A	501	UDP	N3-C2-N1	5.86	122.51	114.89
2	A	501	UDP	O4-C4-C5	-4.00	118.25	125.16
3	A	502	AUO	O32-C1-O5	-3.90	100.42	110.69
3	A	502	AUO	C1-O5-C5	3.87	121.28	113.72
2	A	501	UDP	C5-C4-N3	2.92	118.89	114.80
3	A	502	AUO	O32-C1-C2	2.91	115.25	108.09
3	A	502	AUO	C12-C13-C8	2.87	116.60	111.05
3	A	502	AUO	C1-O32-C32	2.77	124.54	117.98
3	A	502	AUO	C11-O22-C22	2.74	124.46	117.98
3	A	502	AUO	O52-C52-C64	2.68	113.08	106.44
3	A	502	AUO	O22-C11-O51	-2.59	103.87	110.69
2	A	501	UDP	O4-C4-N3	2.46	122.84	119.27
3	A	502	AUO	C63-C53-C43	-2.42	108.42	110.92
3	A	502	AUO	C10-C9-C8	2.35	112.36	108.48
3	A	502	AUO	O51-C51-C61	2.32	112.19	106.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	AUO	O19-C62-C22	2.26	112.73	108.11
3	A	502	AUO	O-C55-C23	-2.20	117.84	122.97
2	A	501	UDP	C1'-N1-C2	2.19	121.53	117.59
3	A	502	AUO	C63-C53-C23	-2.15	111.51	115.12
2	A	501	UDP	O2A-PA-O3A	-2.15	101.46	107.27
3	A	502	AUO	C-C33-C43	-2.09	110.74	113.19
3	A	502	AUO	C21-C31-C41	-2.07	107.20	110.83
3	A	502	AUO	C17-C14-C19	-2.06	124.18	126.70
2	A	501	UDP	O3'-C3'-C4'	2.06	116.99	111.08

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	502	AUO	C2-C1-O32-C32
3	A	502	AUO	C14-C18-O19-C62
3	A	502	AUO	O51-C51-C61-O61
2	A	501	UDP	PB-O3A-PA-O5'
3	A	502	AUO	O5-C1-O32-C32
3	A	502	AUO	C32-C22-O22-C11
2	A	501	UDP	PA-O3A-PB-O1B

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	AUO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	448/461 (97%)	0.02	16 (3%) 46 45	16, 27, 56, 82	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	330	LEU	6.8
1	A	329	PHE	4.0
1	A	460	THR	3.9
1	A	321	TRP	3.0
1	A	333	ARG	3.0
1	A	318	GLY	2.9
1	A	83	HIS	2.5
1	A	332	GLU	2.5
1	A	69	ASP	2.5
1	A	263	ARG	2.4
1	A	288	VAL	2.4
1	A	85	PRO	2.4
1	A	326	PRO	2.3
1	A	175	ARG	2.2
1	A	403	TRP	2.1
1	A	182	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	AUO	A	502	56/56	0.89	0.08	20,20,20,20	0
2	UDP	A	501	25/25	0.97	0.06	18,23,28,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.