



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 2, 2025 – 12:48 am GMT

PDB ID : 6Z2A / pdb_00006z2a
Title : Structure of Clr4 mutant - F256A/F310A/F427A bound to SAH
Authors : Stirpe, A.; Schalch, T.
Deposited on : 2020-05-15
Resolution : 2.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1096 (2.46-2.46)
Clashscore	180529	1178 (2.46-2.46)
Ramachandran outliers	177936	1170 (2.46-2.46)
Sidechain outliers	177891	1170 (2.46-2.46)
RSRZ outliers	164620	1096 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	9% 89% 7% .
2	B	294	7% 87% 6% 6%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4495 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-lysine N-methyltransferase, H3 lysine-9 specific.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	283	Total	C	N	O	S	0	0	0
			2228	1391	384	436	17			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	256	ALA	PHE	engineered mutation	UNP O60016
A	310	ALA	PHE	engineered mutation	UNP O60016
A	427	ALA	PHE	engineered mutation	UNP O60016

- Molecule 2 is a protein called Histone-lysine N-methyltransferase, H3 lysine-9 specific.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	276	Total	C	N	O	S	0	0	0
			2183	1364	376	426	17			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	256	ALA	PHE	engineered mutation	UNP O60016
B	310	ALA	PHE	engineered mutation	UNP O60016
B	427	ALA	PHE	engineered mutation	UNP O60016

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Zn	0	0
			4	4		
3	B	4	Total	Zn	0	0
			4	4		

- Molecule 4 is S-ADENOSYL-L-HOMOCYSTEINE (CCD ID: SAH) (formula: C₁₄H₂₀N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
4	B	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		
5	B	1	Total	Mg	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	12	Total	O	0	0
			12	12		
6	B	10	Total	O	0	0
			10	10		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histone-lysine N-methyltransferase, H3 lysine-9 specific



- Molecule 2: Histone-lysine N-methyltransferase, H3 lysine-9 specific



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	92.44Å 110.29Å 70.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	70.68 – 2.46 70.68 – 2.46	Depositor EDS
% Data completeness (in resolution range)	72.2 (70.68-2.46) 64.0 (70.68-2.46)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.49 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.239 , 0.250 0.238 , 0.247	Depositor DCC
R_{free} test set	975 reflections (3.60%)	wwPDB-VP
Wilson B-factor (Å ²)	24.0	Xtriage
Anisotropy	0.133	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 7.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.54$, $\langle L^2 \rangle = 0.38$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4495	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.80 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0868e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.00	1/2276 (0.0%)	1.26	0/3079
2	B	0.97	0/2231	1.26	1/3019 (0.0%)
All	All	0.99	1/4507 (0.0%)	1.26	1/6098 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	406	ARG	C-O	-5.58	1.16	1.24

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	473	LEU	N-CA-C	-6.55	105.16	113.02

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2228	0	2096	12	0
2	B	2183	0	2056	10	0
3	A	4	0	0	0	0
3	B	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	26	0	19	0	0
4	B	26	0	19	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	12	0	0	0	0
6	B	10	0	0	0	0
All	All	4495	0	4190	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:418:ILE:HD11	1:A:433:LEU:C	2.15	0.72
1:A:418:ILE:HD11	1:A:433:LEU:CA	2.30	0.62
2:B:374:TYR:CE2	2:B:473:LEU:CD2	2.89	0.56
1:A:418:ILE:HD11	1:A:433:LEU:HB3	1.91	0.53
1:A:383:PHE:CZ	1:A:454:ALA:HB1	2.44	0.53
2:B:224:VAL:O	2:B:333:PHE:HA	2.10	0.52
1:A:224:VAL:O	1:A:333:PHE:HA	2.11	0.51
2:B:322:VAL:HG13	2:B:356:CYS:SG	2.55	0.47
2:B:244:ARG:O	2:B:360:GLU:HA	2.15	0.47
2:B:374:TYR:CE2	2:B:473:LEU:HD23	2.50	0.46
2:B:374:TYR:CD2	2:B:473:LEU:HD23	2.50	0.46
1:A:244:ARG:O	1:A:360:GLU:HA	2.17	0.45
2:B:323:GLN:HB2	2:B:423:ARG:NH2	2.33	0.44
1:A:323:GLN:HB2	1:A:423:ARG:NH2	2.33	0.43
1:A:257:GLN:NE2	1:A:424:ASN:OD1	2.51	0.43
1:A:418:ILE:HD11	1:A:433:LEU:CB	2.48	0.43
1:A:240:ILE:HD13	1:A:401:TYR:HB3	2.01	0.43
1:A:322:VAL:HG13	1:A:356:CYS:SG	2.59	0.43
1:A:383:PHE:CE1	1:A:454:ALA:HB1	2.56	0.41
2:B:240:ILE:HD13	2:B:401:TYR:HB3	2.03	0.41
2:B:212:LYS:NZ	2:B:235:LEU:O	2.52	0.41
2:B:265:LEU:HD12	2:B:266:GLY:N	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	277/296 (94%)	268 (97%)	9 (3%)	0	100	100
2	B	272/294 (92%)	262 (96%)	10 (4%)	0	100	100
All	All	549/590 (93%)	530 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	243/256 (95%)	240 (99%)	3 (1%)	67	79
2	B	238/255 (93%)	234 (98%)	4 (2%)	56	70
All	All	481/511 (94%)	474 (98%)	7 (2%)	60	74

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	265	LEU
1	A	272	ASN
1	A	461	GLN
2	B	261	ASN
2	B	264	SER
2	B	272	ASN
2	B	295	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	257	GLN
1	A	272	ASN
1	A	308	ASN
1	A	424	ASN
2	B	261	ASN
2	B	272	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 10 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SAH	B	505	-	24,28,28	0.71	0	25,40,40	0.96	2 (8%)
4	SAH	A	505	-	24,28,28	0.70	0	25,40,40	0.83	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SAH	B	505	-	-	1/11/31/31	0/3/3/3
4	SAH	A	505	-	-	1/11/31/31	0/3/3/3

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	505	SAH	C5-C6-N6	2.65	124.38	120.35
4	B	505	SAH	OXT-C-O	-2.40	118.63	124.09
4	A	505	SAH	C5-C6-N6	2.32	123.88	120.35

There are no chirality outliers.



All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	505	SAH	CB-CG-SD-C5'
4	A	505	SAH	CB-CG-SD-C5'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	283/296 (95%)	0.52	26 (9%) 16 16	13, 31, 62, 84	0
2	B	276/294 (93%)	0.41	21 (7%) 21 21	13, 29, 56, 80	0
All	All	559/590 (94%)	0.46	47 (8%) 18 18	13, 30, 59, 84	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	265	LEU	6.9
1	A	266	GLY	6.1
1	A	473	LEU	5.7
2	B	266	GLY	5.4
1	A	377	ASP	5.0
1	A	265	LEU	5.0
2	B	196	SER	4.6
1	A	376	ASP	4.0
2	B	472	LYS	3.6
1	A	460	VAL	3.6
1	A	461	GLN	3.5
2	B	377	ASP	3.4
1	A	475	ARG	3.4
1	A	195	ASP	3.4
2	B	453	GLY	3.3
1	A	263	SER	3.3
1	A	196	SER	3.3
2	B	263	SER	3.2
2	B	375	ASP	3.0
2	B	489	PHE	3.0
1	A	425	HIS	3.0
1	A	487	TRP	2.9
1	A	373	ASN	2.8
2	B	374	TYR	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	197	TYR	2.7
2	B	197	TYR	2.7
2	B	261	ASN	2.6
1	A	226	GLU	2.6
2	B	425	HIS	2.5
1	A	390	ASP	2.5
2	B	424	ASN	2.5
1	A	198	THR	2.5
1	A	474	ARG	2.5
2	B	264	SER	2.5
1	A	427	ALA	2.4
1	A	378	GLY	2.4
2	B	229	ASP	2.3
1	A	281	ASP	2.3
1	A	391	ALA	2.2
2	B	473	LEU	2.2
1	A	275	ARG	2.1
2	B	391	ALA	2.1
2	B	376	ASP	2.1
2	B	373	ASN	2.1
2	B	226	GLU	2.0
1	A	371	ASP	2.0
1	A	423	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	MG	A	506	1/1	0.67	0.13	30,30,30,30	0
4	SAH	A	505	26/26	0.89	0.10	25,28,30,30	0
5	MG	B	506	1/1	0.93	0.07	31,31,31,31	0
4	SAH	B	505	26/26	0.94	0.07	19,21,22,22	0
3	ZN	A	501	1/1	0.97	0.05	22,22,22,22	0
3	ZN	A	502	1/1	0.97	0.05	21,21,21,21	0
3	ZN	B	501	1/1	0.97	0.05	24,24,24,24	0
3	ZN	B	503	1/1	0.97	0.07	23,23,23,23	0
3	ZN	A	504	1/1	0.98	0.05	22,22,22,22	0
3	ZN	B	504	1/1	0.98	0.05	20,20,20,20	0
3	ZN	A	503	1/1	0.99	0.03	21,21,21,21	0
3	ZN	B	502	1/1	1.00	0.04	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.