



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 2, 2025 – 02:55 am GMT

PDB ID : 7OAT / pdb\_00007oat  
Title : Structural basis for targeted p97 remodelling by ASPL as prerequisite for p97 trimethylation by METTL21D  
Authors : Petrovic, S.; Heinemann, U.; Roske, Y.  
Deposited on : 2021-04-20  
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 1.8.4, CSD as541be (2020)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.46


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2511 (3.00-3.00)
Clashscore	180529	2866 (3.00-3.00)
Ramachandran outliers	177936	2778 (3.00-3.00)
Sidechain outliers	177891	2781 (3.00-3.00)
RSRZ outliers	164620	2523 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	190	 3% 79% 9% • 10%
2	B	481	 % 73% 15% • 10%
3	C	224	 % 80% 13% • 6%

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6558 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tether containing UBX domain for GLUT4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	171	1382	887	242	251	2	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	311	GLY	-	expression tag	UNP Q9BZE9
A	312	SER	-	expression tag	UNP Q9BZE9

- Molecule 2 is a protein called Transitional endoplasmic reticulum ATPase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	434	3385	2124	599	644	18	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	GLY	-	expression tag	UNP P55072
B	1	SER	-	expression tag	UNP P55072

- Molecule 3 is a protein called Protein-lysine methyltransferase METTL21D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	211	1659	1061	269	319	10	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	6	GLY	-	expression tag	UNP Q9H867




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	B	1	31	10	5	13	3	0	0

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
6	B	1	1	1	0	0

- Molecule 7 is S-ADENOSYL-L-HOMOCYSTEINE (CCD ID: SAH) (formula: C<sub>14</sub>H<sub>20</sub>N<sub>6</sub>O<sub>5</sub>S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
7	C	1	26	14	6	5	1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	7	Total	O	0	0
			7	7		
8	B	40	Total	O	0	0
			40	40		
8	C	11	Total	O	0	0
			11	11		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tether containing UBX domain for GLUT4



- Molecule 2: Transitional endoplasmic reticulum ATPase



- Molecule 3: Protein-lysine methyltransferase METTL21D



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.26Å 69.62Å 140.18Å 90.00° 94.41° 90.00°	Depositor
Resolution (Å)	49.32 – 3.00 49.32 – 3.00	Depositor EDS
% Data completeness (in resolution range)	98.1 (49.32-3.00) 98.1 (49.32-3.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.11 (at 3.01Å)	Xtrriage
Refinement program	BUSTER 2.10.3 (6-FEB-2020)	Depositor
R, $R_{free}$	0.225 , 0.270 0.240 , 0.292	Depositor DCC
$R_{free}$ test set	1042 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	73.6	Xtrriage
Anisotropy	0.330	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 65.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6558	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	78.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, M3L, ATP, SAH, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.62	0/1414	0.72	0/1918
2	B	0.64	0/3426	0.74	1/4633 (0.0%)
3	C	0.61	0/1689	0.70	0/2282
All	All	0.63	0/6529	0.73	1/8833 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	315	M3L	O-C-N	8.12	135.99	123.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1382	0	1374	16	0
2	B	3385	0	3416	45	0
3	C	1659	0	1643	15	0
4	B	8	0	12	0	0
4	C	8	0	12	1	0
5	B	31	0	12	0	0
6	B	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	C	26	0	19	0	0
8	A	7	0	0	0	0
8	B	40	0	0	0	0
8	C	11	0	0	0	0
All	All	6558	0	6488	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:372:VAL:HG11	1:A:376:PHE:HD2	1.30	0.95
2:B:284:SER:HA	2:B:287:ARG:HD3	1.53	0.90
2:B:394:VAL:HA	2:B:449:MET:HG2	1.54	0.89
1:A:372:VAL:HG13	1:A:376:PHE:HB3	1.64	0.80
1:A:372:VAL:HG11	1:A:376:PHE:CD2	2.18	0.76
1:A:372:VAL:CG1	1:A:376:PHE:CD2	2.71	0.73
1:A:372:VAL:CG1	1:A:376:PHE:HD2	1.99	0.73
1:A:355:ARG:HG2	2:B:222:LEU:HD21	1.74	0.70
2:B:284:SER:HA	2:B:287:ARG:CD	2.21	0.70
2:B:270:ASN:HB3	2:B:273:GLU:HB3	1.77	0.67
2:B:314:GLU:OE2	3:C:179:GLY:HA3	1.95	0.66
3:C:66:LEU:HB3	3:C:90:ALA:HB2	1.76	0.66
2:B:31:ALA:HA	2:B:83:ARG:HB3	1.79	0.64
1:A:372:VAL:CG1	1:A:376:PHE:HB3	2.26	0.63
2:B:116:VAL:HG22	2:B:165:VAL:HG12	1.79	0.62
1:A:347:LEU:HD13	2:B:226:HIS:CE1	2.36	0.61
1:A:372:VAL:HG13	1:A:376:PHE:CB	2.31	0.60
1:A:412:THR:HA	1:A:448:THR:HA	1.84	0.59
2:B:113:ARG:HG3	2:B:181:VAL:HB	1.87	0.56
1:A:372:VAL:HG13	1:A:376:PHE:CD2	2.42	0.56
2:B:211:LYS:HD3	2:B:211:LYS:H	1.73	0.52
2:B:125:GLY:HA2	2:B:436:THR:HG22	1.92	0.52
2:B:40:SER:HB2	2:B:83:ARG:HB2	1.93	0.51
2:B:136:LYS:O	2:B:140:LEU:HB2	2.12	0.50
1:A:421:ARG:HA	1:A:424:LEU:HD12	1.94	0.50
3:C:153:GLU:O	3:C:157:LYS:HG2	2.11	0.49
2:B:181:VAL:HG12	2:B:183:HIS:CD2	2.48	0.48
1:A:396:LEU:HD23	1:A:461:HIS:NE2	2.28	0.48
1:A:323:VAL:HG23	1:A:404:GLN:HB3	1.96	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:249:THR:HG22	2:B:251:LYS:HD3	1.95	0.47
2:B:119:ILE:O	2:B:122:THR:HG22	2.15	0.46
2:B:74:ASP:HB3	2:B:77:CYS:HB2	1.96	0.46
2:B:185:GLU:H	2:B:185:GLU:HG3	1.44	0.46
2:B:153:LEU:HD11	2:B:160:ALA:HB1	1.97	0.46
2:B:315:M3L:CM2	3:C:43:TRP:CE2	2.99	0.46
3:C:71:VAL:HG13	3:C:92:VAL:HG22	1.97	0.46
1:A:347:LEU:HD13	2:B:226:HIS:NE2	2.31	0.45
2:B:315:M3L:HM13	3:C:147:TYR:CD1	2.53	0.44
2:B:181:VAL:HG12	2:B:183:HIS:HD2	1.83	0.44
2:B:240:GLY:HA2	2:B:343:VAL:O	2.17	0.44
2:B:122:THR:HG23	2:B:161:VAL:HG13	1.99	0.44
2:B:153:LEU:HD23	2:B:162:GLU:HG2	2.00	0.43
2:B:315:M3L:HM13	3:C:147:TYR:CG	2.53	0.43
2:B:187:GLU:HG2	2:B:188:PRO:HD2	2.01	0.43
2:B:136:LYS:HB3	2:B:137:PRO:HD3	2.01	0.43
2:B:56:THR:HG21	2:B:108:VAL:HG21	2.01	0.43
2:B:224:LEU:HD22	2:B:298:PRO:HB2	2.01	0.42
3:C:201:ILE:HD11	3:C:219:TYR:CG	2.54	0.42
2:B:192:GLU:HB2	2:B:195:GLU:HG3	2.02	0.42
2:B:223:PRO:HB3	2:B:237:PRO:HB3	2.00	0.42
2:B:153:LEU:HD21	2:B:160:ALA:HB1	2.00	0.42
2:B:287:ARG:HA	2:B:331:LEU:HD21	2.01	0.42
2:B:26:LEU:HD13	2:B:41:LEU:HD21	2.02	0.41
1:A:347:LEU:HD23	1:A:351:ASP:HB3	2.02	0.41
2:B:106:PRO:HA	2:B:107:ASP:HA	1.87	0.41
2:B:385:THR:HA	2:B:388:MET:HG2	2.03	0.41
3:C:17:VAL:HG12	3:C:31:GLN:HA	2.02	0.41
3:C:152:LEU:HD22	3:C:187:LYS:HB3	2.03	0.41
3:C:169:ILE:HB	3:C:220:ILE:HB	2.03	0.41
2:B:116:VAL:CG1	2:B:163:PHE:HB3	2.51	0.41
3:C:20:LEU:HA	4:C:301:EDO:H22	2.03	0.41
2:B:201:VAL:HG12	2:B:257:ALA:HB2	2.03	0.41
2:B:243:LEU:HD11	2:B:344:MET:HE2	2.03	0.40
2:B:299:ALA:HB3	2:B:341:VAL:HG22	2.02	0.40
3:C:113:HIS:CD2	3:C:114:LEU:HG	2.56	0.40
2:B:315:M3L:HM22	3:C:43:TRP:CE2	2.56	0.40
2:B:359:ARG:HD2	3:C:212:SER:OG	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	165/190 (87%)	155 (94%)	10 (6%)	0	100	100
2	B	429/481 (89%)	413 (96%)	14 (3%)	2 (0%)	25	61
3	C	209/224 (93%)	201 (96%)	7 (3%)	1 (0%)	25	61
All	All	803/895 (90%)	769 (96%)	31 (4%)	3 (0%)	30	66

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	362	ARG
3	C	147	TYR
2	B	106	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	149/164 (91%)	138 (93%)	11 (7%)	11	38
2	B	367/409 (90%)	338 (92%)	29 (8%)	10	35
3	C	180/197 (91%)	167 (93%)	13 (7%)	12	39
All	All	696/770 (90%)	643 (92%)	53 (8%)	11	37

All (53) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	323	VAL
1	A	332	ARG
1	A	347	LEU
1	A	349	VAL
1	A	355	ARG
1	A	364	LYS
1	A	366	LEU
1	A	385	LEU
1	A	454	LEU
1	A	470	VAL
1	A	476	LEU
2	B	24	ASN
2	B	55	ASP
2	B	65	ARG
2	B	108	VAL
2	B	122	THR
2	B	146	ILE
2	B	153	LEU
2	B	155	ARG
2	B	158	MET
2	B	167	GLU
2	B	171	SER
2	B	173	TYR
2	B	185	GLU
2	B	191	ARG
2	B	194	GLU
2	B	200	GLU
2	B	211	LYS
2	B	220	VAL
2	B	242	LEU
2	B	278	LEU
2	B	286	LEU
2	B	287	ARG
2	B	327	GLN
2	B	328	LEU
2	B	329	LEU
2	B	335	LEU
2	B	359	ARG
2	B	402	GLU
2	B	449	MET
3	C	13	LEU
3	C	20	LEU
3	C	34	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
3	C	67	SER
3	C	71	VAL
3	C	73	GLU
3	C	83	LEU
3	C	99	GLU
3	C	109	ASN
3	C	124	LEU
3	C	144	ASP
3	C	159	LEU
3	C	183	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	36	ASN
2	B	103	GLN
2	B	327	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	M3L	B	315	2	10,11,12	0.99	1 (10%)	9,14,16	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	M3L	B	315	2	-	1/9/10/12	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	315	M3L	O-C	3.02	1.32	1.19

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	315	M3L	CE-CD-CG-CB

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	315	M3L	4	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	EDO	B	502	-	3,3,3	0.56	0	2,2,2	0.35	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	EDO	C	301	-	3,3,3	0.57	0	2,2,2	0.34	0
4	EDO	B	501	-	3,3,3	0.64	0	2,2,2	0.29	0
4	EDO	C	303	-	3,3,3	0.55	0	2,2,2	0.35	0
5	ATP	B	503	6	26,33,33	0.62	0	31,52,52	0.62	1 (3%)
7	SAH	C	302	-	24,28,28	0.68	0	25,40,40	1.12	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	502	-	-	0/1/1/1	-
4	EDO	C	301	-	-	0/1/1/1	-
4	EDO	B	501	-	-	0/1/1/1	-
4	EDO	C	303	-	-	0/1/1/1	-
5	ATP	B	503	6	-	3/18/38/38	0/3/3/3
7	SAH	C	302	-	-	5/11/31/31	0/3/3/3

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	302	SAH	CB-CG-SD	-3.23	106.07	113.31
7	C	302	SAH	CB-CA-N	2.78	117.46	110.17
7	C	302	SAH	C5-C6-N6	2.30	123.84	120.35
5	B	503	ATP	C5-C6-N6	2.23	123.75	120.35

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	B	503	ATP	PB-O3B-PG-O2G
5	B	503	ATP	O4'-C4'-C5'-O5'
5	B	503	ATP	C3'-C4'-C5'-O5'
7	C	302	SAH	N-CA-CB-CG
7	C	302	SAH	C-CA-CB-CG
7	C	302	SAH	O-C-CA-N
7	C	302	SAH	OXT-C-CA-N
7	C	302	SAH	CB-CG-SD-C5'


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	301	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	171/190 (90%)	0.38	5 (2%) 54 32	66, 84, 118, 129	0
2	B	433/481 (90%)	0.22	7 (1%) 70 49	52, 75, 94, 99	0
3	C	211/224 (94%)	0.20	3 (1%) 73 52	61, 75, 88, 102	0
All	All	815/895 (91%)	0.25	15 (1%) 67 45	52, 77, 101, 129	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	372	VAL	4.8
2	B	23	PRO	3.7
1	A	373	THR	3.2
3	C	13	LEU	3.1
1	A	366	LEU	3.1
3	C	61	ASP	2.9
2	B	364	ASP	2.8
3	C	152	LEU	2.6
2	B	244	TYR	2.6
2	B	348	ASN	2.3
2	B	338	ARG	2.2
2	B	193	ASP	2.2
1	A	333	LEU	2.1
1	A	385	LEU	2.1
2	B	106	PRO	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	M3L	B	315	12/13	0.93	0.13	79,79,81,129	0

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	EDO	C	303	4/4	0.63	0.21	87,87,87,87	0
4	EDO	B	502	4/4	0.80	0.15	72,72,72,72	0
4	EDO	B	501	4/4	0.88	0.12	60,60,60,60	0
5	ATP	B	503	31/31	0.89	0.12	75,77,81,81	0
6	MG	B	504	1/1	0.89	0.07	72,72,72,72	0
4	EDO	C	301	4/4	0.90	0.14	73,73,73,73	0
7	SAH	C	302	26/26	0.92	0.11	75,77,77,77	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around ATP B 503:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around SAH C 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.