



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 24, 2025 – 06:26 am BST

PDB ID : 7OBN / pdb_00007obn
Title : Structural investigations of a new L3 DNA ligase: structure-function analysis
Authors : Leiros, H.-K.S.; Williamson, A.
Deposited on : 2021-04-23
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1096 (2.46-2.46)
Clashscore	180529	1178 (2.46-2.46)
Ramachandran outliers	177936	1170 (2.46-2.46)
Sidechain outliers	177891	1170 (2.46-2.46)
RSRZ outliers	164620	1096 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	21	 81% 19%
2	C	21	 81% 14% 5%
3	A	321	 19% 60% 28% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	AMP	A	401	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3303 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*TP*TP*CP*CP*GP*AP*TP*AP*GP*TP*GP*GP*GP*TP*CP*GP*CP*AP*AP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	21	Total	C	N	O	P	0	0	0
			431	206	79	126	20			


- Molecule 2 is a DNA chain called DNA/RNA (5'-D(*AP*TP*TP*GP*CP*GP*AP*C)-R(P*(OMC))-D(P*CP*CP*AP*CP*TP*AP*TP*CP*GP*GP*AP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	21	Total	C	N	O	P	0	0	0
			426	204	79	123	20			

- Molecule 3 is a protein called DNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	299	Total	C	N	O	S	53	1	0
			2359	1494	420	432	13			

- Molecule 4 is ADENOSINE MONOPHOSPHATE (CCD ID: AMP) (formula: C₁₀H₁₄N₅O₇P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			22	10	5	6	1		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	12	Total 12	O 12	0	0
6	C	13	Total 13	O 13	0	0
6	A	35	Total 35	O 35	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA (5'-D(*TP*TP*CP*CP*GP*AP*TP*AP*GP*TP*GP*GP*GP*GP*TP*CP*GP*CP*AP*AP*T)-3')

Chain B: 




- Molecule 2: DNA/RNA (5'-D(*AP*TP*TP*GP*CP*GP*AP*C)-R(P*(OMC))-D(P*CP*CP*AP*CP*TP*AP*TP*CP*GP*GP*AP*A)-3')

Chain C: 



- Molecule 3: DNA ligase

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	125.47Å 45.55Å 111.46Å 90.00° 103.94° 90.00°	Depositor
Resolution (Å)	24.62 – 2.45 24.62 – 2.45	Depositor EDS
% Data completeness (in resolution range)	96.0 (24.62-2.45) 95.9 (24.62-2.45)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.17 (at 2.44Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874, PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.257 , 0.283 0.258 , 0.283	Depositor DCC
R_{free} test set	2028 reflections (8.85%)	wwPDB-VP
Wilson B-factor (Å ²)	56.1	Xtriage
Anisotropy	0.328	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 64.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3303	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: OMC, AMP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.19	0/483	0.39	0/745
2	C	0.21	0/454	0.41	0/698
3	A	0.60	5/2412 (0.2%)	1.20	27/3252 (0.8%)
All	All	0.52	5/3349 (0.1%)	1.02	27/4695 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	A	0	8

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	22	LYS	CD-CE	12.50	1.90	1.52
3	A	308	HIS	CA-CB	-7.44	1.47	1.52
3	A	308	HIS	N-CA	7.28	1.51	1.46
3	A	22	LYS	CE-NZ	7.17	1.70	1.49
3	A	308	HIS	C-O	-5.44	1.21	1.23

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	211	GLU	CG-CD-OE1	19.57	163.41	118.40
3	A	211	GLU	OE1-CD-OE2	-18.43	78.66	122.90
3	A	211	GLU	CG-CD-OE2	-14.46	85.14	118.40
3	A	215	ARG	NE-CZ-NH1	-10.22	111.28	121.50
3	A	161	LEU	CB-CG-CD2	-9.25	82.96	110.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	22	LYS	CB-CG-CD	-8.96	90.70	111.30
3	A	22	LYS	CB-CA-C	8.26	123.58	110.19
3	A	7	PHE	CA-C-N	8.07	136.77	122.48
3	A	7	PHE	C-N-CA	8.07	136.77	122.48
3	A	21	ILE	CA-C-N	-7.61	112.65	122.77
3	A	21	ILE	C-N-CA	-7.61	112.65	122.77
3	A	161	LEU	CD1-CG-CD2	-7.56	94.17	110.80
3	A	199	ARG	CG-CD-NE	7.24	127.93	112.00
3	A	215	ARG	CG-CD-NE	7.01	127.41	112.00
3	A	161	LEU	CB-CG-CD1	6.88	131.35	110.70
3	A	22	LYS	CD-CE-NZ	-6.81	90.11	111.90
3	A	105	PHE	CB-CA-C	6.26	120.00	110.37
3	A	291	ARG	CB-CA-C	-6.23	99.02	109.48
3	A	211	GLU	CB-CG-CD	6.13	123.02	112.60
3	A	199	ARG	CD-NE-CZ	6.06	132.89	124.40
3	A	215	ARG	CD-NE-CZ	6.00	132.80	124.40
3	A	291	ARG	CA-CB-CG	6.00	126.09	114.10
3	A	294	LYS	CB-CG-CD	5.71	124.44	111.30
3	A	199	ARG	NE-CZ-NH1	-5.54	115.96	121.50
3	A	79	PRO	N-CA-C	5.48	123.76	112.47
3	A	105	PHE	N-CA-C	-5.48	106.05	114.16
3	A	159	ASP	N-CA-C	-5.39	103.38	110.33

There are no chirality outliers.

All (8) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	A	105	PHE	Sidechain
3	A	199	ARG	Sidechain
3	A	215	ARG	Sidechain
3	A	22	LYS	Peptide
3	A	23	PHE	Peptide
3	A	4	PRO	Peptide
3	A	5	THR	Peptide
3	A	78	SER	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	431	0	239	3	0
2	C	426	0	239	5	0
3	A	2359	0	2303	100	0
4	A	22	0	12	10	0
5	A	5	0	0	0	0
6	A	35	0	0	3	0
6	B	12	0	0	1	0
6	C	13	0	0	2	0
All	All	3303	0	2793	104	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (104) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:22:LYS:NZ	3:A:22:LYS:CE	1.70	1.52
3:A:22:LYS:CE	3:A:22:LYS:CD	1.90	1.48
3:A:138:ARG:C	3:A:143:ASP:CA	2.45	0.90
3:A:72:GLY:HA2	3:A:105:PHE:CD1	2.17	0.79
3:A:78:SER:O	3:A:80:THR:N	2.15	0.78
3:A:66:LEU:O	3:A:130:ARG:NH2	2.20	0.75
3:A:22:LYS:NZ	3:A:22:LYS:CD	2.55	0.70
3:A:234:GLN:OE1	6:A:501:HOH:O	2.11	0.68
3:A:81:ASP:O	3:A:85:MET:HG3	1.94	0.67
3:A:7:PHE:HE1	3:A:9:PRO:HA	1.60	0.67
3:A:304:ASP:O	6:A:502:HOH:O	2.15	0.64
3:A:7:PHE:CE2	3:A:39:PHE:HE2	2.16	0.64
3:A:7:PHE:CE2	3:A:39:PHE:CE2	2.86	0.63
3:A:53:ASN:OD1	3:A:55:VAL:HG12	1.98	0.63
3:A:86:GLN:C	3:A:88:SER:H	2.06	0.63
3:A:51:ILE:HB	3:A:57:GLN:HE21	1.63	0.63
3:A:92:MET:HG3	3:A:93:SER:N	2.14	0.63
3:A:215:ARG:NH2	3:A:238:VAL:HG11	2.14	0.62
2:C:33:DA:N7	6:C:102:HOH:O	2.31	0.62
3:A:30:LYS:NZ	3:A:171:GLU:OE2	2.29	0.62
3:A:106:ASP:OD1	3:A:124:ARG:NH1	2.33	0.61
3:A:174:MET:HE2	4:A:401:AMP:C2	2.37	0.59
3:A:105:PHE:HE2	4:A:401:AMP:C4	2.21	0.59
3:A:7:PHE:HE2	3:A:39:PHE:CE2	2.21	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:84:CYS:C	3:A:86:GLN:H	2.11	0.58
3:A:215:ARG:NH2	3:A:217:GLU:OE1	2.37	0.58
3:A:76:VAL:HG22	3:A:85:MET:HA	1.86	0.57
3:A:105:PHE:CE2	4:A:401:AMP:C2	2.92	0.57
3:A:34:ILE:C	3:A:73:GLU:HG3	2.30	0.56
3:A:171:GLU:CD	3:A:198:LYS:NZ	2.63	0.56
3:A:75:THR:OG1	3:A:85:MET:SD	2.62	0.56
3:A:62:ALA:O	3:A:138:ARG:NH1	2.38	0.56
3:A:105:PHE:HE2	4:A:401:AMP:N3	2.04	0.55
3:A:59:PHE:CD1	3:A:100:PHE:HZ	2.24	0.54
3:A:76:VAL:HG12	3:A:100:PHE:HA	1.88	0.54
3:A:80:THR:OG1	3:A:81:ASP:N	2.41	0.53
3:A:105:PHE:HE2	4:A:401:AMP:C2	2.26	0.53
3:A:212:GLU:HB3	3:A:282:TRP:CH2	2.43	0.53
3:A:212:GLU:HG3	3:A:214:MET:HE2	1.89	0.53
3:A:86:GLN:O	3:A:86:GLN:HG2	2.08	0.53
3:A:72:GLY:HA2	3:A:105:PHE:CE1	2.43	0.52
3:A:106:ASP:OD1	3:A:107:TRP:N	2.43	0.52
3:A:102:PHE:CE2	3:A:104:VAL:HG22	2.45	0.52
3:A:282:TRP:O	3:A:285:ARG:NH2	2.42	0.52
3:A:52:PRO:HG2	3:A:89:MET:HE3	1.92	0.51
3:A:119:MET:SD	3:A:123:ARG:HD3	2.50	0.51
3:A:223:ASN:OD1	3:A:227:ARG:N	2.41	0.51
3:A:171:GLU:CD	3:A:198:LYS:HZ2	2.19	0.51
3:A:56:VAL:O	3:A:59:PHE:HB3	2.11	0.51
3:A:122:ASP:N	3:A:178:HIS:O	2.45	0.50
3:A:250:ARG:NH2	3:A:304:ASP:HA	2.26	0.50
3:A:56:VAL:HG12	3:A:100:PHE:CD2	2.46	0.50
3:A:215:ARG:HH21	3:A:238:VAL:CG1	2.25	0.50
3:A:16:THR:N	3:A:189:GLU:OE2	2.41	0.50
3:A:207:ILE:HA	3:A:248:VAL:HG12	1.94	0.49
3:A:250:ARG:HH22	3:A:305:ALA:H	1.60	0.49
3:A:119:MET:HB3	3:A:124:ARG:HG3	1.95	0.49
3:A:215:ARG:NH2	3:A:238:VAL:CG1	2.75	0.49
3:A:7:PHE:HE2	3:A:39:PHE:CZ	2.30	0.49
2:C:32:DC:OP2	3:A:46:ARG:NH1	2.46	0.48
3:A:91:VAL:HG12	3:A:91:VAL:O	2.13	0.48
3:A:12:ALA:HB2	3:A:174:MET:HE1	1.97	0.47
2:C:36:DA:N7	6:C:103:HOH:O	2.35	0.47
3:A:159:ASP:C	3:A:161:LEU:H	2.23	0.47
3:A:292:VAL:HG11	3:A:315:ARG:HD3	1.95	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:198:LYS:HE2	3:A:298:PHE:CD2	2.50	0.47
3:A:54:PRO:HD3	3:A:91:VAL:HA	1.96	0.46
3:A:33:GLY:HA3	3:A:74:LEU:O	2.15	0.46
3:A:7:PHE:CE1	3:A:9:PRO:HA	2.46	0.46
3:A:153:VAL:HG12	3:A:157:ASP:OD2	2.15	0.46
3:A:198:LYS:HD3	3:A:298:PHE:CE2	2.50	0.46
2:C:30:OMC:OP1	3:A:47:SER:OG	2.33	0.46
3:A:73:GLU:OE1	4:A:401:AMP:C8	2.69	0.46
3:A:131:ARG:O	3:A:133:ALA:N	2.49	0.45
3:A:127:ILE:O	3:A:130:ARG:HG3	2.17	0.45
3:A:78:SER:HA	3:A:85:MET:HE2	1.98	0.45
3:A:72:GLY:HA2	3:A:105:PHE:HD1	1.72	0.45
3:A:250:ARG:HH12	3:A:265:PRO:HD2	1.81	0.44
3:A:86:GLN:C	3:A:88:SER:N	2.71	0.44
3:A:105:PHE:HD2	4:A:401:AMP:C6	2.34	0.44
3:A:131:ARG:O	3:A:132:ILE:C	2.60	0.44
3:A:215:ARG:HH21	3:A:238:VAL:HG11	1.82	0.44
2:C:32:DC:H2"	3:A:308:HIS:NE2	2.33	0.44
3:A:119:MET:HE1	3:A:123:ARG:HG2	1.99	0.44
3:A:291:ARG:NH2	3:A:316:HIS:O	2.51	0.44
3:A:308:HIS:N	3:A:309:PRO:HD3	2.32	0.44
3:A:105:PHE:CD2	4:A:401:AMP:C6	3.06	0.43
3:A:212:GLU:HB3	3:A:282:TRP:CZ2	2.54	0.43
3:A:174:MET:HG3	4:A:401:AMP:C6	2.53	0.43
3:A:7:PHE:CE2	3:A:39:PHE:CZ	3.05	0.43
3:A:70:LEU:HD23	3:A:107:TRP:CD1	2.53	0.42
1:B:8:DA:OP1	3:A:228:THR:OG1	2.37	0.42
3:A:286:ASP:OD1	6:A:503:HOH:O	2.22	0.42
3:A:32:ASP:HB3	3:A:82:ALA:HB2	2.02	0.42
3:A:174:MET:HE2	4:A:401:AMP:N3	2.35	0.42
1:B:16:DC:H2"	1:B:17:DG:C8	2.54	0.41
3:A:78:SER:HB2	3:A:79:PRO:HD3	2.01	0.41
3:A:71:ASP:O	3:A:105:PHE:HD1	2.04	0.41
3:A:123:ARG:O	3:A:126:GLU:HB3	2.20	0.41
1:B:12:DG:N7	6:B:103:HOH:O	2.37	0.41
3:A:32:ASP:OD1	3:A:171:GLU:OE1	2.38	0.41
3:A:67:LEU:HD23	3:A:70:LEU:HD12	2.03	0.41
3:A:51:ILE:HG21	3:A:56:VAL:HG21	2.04	0.40
3:A:234:GLN:HA	3:A:237:LEU:HG	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	A	290/321 (90%)	246 (85%)	36 (12%)	8 (3%)	4 2

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	23	PHE
3	A	78	SER
3	A	79	PRO
3	A	4	PRO
3	A	5	THR
3	A	24	PRO
3	A	109	PRO
3	A	132	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	A	244/260 (94%)	244 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	86	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	A	234	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	OMC	C	30	1,2	19,22,23	1.47	3 (15%)	26,31,34	0.92	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OMC	C	30	1,2	-	3/9/27/28	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	30	OMC	C4-N4	3.86	1.43	1.33
2	C	30	OMC	C2-N1	-3.25	1.32	1.40
2	C	30	OMC	C6-C5	2.76	1.41	1.35

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	30	OMC	O2-C2-N3	-2.14	118.84	122.33

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	30	OMC	C3'-C4'-C5'-O5'
2	C	30	OMC	O4'-C4'-C5'-O5'
2	C	30	OMC	C3'-C2'-O2'-CM2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	30	OMC	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	AMP	A	401	-	18,24,25	1.10	2 (11%)	18,35,38	1.46	3 (16%)
5	SO4	A	402	-	4,4,4	0.17	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsions and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	AMP	A	401	-	-	1/3/25/26	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	401	AMP	C5-C4	2.64	1.47	1.40
4	A	401	AMP	C2-N3	2.26	1.35	1.32

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	401	AMP	N3-C2-N1	-3.59	123.07	128.68
4	A	401	AMP	C3'-C2'-C1'	3.03	105.54	100.98
4	A	401	AMP	C4-C5-N7	-2.46	106.83	109.40

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	401	AMP	C4'-C5'-O5'-P

There are no ring outliers.

1 monomer is involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	401	AMP	10	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	B	21/21 (100%)	0.17	0	100 100	47, 73, 103, 113	0
2	C	20/21 (95%)	0.20	0	100 100	50, 74, 95, 98	0
3	A	299/321 (93%)	1.17	62 (20%)	3 3	25, 66, 108, 133	25 (8%)
All	All	340/363 (93%)	1.05	62 (18%)	4 4	25, 67, 108, 133	25 (7%)

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	87	ASN	8.1
3	A	303	VAL	5.2
3	A	125	SER	4.9
3	A	99	ASP	4.8
3	A	227	ARG	4.4
3	A	110	LEU	4.4
3	A	3	ILE	4.3
3	A	77	GLY	4.1
3	A	92	MET	3.9
3	A	59	PHE	3.8
3	A	252	GLU	3.8
3	A	90	ALA	3.8
3	A	91	VAL	3.6
3	A	234	GLN	3.5
3	A	126	GLU	3.5
3	A	61	ALA	3.3
3	A	298	PHE	3.2
3	A	78	SER	3.2
3	A	101	THR	3.2
3	A	302	THR	3.1
3	A	7	PHE	3.1
3	A	63	TYR	3.1
3	A	64	ALA	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	A	132	ILE	2.9
3	A	21	ILE	2.8
3	A	300	HIS	2.8
3	A	41	GLY	2.8
3	A	144	ILE	2.8
3	A	160	THR	2.8
3	A	200	PHE	2.8
3	A	5	THR	2.7
3	A	111	ARG	2.7
3	A	2	THR	2.7
3	A	233	ALA	2.7
3	A	82	ALA	2.6
3	A	6	GLY	2.6
3	A	54	PRO	2.6
3	A	260	TYR	2.6
3	A	232	THR	2.6
3	A	65	SER	2.5
3	A	81	ASP	2.5
3	A	94	LYS	2.5
3	A	24	PRO	2.5
3	A	86	GLN	2.5
3	A	226	GLY	2.5
3	A	79	PRO	2.5
3	A	105	PHE	2.5
3	A	93	SER	2.4
3	A	128	VAL	2.3
3	A	177	ALA	2.3
3	A	55	VAL	2.2
3	A	119	MET	2.2
3	A	117	ASP	2.2
3	A	308	HIS	2.2
3	A	67	LEU	2.2
3	A	138	ARG	2.2
3	A	261	GLY	2.1
3	A	76	VAL	2.1
3	A	220	ALA	2.1
3	A	109	PRO	2.1
3	A	85	MET	2.1
3	A	13	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	OMC	C	30	21/22	0.92	0.13	56,64,72,80	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	AMP	A	401	22/23	0.77	0.16	48,56,73,83	22
5	SO4	A	402	5/5	0.97	0.08	46,49,64,79	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.