



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2025 – 07:03 pm BST

PDB ID : 7PRT / pdb\_00007prt  
Title : Crystal structure of human heparanase in complex with covalent inhibitor CB678  
Authors : Wu, L.; Armstrong, Z.; Davies, G.J.  
Deposited on : 2021-09-22  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.44


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	5161 (1.70-1.70)
Clashscore	180529	5671 (1.70-1.70)
Ramachandran outliers	177936	5594 (1.70-1.70)
Sidechain outliers	177891	5594 (1.70-1.70)
RSRZ outliers	164620	5159 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	389	
2	B	74	
3	C	2	
4	D	2	

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 3883 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heparanase 50 kDa subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	384	3080	1990	522	556	12	0	4	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	155	ASP	-	expression tag	UNP Q9Y251
A	156	PRO	-	expression tag	UNP Q9Y251
A	157	GLY	-	expression tag	UNP Q9Y251
A	307	ARG	LYS	variant	UNP Q9Y251

- Molecule 2 is a protein called Heparanase 8 kDa subunit.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	74	595	388	97	110	0	1	0

- Molecule 3 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	2	24	14	1	9	0	0	0

- Molecule 4 is an oligosaccharide called 2-deoxy-alpha-D-arabino-hexopyranose-(1-4)-(2R,3S,5R,6R)-2,3,4,5,6-pentakis(oxidanyl)cyclohexane-1-carboxylic acid.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
4	D	2	Total	C	O	0	0	0
			23	13	10			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	2	Total Cl 2 2	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	76	Total O 76 76	0	0
8	B	15	Total O 15 15	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Heparanase 50 kDa subunit




- Molecule 2: Heparanase 8 kDa subunit



- Molecule 3: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-deoxy-alpha-D-arabino-hexopyranose-(1-4)-(2R,3S,5R,6R)-2,3,4,5,6-pentakis(oxidanyl)cyclohexane-1-carboxylic acid



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.81Å 71.62Å 78.74Å 90.00° 94.89° 90.00°	Depositor
Resolution (Å)	52.95 – 1.70 52.95 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.7 (52.95-1.70) 99.7 (52.95-1.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.12 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.192 , 0.236 0.200 , 0.246	Depositor DCC
$R_{free}$ test set	2806 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.8	Xtrriage
Anisotropy	0.547	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 30.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3883	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.59% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, FUC, 8I4, Z61, EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.02	0/3155	1.31	2/4268 (0.0%)
2	B	1.02	0/609	1.19	0/826
All	All	1.02	0/3764	1.29	2/5094 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	225	GLU	CB-CA-C	6.06	116.27	111.00
1	A	270	GLN	CB-CA-C	5.46	118.07	109.27

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3080	0	3097	16	0
2	B	595	0	607	6	0
3	C	24	0	22	0	0
4	D	23	0	0	0	0
5	A	56	0	52	0	0
6	A	12	0	18	1	0
7	A	2	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	A	76	0	0	0	0
8	B	15	0	0	0	0
All	All	3883	0	3796	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:298[A]:TYR:CE2	1:A:345:SER:OG	2.35	0.79
1:A:298[A]:TYR:HE2	1:A:345:SER:OG	1.71	0.72
2:B:26:ILE:HG12	2:B:57:LEU:HD11	1.72	0.71
2:B:1:GLN:HE21	2:B:1:GLN:HA	1.57	0.68
1:A:502[A]:MET:SD	1:A:508:LEU:HD23	2.44	0.57
1:A:315:VAL:O	1:A:318:ILE:HG12	2.07	0.55
2:B:1:GLN:HA	2:B:1:GLN:NE2	2.25	0.48
1:A:430[B]:LYS:HE2	1:A:458:HIS:CE1	2.51	0.46
1:A:398:PHE:HB2	2:B:44:LYS:HD3	1.99	0.45
1:A:164:THR:HA	2:B:68:ILE:O	2.18	0.43
1:A:489:LEU:HG	6:A:606:EDO:H22	2.00	0.43
1:A:313:PRO:HG3	1:A:430[B]:LYS:HG2	1.99	0.43
2:B:9:PHE:CZ	2:B:11[A]:GLN:HB2	2.54	0.42
1:A:191:LEU:HD23	1:A:191:LEU:HA	1.92	0.42
1:A:365:TRP:CH2	1:A:382:ARG:HB2	2.55	0.41
1:A:430[B]:LYS:HE2	1:A:458:HIS:ND1	2.34	0.41
1:A:295:TRP:CD2	1:A:326:VAL:HG21	2.55	0.41
1:A:404:TYR:CD1	1:A:404:TYR:C	2.99	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/389 (99%)	375 (97%)	10 (3%)	1 (0%)	37	23
2	B	73/74 (99%)	72 (99%)	1 (1%)	0	100	100
All	All	459/463 (99%)	447 (97%)	11 (2%)	1 (0%)	44	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	428	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/338 (100%)	324 (96%)	13 (4%)	27	11
2	B	67/66 (102%)	66 (98%)	1 (2%)	60	47
All	All	404/404 (100%)	390 (96%)	14 (4%)	31	15

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	163	SER
1	A	197	LEU
1	A	231	LYS
1	A	234	ASP
1	A	273	ARG
1	A	284	LYS
1	A	390	ASN
1	A	391	TYR
1	A	392	HIS
1	A	396	GLU
1	A	429	ARG
1	A	431	LEU
1	A	498	LEU
2	B	1	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	248	GLN
1	A	486	HIS
1	A	494	GLN
2	B	1	GLN
2	B	15	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	C	1	1,3	14,14,15	0.94	0	17,19,21	2.17	4 (23%)
3	FUC	C	2	3	10,10,11	0.89	0	14,14,16	1.49	2 (14%)
4	8I4	D	1	1,4	13,13,14	0.98	1 (7%)	17,19,21	1.00	0
4	Z61	D	2	4	10,10,11	0.40	0	13,13,15	1.14	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	3/6/23/26	0/1/1/1
3	FUC	C	2	3	-	-	0/1/1/1
4	8I4	D	1	1,4	-	0/4/24/28	0/1/1/1
4	Z61	D	2	4	-	0/2/16/18	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	1	8I4	O6B-C6	-3.14	1.20	1.30

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	NAG	C1-C2-N2	-5.11	101.75	110.49
3	C	1	NAG	C2-N2-C7	4.16	128.82	122.90
3	C	1	NAG	C8-C7-N2	3.42	121.89	116.10
4	D	2	Z61	C3-C4-C5	-3.20	106.78	109.97
3	C	2	FUC	C1-O5-C5	2.68	118.86	112.78
3	C	2	FUC	O3-C3-C4	-2.31	105.00	110.35
3	C	1	NAG	O7-C7-N2	-2.21	117.88	121.95

There are no chirality outliers.


All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C8-C7-N2-C2
3	C	1	NAG	O7-C7-N2-C2
3	C	1	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
6	EDO	A	606	-	3,3,3	0.36	0	2,2,2	0.18	0
5	NAG	A	604	1	14,14,15	0.66	0	17,19,21	1.15	1 (5%)
6	EDO	A	607	-	3,3,3	0.59	0	2,2,2	0.81	0
5	NAG	A	602	1	14,14,15	1.05	1 (7%)	17,19,21	1.63	5 (29%)
5	NAG	A	601	1	14,14,15	0.72	1 (7%)	17,19,21	1.25	1 (5%)
5	NAG	A	603	1	14,14,15	0.69	0	17,19,21	2.26	4 (23%)
6	EDO	A	605	-	3,3,3	0.45	0	2,2,2	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	A	606	-	-	1/1/1/1	-
5	NAG	A	604	1	-	0/6/23/26	0/1/1/1
6	EDO	A	607	-	-	0/1/1/1	-
5	NAG	A	602	1	-	2/6/23/26	0/1/1/1
5	NAG	A	601	1	-	2/6/23/26	0/1/1/1
5	NAG	A	603	1	-	0/6/23/26	0/1/1/1
6	EDO	A	605	-	-	0/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	602	NAG	C1-C2	2.95	1.56	1.52
5	A	601	NAG	C1-C2	2.27	1.55	1.52

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	603	NAG	C1-O5-C5	7.37	122.17	112.19
5	A	601	NAG	C1-O5-C5	3.73	117.25	112.19
5	A	603	NAG	C6-C5-C4	-3.69	104.36	113.00
5	A	602	NAG	O5-C5-C6	3.59	112.83	107.20
5	A	604	NAG	C1-C2-N2	-2.57	106.10	110.49
5	A	603	NAG	O5-C5-C4	2.40	116.67	110.83
5	A	602	NAG	O3-C3-C2	2.39	114.40	109.47
5	A	602	NAG	C1-O5-C5	2.15	115.10	112.19
5	A	602	NAG	C3-C4-C5	-2.09	106.51	110.24
5	A	602	NAG	C2-N2-C7	2.04	125.81	122.90
5	A	603	NAG	C3-C4-C5	2.02	113.84	110.24

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	601	NAG	O5-C5-C6-O6
5	A	602	NAG	O5-C5-C6-O6
5	A	601	NAG	C4-C5-C6-O6
5	A	602	NAG	C4-C5-C6-O6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
6	A	606	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	606	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	384/389 (98%)	0.47	23 (5%) 29 30	14, 37, 61, 96	4 (1%)
2	B	74/74 (100%)	0.55	3 (4%) 42 45	17, 39, 56, 83	1 (1%)
All	All	458/463 (98%)	0.49	26 (5%) 30 32	14, 38, 62, 96	5 (1%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	160	PHE	5.5
1	A	543	ILE	3.6
1	A	258	PHE	3.5
1	A	298[A]	TYR	3.3
1	A	429	ARG	3.3
1	A	197	LEU	3.3
1	A	427	LYS	3.3
1	A	425	GLY	3.2
1	A	195	ALA	3.2
1	A	303	ARG	3.1
1	A	428	ARG	3.0
1	A	426	SER	2.7
1	A	512	MET	2.5
1	A	440	THR	2.4
2	B	66	PHE	2.4
1	A	318	ILE	2.4
1	A	430[A]	LYS	2.3
1	A	237	ILE	2.3
1	A	498	LEU	2.3
1	A	206	LEU	2.2
1	A	468	TYR	2.2
1	A	272	ARG	2.1
2	B	3	VAL	2.1
2	B	4	VAL	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	232	LYS	2.0
1	A	472	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	C	1	14/15	0.46	0.19	76,83,92,93	0
3	FUC	C	2	10/11	0.79	0.16	61,71,79,80	0
4	Z61	D	2	10/11	0.92	0.09	33,36,44,47	0
4	8I4	D	1	13/14	0.96	0.06	28,31,34,35	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	NAG	A	601	14/15	0.59	0.16	82,89,97,105	0
5	NAG	A	603	14/15	0.60	0.17	74,87,91,93	0
5	NAG	A	602	14/15	0.79	0.13	55,68,71,75	0
5	NAG	A	604	14/15	0.79	0.12	56,67,71,75	0
6	EDO	A	607	4/4	0.88	0.13	38,41,43,56	0
6	EDO	A	606	4/4	0.90	0.19	36,37,38,41	0
6	EDO	A	605	4/4	0.92	0.11	38,39,41,43	0
7	CL	A	608	1/1	0.97	0.05	38,38,38,38	0
7	CL	A	609	1/1	0.97	0.07	48,48,48,48	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.