



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 23, 2025 – 11:30 pm BST

PDB ID : 7R0Y / pdb_00007r0y
Title : ATAD2 in complex with PepLite-Glu
Authors : Turberville, S.; Martin, M.P.; Hope, I.; Noble, M.E.M.
Deposited on : 2022-02-02
Resolution : 1.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2809 (1.46-1.42)
Clashscore	180529	3008 (1.46-1.42)
Ramachandran outliers	177936	2971 (1.46-1.42)
Sidechain outliers	177891	2971 (1.46-1.42)
RSRZ outliers	164620	2809 (1.46-1.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	130	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	HNU	AAA	1203	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 1392 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATPase family AAA domain-containing protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	130	Total	C	N	O	S	0	5	0
			1127	704	199	219	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	979	SER	-	expression tag	UNP Q6PL18
AAA	980	MET	-	expression tag	UNP Q6PL18

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	AAA	1	Total	O	S	0	0
			5	4	1		
2	AAA	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	AAA	1	Total Cl 1 1	0	0

- Molecule 4 is (2 {S})-2-acetamido- {N}-prop-2-enyl-pentanediamide (CCD ID: HNU) (formula: C₁₀H₁₇N₃O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	AAA	1	Total C N O 16 10 3 3	0	0

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	AAA	1	Total	C	O	0	0
			4	2	2		
5	AAA	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	AAA	230	Total	O	0	0
			230	230		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ATPase family AAA domain-containing protein 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	79.29Å 79.29Å 138.01Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	61.48 – 1.43 61.48 – 1.43	Depositor EDS
% Data completeness (in resolution range)	99.3 (61.48-1.43) 99.3 (61.48-1.43)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 1.43Å)	Xtriage
Refinement program	REFMAC 5.8.0258, REFMAC 5.8.0258	Depositor
R, R_{free}	0.195 , 0.213 0.195 , 0.213	Depositor DCC
R_{free} test set	2335 reflections (4.71%)	wwPDB-VP
Wilson B-factor (Å ²)	22.0	Xtriage
Anisotropy	0.107	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 51.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	1392	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, HNU, EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.16	2/1145 (0.2%)	1.61	11/1545 (0.7%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	1078	ALA	C-O	5.91	1.31	1.24
1	AAA	998	HIS	CE1-NE2	5.81	1.38	1.32

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	1076	HIS	CA-C-O	-6.92	113.08	120.42
1	AAA	1078	ALA	CA-C-N	6.19	128.57	120.28
1	AAA	1078	ALA	C-N-CA	6.19	128.57	120.28
1	AAA	1067	ARG	NE-CZ-NH1	-6.02	115.48	121.50
1	AAA	1060	ALA	CA-C-N	5.96	128.27	120.28
1	AAA	1060	ALA	C-N-CA	5.96	128.27	120.28
1	AAA	1059	ASN	OD1-CG-ND2	5.86	128.46	122.60
1	AAA	994	ARG	CG-CD-NE	-5.77	99.31	112.00
1	AAA	1041	HIS	ND1-CE1-NE2	5.29	113.69	108.40
1	AAA	1102	GLU	CB-CG-CD	5.21	121.45	112.60
1	AAA	1078	ALA	CA-C-O	-5.13	114.98	120.42

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1127	0	1112	24	7
2	AAA	10	0	0	0	0
3	AAA	1	0	0	1	0
4	AAA	16	0	0	6	5
5	AAA	8	0	12	3	0
6	AAA	230	0	0	15	8
All	All	1392	0	1124	28	13

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:1075:ARG:HB2	6:AAA:1313:HOH:O	1.44	1.14
4:AAA:1203:HNU:C3	6:AAA:1321:HOH:O	2.13	0.95
1:AAA:1027[B]:GLN:OE1	6:AAA:1302:HOH:O	1.87	0.91
1:AAA:995:ASN:ND2	6:AAA:1303:HOH:O	2.05	0.89
1:AAA:998:HIS:HD2	5:AAA:1205:EDO:H11	1.35	0.89
1:AAA:998:HIS:CD2	5:AAA:1205:EDO:H11	2.14	0.82
1:AAA:981[A]:GLN:HE21	1:AAA:981[A]:GLN:H	1.29	0.80
1:AAA:1090:LYS:HE2	6:AAA:1485:HOH:O	1.86	0.75
1:AAA:1066:ASP:OD1	6:AAA:1304:HOH:O	2.07	0.70
1:AAA:983:GLU:OE2	6:AAA:1305:HOH:O	2.10	0.70
3:AAA:1202:CL:CL	6:AAA:1431:HOH:O	2.51	0.65
1:AAA:1075:ARG:HH21	4:AAA:1203:HNU:C7	2.10	0.64
1:AAA:981[A]:GLN:NE2	6:AAA:1311:HOH:O	2.30	0.61
1:AAA:1020:ASP:OD2	6:AAA:1306:HOH:O	2.16	0.61
4:AAA:1203:HNU:C5	6:AAA:1313:HOH:O	2.49	0.60
1:AAA:995:ASN:ND2	6:AAA:1314:HOH:O	2.36	0.57
1:AAA:987[A]:ARG:NH2	6:AAA:1315:HOH:O	2.38	0.55
1:AAA:1075:ARG:HH21	4:AAA:1203:HNU:C8	2.19	0.55
1:AAA:1081[B]:LEU:C	1:AAA:1081[B]:LEU:HD23	2.34	0.53
1:AAA:1082:ARG:HB3	1:AAA:1082:ARG:HH11	1.74	0.52
1:AAA:980:MET:HE3	1:AAA:981[A]:GLN:HG3	1.91	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:1030:ASP:OD2	1:AAA:1032:SER:OG	2.24	0.51
1:AAA:998:HIS:HD2	5:AAA:1205:EDO:C1	2.16	0.50
1:AAA:1075:ARG:NH2	4:AAA:1203:HNU:C7	2.76	0.48
4:AAA:1203:HNU:C6	4:AAA:1203:HNU:C10	2.92	0.47
1:AAA:987[B]:ARG:NH2	6:AAA:1319:HOH:O	2.43	0.46
1:AAA:1028:PRO:HD2	6:AAA:1493:HOH:O	2.16	0.44
1:AAA:981[A]:GLN:HE21	1:AAA:981[A]:GLN:N	2.06	0.43

All (13) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:981[B]:GLN:NE2	4:AAA:1203:HNU:O1[8_665]	1.18	1.02
1:AAA:981[B]:GLN:NE2	4:AAA:1203:HNU:C4[8_665]	1.49	0.71
1:AAA:981[B]:GLN:NE2	4:AAA:1203:HNU:N3[8_665]	1.70	0.50
6:AAA:1315:HOH:O	6:AAA:1328:HOH:O[6_664]	1.71	0.49
1:AAA:981[B]:GLN:NE2	4:AAA:1203:HNU:C5[8_665]	1.76	0.44
1:AAA:981[B]:GLN:CD	4:AAA:1203:HNU:O1[8_665]	1.78	0.42
6:AAA:1460:HOH:O	6:AAA:1475:HOH:O[12_564]	1.78	0.42
6:AAA:1381:HOH:O	6:AAA:1456:HOH:O[12_564]	1.82	0.38
1:AAA:981[A]:GLN:O	6:AAA:1301:HOH:O[8_665]	2.06	0.14
6:AAA:1462:HOH:O	6:AAA:1509:HOH:O[10_665]	2.06	0.14
6:AAA:1383:HOH:O	6:AAA:1461:HOH:O[8_565]	2.07	0.13
1:AAA:985:THR:OG1	6:AAA:1301:HOH:O[8_665]	2.14	0.06
6:AAA:1328:HOH:O	6:AAA:1392:HOH:O[8_565]	2.16	0.04

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	133/130 (102%)	132 (99%)	1 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	AAA	128/123 (104%)	123 (96%)	5 (4%)	27 3

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	980	MET
1	AAA	981[A]	GLN
1	AAA	981[B]	GLN
1	AAA	1067	ARG
1	AAA	1082	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	AAA	1206	-	4,4,4	0.37	0	6,6,6	0.08	0
5	EDO	AAA	1204	-	3,3,3	0.63	0	2,2,2	0.46	0
4	HNU	AAA	1203	-	15,15,15	0.91	1 (6%)	18,18,18	2.73	4 (22%)
5	EDO	AAA	1205	-	3,3,3	0.27	0	2,2,2	0.20	0
2	SO4	AAA	1201	-	4,4,4	0.82	0	6,6,6	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HNU	AAA	1203	-	-	7/17/17/17	-
5	EDO	AAA	1205	-	-	1/1/1/1	-
5	EDO	AAA	1204	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	AAA	1203	HNU	C3-N1	-2.28	1.39	1.46

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	AAA	1203	HNU	C5-C4-N1	7.08	130.81	116.54
4	AAA	1203	HNU	O1-C4-N1	-6.94	108.09	122.99
4	AAA	1203	HNU	C2-C3-N1	5.02	132.94	112.92
4	AAA	1203	HNU	C5-N3-C9	2.11	127.75	121.48

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	AAA	1203	HNU	C6-C5-N3-C9
4	AAA	1203	HNU	C10-C9-N3-C5
4	AAA	1203	HNU	O3-C9-N3-C5
4	AAA	1203	HNU	C5-C4-N1-C3
4	AAA	1203	HNU	O1-C4-N1-C3
4	AAA	1203	HNU	C5-C6-C7-C8
4	AAA	1203	HNU	C1-C2-C3-N1
5	AAA	1205	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	AAA	1203	HNU	6	5
5	AAA	1205	EDO	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	130/130 (100%)	0.70	16 (12%) 9 8	9, 25, 62, 101	5 (3%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	980	MET	4.5
1	AAA	1015	PRO	4.2
1	AAA	1022	VAL	4.1
1	AAA	1019	PRO	3.9
1	AAA	1018	VAL	3.9
1	AAA	1024	VAL	3.1
1	AAA	1023	THR	2.9
1	AAA	1079	CYS	2.7
1	AAA	1014	ASP	2.6
1	AAA	1108	ARG	2.5
1	AAA	981[A]	GLN	2.5
1	AAA	1107[A]	SER	2.5
1	AAA	1016	ASP	2.4
1	AAA	1017	GLU	2.1
1	AAA	1105	GLN	2.0
1	AAA	1068	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	HNU	AAA	1203	16/16	0.58	0.29	31,68,80,99	0
5	EDO	AAA	1205	4/4	0.71	0.28	54,62,63,66	0
2	SO4	AAA	1206	5/5	0.75	0.18	102,107,115,116	5
3	CL	AAA	1202	1/1	0.93	0.11	67,67,67,67	0
5	EDO	AAA	1204	4/4	0.97	0.06	20,22,28,28	0
2	SO4	AAA	1201	5/5	0.97	0.08	29,31,34,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.