Full wwPDB X-ray Structure Validation Report (i)

Mar 7, 2026 – 01:45 AM UTC

PDB ID	:	7RVX / pdb_00007rvx
Title	:	Structure of the SARS-CoV-2 main protease in complex with inhibitor MPI24
Authors	:	Yang, K.; Liu, W.
Deposited on	:	2021-08-19
Resolution	:	1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	13% 77% 20% .
1	B	306	19% 79% 20% .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5069 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3C-like proteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	306	Total	C	N	O	S	0	2	0
			2384	1507	405	450	22			
1	B	306	Total	C	N	O	S	0	2	0
			2384	1507	405	450	22			

- Molecule 2 is benzyl [(1S)-1-cyclopropyl-2-{[(2S)-3-cyclopropyl-1-({(2S)-1-hydroxy-3-[(3S)-2-oxopyrrolidin-3-yl]propan-2-yl}amino)-1-oxopropan-2-yl]amino}-2-oxoethyl]carbamate (CCD ID: 7YI) (formula: $C_{26}H_{36}N_4O_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			36	26	4	6		
2	B	1	Total	C	N	O	0	0
			36	26	4	6		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	127	Total O 127 127	0	0
3	B	102	Total O 102 102	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 3C-like proteinase

Chain A:



• Molecule 1: 3C-like proteinase

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	56.06Å 61.15Å 63.76Å 79.94° 68.33° 70.17°	Depositor
Resolution (Å)	49.69 – 1.85 49.69 – 1.85	Depositor EDS
% Data completeness (in resolution range)	88.9 (49.69-1.85) 88.9 (49.69-1.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.91 (at 1.86Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.235 , 0.265 0.239 , 0.271	Depositor DCC
R_{free} test set	2799 reflections (4.43%)	wwPDB-VP
Wilson B-factor (Å^2)	30.7	Xtriage
Anisotropy	0.185	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 39.0	EDS
L-test for twinning[2]	$< \|L\| > = 0.52$, $< L^2 > = 0.36$	Xtriage
Estimated twinning fraction	0.147 for -h,-l,-k	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	5069	wwPDB-VP
Average B, all atoms (Å^2)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.71% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 7YI, CSO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.46	0/2421	0.80	6/3287 (0.2%)
1	B	0.57	3/2421 (0.1%)	0.89	12/3287 (0.4%)
All	All	0.52	3/4842 (0.1%)	0.84	18/6574 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	188	ARG	CB-CG	-7.70	1.29	1.52
1	B	188	ARG	CG-CD	-6.47	1.33	1.52
1	B	60	ARG	CD-NE	-5.54	1.38	1.46

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	60	ARG	CD-NE-CZ	10.87	139.62	124.40
1	A	221	ASN	CA-C-N	6.84	129.77	120.54
1	A	221	ASN	C-N-CA	6.84	129.77	120.54
1	B	60	ARG	CA-CB-CG	6.70	127.51	114.10
1	A	5	LYS	N-CA-CB	6.69	120.57	109.60
1	B	188	ARG	CA-CB-CG	-6.43	101.23	114.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	222	ARG	CG-CD-NE	-6.30	98.13	112.00
1	B	188	ARG	CG-CD-NE	-6.18	98.39	112.00
1	A	298	ARG	CB-CG-CD	6.09	125.32	111.30
1	A	49	MET	CB-CG-SD	5.95	130.55	112.70
1	B	60	ARG	NE-CZ-NH1	-5.61	115.89	121.50
1	B	49	MET	CB-CG-SD	5.45	129.05	112.70
1	B	69	GLN	N-CA-C	5.41	116.95	109.15
1	B	100	LYS	CA-CB-CG	-5.29	103.53	114.10
1	B	74	GLN	CA-CB-CG	-5.21	103.68	114.10
1	B	69	GLN	CA-C-N	5.12	129.97	122.85
1	B	69	GLN	C-N-CA	5.12	129.97	122.85
1	A	222	ARG	CB-CG-CD	-5.05	99.69	111.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	168	PRO	Peptide
1	A	300	CSO	Mainchain
1	B	300	CSO	Mainchain
1	B	60	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2384	0	2323	54	1
1	B	2384	0	2323	44	2
2	A	36	0	0	0	0
2	B	36	0	0	0	0
3	A	127	0	0	14	0
3	B	102	0	0	7	0
All	All	5069	0	4646	92	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:ALA:O	3:A:501:HOH:O	1.83	0.96
1:A:90:LYS:NZ	3:A:504:HOH:O	2.07	0.87
1:A:206:ALA:HB3	3:A:511:HOH:O	1.74	0.87
1:A:3:PHE:O	3:A:502:HOH:O	1.93	0.85
1:B:194:ALA:O	3:B:501:HOH:O	1.94	0.84
1:A:245:ASP:O	1:A:249:ILE:HD12	1.87	0.75
1:A:222:ARG:HD3	1:A:222:ARG:H	1.50	0.74
1:A:270:GLU:OE2	3:A:503:HOH:O	2.06	0.74
1:A:22:CYS:SG	1:A:61:LYS:NZ	2.61	0.74
1:B:245:ASP:O	1:B:249:ILE:HD12	1.90	0.71
1:A:203:ASN:HA	3:A:511:HOH:O	1.91	0.70
1:A:168:PRO:O	3:A:505:HOH:O	2.11	0.68
1:B:51:ASN:OD1	1:B:188:ARG:NH2	2.27	0.68
1:B:86:VAL:HG13	1:B:179:GLY:HA2	1.75	0.68
1:A:53:ASN:O	1:A:56:ASP:HB2	1.93	0.68
1:A:61:LYS:NZ	3:A:512:HOH:O	2.28	0.67
1:B:113:SER:OG	1:B:127:GLN:OE1	2.08	0.67
1:B:270:GLU:OE1	3:B:504:HOH:O	2.12	0.67
1:B:226:THR:HG23	1:B:229:ASP:H	1.60	0.67
1:A:100:LYS:NZ	3:A:513:HOH:O	2.28	0.66
1:B:40:ARG:HD3	1:B:85:CYS:HA	1.77	0.65
1:A:198:THR:HG22	1:A:238:ASN:OD1	1.97	0.65
1:A:169:THR:HG21	3:A:621:HOH:O	1.96	0.64
1:B:57:LEU:O	1:B:60:ARG:HB2	1.98	0.64
1:A:71:GLY:HA3	3:A:507:HOH:O	1.97	0.63
1:A:222:ARG:H	1:A:222:ARG:CD	2.10	0.63
1:B:198:THR:OG1	1:B:240:GLU:OE1	2.07	0.62
1:A:222:ARG:HD3	1:A:222:ARG:N	2.14	0.62
1:A:226:THR:HG23	1:A:229:ASP:H	1.64	0.61
1:B:54:TYR:HB3	1:B:82:MET:HE1	1.83	0.60
1:A:128:CSO:HA	3:A:506:HOH:O	2.01	0.60
1:B:139:SER:OG	3:B:503:HOH:O	2.11	0.59
1:B:277:ASN:OD1	3:B:505:HOH:O	2.16	0.58
1:B:59:ILE:HG13	1:B:60:ARG:N	2.18	0.58
1:B:225:THR:O	1:B:262:LEU:HD13	2.03	0.58
1:A:12:LYS:NZ	1:A:306:GLN:OE1	2.37	0.56
1:A:233:VAL:HG11	1:A:269:LYS:HG3	1.88	0.55
1:B:53:ASN:O	1:B:56:ASP:HB2	2.06	0.55
1:A:266:ALA:O	1:A:270:GLU:HG2	2.06	0.55
1:B:288:GLU:CD	3:B:513:HOH:O	2.50	0.55
1:A:163:HIS:HE1	1:A:172:HIS:HB3	1.72	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:298:ARG:HG3	1:A:299:GLN:N	2.22	0.54
1:A:86:VAL:HG13	1:A:179:GLY:HA2	1.91	0.53
1:A:5:LYS:NZ	1:B:4:ARG:HH22	2.06	0.53
1:A:169:THR:HG22	1:A:171:VAL:HG13	1.91	0.53
1:B:27:LEU:HD13	1:B:39:PRO:HD2	1.91	0.52
1:B:163:HIS:HE1	1:B:172:HIS:HB3	1.75	0.52
1:B:198:THR:HG22	1:B:238:ASN:OD1	2.09	0.52
1:A:226:THR:CG2	1:A:229:ASP:H	2.24	0.51
1:A:40:ARG:O	1:A:43:ILE:HG12	2.11	0.51
1:B:58:LEU:HD22	1:B:82:MET:HE3	1.94	0.49
1:A:163:HIS:CE1	1:A:172:HIS:HB3	2.48	0.49
1:A:27:LEU:HD13	1:A:39:PRO:HD2	1.96	0.48
1:B:221:ASN:OD1	1:B:223:PHE:HB2	2.14	0.47
1:A:1:SER:HB2	1:A:214:ASN:ND2	2.30	0.47
1:A:45:THR:O	1:A:49:MET:HG2	2.14	0.47
1:B:163:HIS:CE1	1:B:172:HIS:HB3	2.50	0.47
1:A:285:ALA:HB2	1:B:286:LEU:HD21	1.97	0.46
1:A:115:LEU:HD11	1:A:122:PRO:HB3	1.98	0.46
1:A:123:SER:HB2	1:B:304:THR:HG22	1.98	0.46
1:A:72:ASN:N	3:A:507:HOH:O	2.21	0.45
1:B:40:ARG:CD	1:B:85:CYS:HA	2.45	0.45
1:A:218:TRP:HH2	1:A:276:MET:HG2	1.80	0.45
1:B:56:ASP:O	1:B:60:ARG:HG2	2.16	0.45
1:A:57:LEU:O	1:A:60:ARG:HB2	2.18	0.44
1:A:31:TRP:CE2	1:A:95:ASN:HB2	2.53	0.44
1:A:5:LYS:HG2	1:A:291:PHE:CZ	2.53	0.43
1:A:5:LYS:HZ2	1:B:4:ARG:HH22	1.67	0.43
1:A:153:ASP:O	1:A:155:ASP:N	2.51	0.43
1:A:298:ARG:CG	1:A:299:GLN:N	2.81	0.43
1:B:111:THR:HG22	1:B:129:ALA:HB2	2.01	0.43
1:B:52:PRO:O	1:B:188:ARG:NH1	2.52	0.42
1:B:222:ARG:NH2	1:B:223:PHE:HE1	2.18	0.42
1:B:45:THR:H	1:B:48:ASP:HB2	1.85	0.41
1:B:57:LEU:HA	1:B:60:ARG:CG	2.50	0.41
1:B:226:THR:CG2	1:B:229:ASP:HB2	2.50	0.41
1:A:27:LEU:HD21	1:A:42:VAL:HB	2.02	0.41
1:A:221:ASN:HB2	1:A:222:ARG:CZ	2.51	0.41
1:B:75:LEU:O	3:B:507:HOH:O	2.22	0.41
1:B:113:SER:O	1:B:149:GLY:HA2	2.21	0.41
1:B:40:ARG:O	1:B:43:ILE:HG12	2.21	0.41
1:B:167:LEU:HB3	1:B:168:PRO:HD2	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:277:ASN:N	3:B:506:HOH:O	2.53	0.41
1:A:31:TRP:CZ3	1:A:75:LEU:HD11	2.56	0.40
1:A:252:PRO:HB2	3:A:529:HOH:O	2.21	0.40
1:A:285:ALA:HB3	1:B:285:ALA:HB3	2.02	0.40
1:A:10:SER:O	1:A:14:GLU:HG3	2.21	0.40
1:A:1:SER:HB2	1:A:214:ASN:HD22	1.86	0.40
1:A:109:GLY:HA2	1:A:200:ILE:HD13	2.04	0.40
1:A:286:LEU:HD11	1:B:280:THR:HG21	2.02	0.40
1:B:190:THR:O	1:B:192:GLN:HG3	2.22	0.40
1:B:282:LEU:HD23	1:B:282:LEU:HA	1.89	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:60:ARG:NH2	1:B:220:LEU:O[1_654]	1.78	0.42
1:A:60:ARG:NH1	1:A:220:LEU:O[1_465]	1.93	0.27
1:B:60:ARG:NH1	1:B:217:ARG:O[1_654]	2.08	0.12

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	304/306 (99%)	293 (96%)	8 (3%)	3 (1%)	12 4
1	B	304/306 (99%)	290 (95%)	13 (4%)	1 (0%)	36 25
All	All	608/612 (99%)	583 (96%)	21 (4%)	4 (1%)	18 8

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	154	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	154	TYR
1	A	49	MET
1	A	169	THR

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	263/261 (101%)	263 (100%)	0	100 100
1	B	263/261 (101%)	261 (99%)	2 (1%)	73 67
All	All	526/522 (101%)	524 (100%)	2 (0%)	87 82

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	46[A]	SER
1	B	46[B]	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	GLN
1	A	119	ASN
1	A	189	GLN
1	A	214	ASN
1	A	256	GLN
1	B	53	ASN
1	B	164	HIS
1	B	274	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	CSO	B	300	1	3,6,7	0.68	0	1,6,8	0.39	0
1	CSO	A	128	1	3,6,7	1.06	0	1,6,8	1.30	0
1	CSO	A	300	1	3,6,7	0.77	0	1,6,8	0.02	0
1	CSO	B	128	1	3,6,7	0.98	0	1,6,8	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	B	300	1	-	0/1/5/7	-
1	CSO	A	128	1	-	0/1/5/7	-
1	CSO	A	300	1	-	0/1/5/7	-
1	CSO	B	128	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	128	CSO	1	0

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	7YI	B	401	1	39,39,39	1.04	1 (2%)	46,53,53	1.90	6 (13%)
2	7YI	A	401	1	39,39,39	1.01	1 (2%)	46,53,53	1.98	7 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	7YI	B	401	1	-	0/39/53/53	0/4/4/4
2	7YI	A	401	1	-	1/39/53/53	0/4/4/4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	7YI	O8-C9	5.18	1.45	1.35
2	A	401	7YI	O8-C9	5.01	1.45	1.35

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	401	7YI	C34-C8-C28	-9.99	105.82	119.71
2	B	401	7YI	C34-C8-C28	-9.24	106.86	119.71
2	A	401	7YI	C34-C8-C29	-4.77	113.08	119.71
2	B	401	7YI	C8-C34-C14	-4.26	108.80	114.52
2	A	401	7YI	C8-C34-C14	-4.10	109.01	114.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	401	7YI	C34-C8-C29	-3.98	114.17	119.71
2	B	401	7YI	O26-C24-C20	-3.29	122.40	126.21
2	B	401	7YI	O8-C9-N10	2.74	116.32	110.45
2	B	401	7YI	O8-C9-O29	-2.57	119.34	124.26
2	A	401	7YI	O26-C24-C20	-2.52	123.29	126.21
2	A	401	7YI	O8-C9-N10	2.30	115.37	110.45
2	A	401	7YI	C7-O8-C9	2.29	121.07	115.93
2	A	401	7YI	O8-C9-O29	-2.26	119.92	124.26

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	7YI	N10-C11-C12-O33

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 7YI B 401



Bond lengths



Bond angles



Torsions



Rings

Ligand 7YI A 401



Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	304/306 (99%)	1.14	41 (13%)	7	6	15, 37, 59, 90	2 (0%)
1	B	304/306 (99%)	1.27	57 (18%)	3	3	16, 38, 65, 93	2 (0%)
All	All	608/612 (99%)	1.20	98 (16%)	4	4	15, 37, 63, 93	4 (0%)

All (98) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	154	TYR	9.2
1	A	154	TYR	7.5
1	B	60	ARG	6.5
1	B	50	LEU	6.1
1	A	50	LEU	5.9
1	A	49	MET	5.4
1	A	46[A]	SER	5.0
1	B	49	MET	5.0
1	A	72	ASN	4.8
1	B	45	THR	4.7
1	A	298	ARG	4.7
1	A	45	THR	4.6
1	B	303	VAL	4.5
1	B	223	PHE	4.2
1	B	222	ARG	4.1
1	B	74	GLN	4.1
1	B	72	ASN	4.0
1	B	235	MET	4.0
1	B	193	ALA	3.9
1	A	74	GLN	3.9
1	B	68	VAL	3.9
1	B	228	ASN	3.9
1	B	306	GLN	3.9
1	B	304	THR	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	44	CYS	3.8
1	A	56	ASP	3.7
1	B	305	PHE	3.7
1	A	153	ASP	3.7
1	A	44	CYS	3.6
1	B	59	ILE	3.6
1	A	222	ARG	3.6
1	A	73	VAL	3.5
1	A	303	VAL	3.4
1	B	224	THR	3.4
1	B	294	PHE	3.4
1	B	226	THR	3.3
1	A	294	PHE	3.3
1	B	48	ASP	3.3
1	B	57	LEU	3.2
1	A	306	GLN	3.2
1	B	46[A]	SER	3.2
1	A	194	ALA	3.1
1	B	56	ASP	3.0
1	B	232	LEU	3.0
1	A	51	ASN	3.0
1	B	277	ASN	3.0
1	A	76	ARG	3.0
1	A	305	PHE	2.9
1	A	59	ILE	2.9
1	A	47	GLU	2.9
1	B	52	PRO	2.8
1	A	168	PRO	2.8
1	A	226	THR	2.7
1	B	51	ASN	2.7
1	B	55	GLU	2.7
1	A	195	GLY	2.7
1	B	43	ILE	2.7
1	A	104	VAL	2.6
1	B	188	ARG	2.6
1	A	48	ASP	2.5
1	A	232	LEU	2.5
1	A	213	ILE	2.5
1	B	249	ILE	2.5
1	B	104	VAL	2.5
1	B	71	GLY	2.4
1	B	301	SER	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	155	ASP	2.4
1	A	252	PRO	2.4
1	A	304	THR	2.3
1	A	57	LEU	2.3
1	B	181	PHE	2.3
1	A	53	ASN	2.3
1	B	262	LEU	2.3
1	B	73	VAL	2.2
1	A	122	PRO	2.2
1	B	194	ALA	2.2
1	B	245	ASP	2.2
1	A	43	ILE	2.2
1	A	55	GLU	2.2
1	B	169	THR	2.2
1	B	242	LEU	2.2
1	B	53	ASN	2.1
1	A	67	LEU	2.1
1	A	255	ALA	2.1
1	B	86	VAL	2.1
1	B	244	GLN	2.1
1	B	106	ILE	2.1
1	A	60	ARG	2.1
1	B	26	THR	2.1
1	B	47	GLU	2.1
1	B	236	LYS	2.1
1	B	278	GLY	2.1
1	A	142	ASN	2.1
1	B	69	GLN	2.0
1	B	184	PRO	2.0
1	B	67	LEU	2.0
1	B	152	ILE	2.0
1	B	298	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
1	CSO	B	128	7/8	0.88	0.12	23,27,35,42	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
1	CSO	B	300	7/8	0.88	0.14	31,35,44,45	0
1	CSO	A	128	7/8	0.89	0.12	22,24,31,43	0
1	CSO	A	300	7/8	0.89	0.14	32,38,44,48	0

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	7YI	B	401	36/36	0.89	0.12	31,39,59,60	0
2	7YI	A	401	36/36	0.91	0.10	29,38,51,55	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 7YI B 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around 7YI A 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.