



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 24, 2026 – 03:34 pm GMT

PDB ID : 8AKD / pdb_00008akd
Title : Human Sirt6 in complex with ADP-ribose and fragment 1-methyl-L-histidine
Authors : You, W.; Steegborn, C.
Deposited on : 2022-07-29
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.48.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2888 (1.76-1.76)
Clashscore	180529	3097 (1.76-1.76)
Ramachandran outliers	177936	3072 (1.76-1.76)
Sidechain outliers	177891	3072 (1.76-1.76)
RSRZ outliers	164620	2887 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	85% 8% 7%
1	B	302	83% 8% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	SO4	B	408	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4658 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NAD-dependent protein deacetylase sirtuin-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	282	Total	C	N	O	S	0	0	0
			2206	1388	404	403	11			
1	B	277	Total	C	N	O	S	0	0	0
			2158	1358	396	393	11			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	GLY	-	expression tag	UNP Q8N6T7
A	8	ILE	-	expression tag	UNP Q8N6T7
A	9	ASP	-	expression tag	UNP Q8N6T7
A	10	PRO	-	expression tag	UNP Q8N6T7
A	11	PHE	-	expression tag	UNP Q8N6T7
A	12	THR	-	expression tag	UNP Q8N6T7
B	7	GLY	-	expression tag	UNP Q8N6T7
B	8	ILE	-	expression tag	UNP Q8N6T7
B	9	ASP	-	expression tag	UNP Q8N6T7
B	10	PRO	-	expression tag	UNP Q8N6T7
B	11	PHE	-	expression tag	UNP Q8N6T7
B	12	THR	-	expression tag	UNP Q8N6T7

- Molecule 2 is [(2R,3S,4R,5R)-5-(6-AMINOPURIN-9-YL)-3,4-DIHYDROXY-OXOLAN-2-YL]METHYL[HYDROXY-[(2R,3S,4R,5S)-3,4,5-TRIHYDROXYOXOLAN-2-YL]METHOXY]PHOSPHORYL] HYDROGEN PHOSPHATE (CCD ID: AR6) (formula: C₁₅H₂₃N₅O₁₄P₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	36	15	5	14	2	0	0
2	B	1	36	15	5	14	2	0	0

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
3	A	1	1	1	0	0
3	B	1	1	1	0	0

- Molecule 4 is 4-METHYL-HISTIDINE (CCD ID: HIC) (formula: C₇H₁₁N₃O₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			12	7	3	2		
4	B	1	Total	C	N	O	0	0
			12	7	3	2		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	84	Total	O	0	0
			84	84		
7	B	63	Total	O	0	0
			63	63		

4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	91.06Å 91.06Å 143.30Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.53 – 1.76 45.53 – 1.76	Depositor EDS
% Data completeness (in resolution range)	98.9 (45.53-1.76) 98.9 (45.53-1.76)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.23 (at 1.76Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.176 , 0.202 0.179 , 0.202	Depositor DCC
R_{free} test set	2100 reflections (3.19%)	wwPDB-VP
Wilson B-factor (Å ²)	30.4	Xtrriage
Anisotropy	0.532	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 32.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.22$	Xtrriage
Estimated twinning fraction	0.387 for h,-h-k,-l	Xtrriage
Reported twinning fraction	0.602 for H, K, L 0.398 for -K, -H, -L	Depositor
Outliers	0 of 65810 reflections	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4658	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HIC, AR6, PEG, ZN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.11	3/2255 (0.1%)	1.47	4/3058 (0.1%)
1	B	1.06	1/2205 (0.0%)	1.43	4/2991 (0.1%)
All	All	1.08	4/4460 (0.1%)	1.45	8/6049 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	95	HIS	CE1-NE2	6.39	1.39	1.32
1	A	56	SER	CA-CB	-5.14	1.46	1.53
1	A	213	LEU	C-O	5.05	1.29	1.24
1	B	50	HIS	CE1-NE2	-5.03	1.27	1.32

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	108	ARG	CG-CD-NE	-5.79	99.27	112.00
1	A	94	THR	CA-C-O	-5.78	114.94	120.90
1	B	151	ASP	CA-CB-CG	5.74	118.34	112.60
1	B	280	ARG	CG-CD-NE	-5.47	99.97	112.00
1	A	287	PRO	N-CA-C	-5.24	105.44	110.47
1	B	108	ARG	CG-CD-NE	5.19	123.42	112.00
1	A	251	ASP	N-CA-C	-5.17	106.48	112.89
1	B	94	THR	CA-C-O	-5.05	115.70	121.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2206	0	2219	14	0
1	B	2158	0	2171	19	0
2	A	36	0	21	1	0
2	B	36	0	21	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	12	0	10	1	0
4	B	12	0	10	0	0
5	A	7	0	10	0	0
5	B	7	0	10	0	0
6	A	15	0	0	0	0
6	B	20	0	0	2	0
7	A	84	0	0	5	0
7	B	63	0	0	6	0
All	All	4658	0	4472	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:68:HIS:CE1	1:B:73:MET:HE2	2.26	0.70
1:A:73:MET:HE2	1:B:68:HIS:CE1	2.29	0.68
1:B:298:GLU:HB3	7:B:538:HOH:O	1.92	0.67
1:B:253:ARG:HD2	7:B:512:HOH:O	1.95	0.64
1:A:50:HIS:HE1	1:A:210:SER:OG	1.83	0.62
1:A:104:VAL:HB	1:A:291:ARG:HB3	1.83	0.60
1:A:189:GLU:HB2	7:A:579:HOH:O	2.03	0.58
1:B:104:VAL:HB	1:B:291:ARG:HB3	1.87	0.56
1:B:74:GLU:HB2	7:B:546:HOH:O	2.05	0.54
1:B:253:ARG:CD	7:B:512:HOH:O	2.52	0.54
1:B:245:LYS:HB3	1:B:246:HIS:CD2	2.43	0.53
1:B:221:PRO:HB3	6:B:408:SO4:O3	2.09	0.53
1:B:255:HIS:CE1	7:B:512:HOH:O	2.61	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:220:ARG:HB3	6:B:408:SO4:O2	2.12	0.50
1:B:298:GLU:C	7:B:538:HOH:O	2.56	0.48
1:A:63:ASP:OD2	2:A:401:AR6:H8	2.13	0.48
1:B:229:THR:O	1:B:234:GLY:HA3	2.13	0.48
1:B:28:GLU:H	1:B:28:GLU:CD	2.23	0.47
1:B:238:ILE:HD12	1:B:250:ALA:HB2	1.97	0.47
1:A:28:GLU:H	1:A:28:GLU:CD	2.23	0.46
1:B:63:ASP:OD2	2:B:402:AR6:H8	2.17	0.45
1:B:240:ASN:ND2	2:B:402:AR6:O3'	2.46	0.44
1:A:32:ARG:HD2	7:A:570:HOH:O	2.16	0.44
1:A:280:ARG:HH11	1:A:280:ARG:HG2	1.83	0.44
1:B:46:SER:OG	1:B:108:ARG:NH1	2.51	0.44
1:A:238:ILE:HD12	1:A:250:ALA:HB2	2.00	0.42
1:A:50:HIS:HA	1:A:111:VAL:O	2.18	0.42
4:A:403:HIC:HZ1	7:A:578:HOH:O	2.20	0.41
1:A:160:LYS:HE2	7:A:506:HOH:O	2.19	0.41
1:A:227:LEU:HD11	1:A:246:HIS:ND1	2.35	0.41
1:B:108:ARG:HH11	1:B:108:ARG:HG2	1.85	0.41
1:A:50:HIS:HD2	7:A:560:HOH:O	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	278/302 (92%)	273 (98%)	5 (2%)	0	100	100
1	B	273/302 (90%)	268 (98%)	5 (2%)	0	100	100
All	All	551/604 (91%)	541 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	240/256 (94%)	238 (99%)	2 (1%)	79	71
1	B	234/256 (91%)	233 (100%)	1 (0%)	89	86
All	All	474/512 (93%)	471 (99%)	3 (1%)	84	78

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	210	SER
1	A	244	THR
1	B	210	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	HIS
1	A	93	GLN
1	A	255	HIS
1	B	246	HIS
1	B	255	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 15 ligands modelled in this entry, 2 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	HIC	A	403	-	9,12,12	0.98	1 (11%)	10,16,16	1.17	2 (20%)
6	SO4	B	406	-	4,4,4	0.39	0	6,6,6	0.05	0
2	AR6	B	402	-	34,39,39	0.61	0	40,60,60	0.97	1 (2%)
6	SO4	B	407	-	4,4,4	0.38	0	6,6,6	0.09	0
6	SO4	B	408	-	4,4,4	0.37	0	6,6,6	0.06	0
4	HIC	B	404	-	9,12,12	1.01	1 (11%)	10,16,16	1.24	2 (20%)
5	PEG	A	404	-	6,6,6	0.17	0	5,5,5	0.18	0
6	SO4	A	407	-	4,4,4	0.34	0	6,6,6	0.11	0
6	SO4	A	405	-	4,4,4	0.41	0	6,6,6	0.09	0
6	SO4	B	405	-	4,4,4	0.52	0	6,6,6	0.12	0
2	AR6	A	401	-	34,39,39	0.71	0	40,60,60	0.87	0
6	SO4	A	406	-	4,4,4	0.39	0	6,6,6	0.11	0
5	PEG	B	401	-	6,6,6	0.15	0	5,5,5	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HIC	A	403	-	-	4/8/8/8	0/1/1/1
2	AR6	B	402	-	-	2/18/54/54	0/4/4/4
4	HIC	B	404	-	-	4/8/8/8	0/1/1/1
5	PEG	A	404	-	-	2/4/4/4	-
2	AR6	A	401	-	-	2/18/54/54	0/4/4/4
5	PEG	B	401	-	-	1/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	403	HIC	OXT-C	-2.17	1.23	1.30
4	B	404	HIC	OXT-C	-2.10	1.23	1.30

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	404	HIC	OXT-C-O	-2.44	118.55	124.09
2	B	402	AR6	O3'-C3'-C2'	-2.29	104.42	111.82
4	B	404	HIC	OXT-C-CA	2.21	120.92	113.38
4	A	403	HIC	OXT-C-O	-2.20	119.10	124.09
4	A	403	HIC	OXT-C-CA	2.07	120.45	113.38

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	402	AR6	C5D-O5D-PB-O2B
4	A	403	HIC	C-CA-CB-CG
4	B	404	HIC	C-CA-CB-CG
5	B	401	PEG	O2-C3-C4-O4
4	A	403	HIC	OXT-C-CA-N
5	A	404	PEG	O1-C1-C2-O2
5	A	404	PEG	O2-C3-C4-O4
4	A	403	HIC	O-C-CA-N
4	A	403	HIC	N-CA-CB-CG
4	B	404	HIC	N-CA-CB-CG
4	B	404	HIC	CA-CB-CG-ND1
2	B	402	AR6	O4'-C4'-C5'-O5'
2	A	401	AR6	O4'-C4'-C5'-O5'
2	A	401	AR6	C5D-O5D-PB-O2B
4	B	404	HIC	CA-CB-CG-CD2


There are no ring outliers.


4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	403	HIC	1	0
2	B	402	AR6	2	0
6	B	408	SO4	2	0
2	A	401	AR6	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	282/302 (93%)	-0.47	1 (0%) 89 92	24, 37, 60, 77	0
1	B	277/302 (91%)	-0.35	0 100 100	27, 43, 71, 88	0
All	All	559/604 (92%)	-0.41	1 (0%) 92 94	24, 40, 68, 88	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	10	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	PEG	A	404	7/7	0.96	0.09	64,68,71,73	0
4	HIC	B	404	12/12	0.97	0.09	79,84,85,86	0
6	SO4	A	406	5/5	0.97	0.06	61,63,65,70	0
6	SO4	B	407	5/5	0.97	0.05	63,63,65,66	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	SO4	A	405	5/5	0.98	0.05	59,60,62,63	0
4	HIC	A	403	12/12	0.98	0.07	55,62,65,65	0
6	SO4	A	407	5/5	0.98	0.04	53,61,63,66	0
6	SO4	B	406	5/5	0.98	0.05	63,64,69,71	0
5	PEG	B	401	7/7	0.98	0.09	55,56,59,63	0
6	SO4	B	408	5/5	0.98	0.04	61,65,67,68	0
6	SO4	B	405	5/5	0.99	0.04	41,43,47,48	0
2	AR6	B	402	36/36	0.99	0.05	27,35,46,49	0
3	ZN	B	403	1/1	0.99	0.04	61,61,61,61	0
2	AR6	A	401	36/36	0.99	0.05	21,30,35,37	0
3	ZN	A	402	1/1	1.00	0.04	56,56,56,56	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around HIC A 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around AR6 B 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.