

# Full wwPDB X-ray Structure Validation Report (i)

Dec 2, 2025 - 01:18 am GMT

PDB ID : 8B52 / pdb 00008b52

Title : Usutu virus methyltransferase domain in complex with sinefungin Authors : Ferrero, D.S.; Albentosa Gonzalez, L.; Mas, A.; Verdaguer, N.

Deposited on : 2022-09-21

Resolution : 2.22 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 2.0

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.010 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.46

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},\ {\rm resolution\ range}({\rm \AA})) \end{array}$
$R_{free}$	164625	7167 (2.24-2.20)
Clashscore	180529	8096 (2.24-2.20)
Ramachandran outliers	177936	8010 (2.24-2.20)
Sidechain outliers	177891	8011 (2.24-2.20)
RSRZ outliers	164620	7166 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	265	18% 80%	18%	
1	В	265	12% 82%	17%	



# 2 Entry composition (i)


There are 6 unique types of molecules in this entry. The entry contains 4368 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Genome polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	261	Total	С	N	О	S	0	0	0
	1		2072	1295	383	379	15	Ŭ	Ů	
1	B	262	Total	С	N	Ο	S	0	1	0
1	ע	202	2088	1305	385	383	15		1	


 $\bullet$  Molecule 2 is SULFATE ION (CCD ID: SO4) (formula:  $\mathrm{O_4S}).$ 



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	В	1	Total O S 5 4 1	0	0


• Molecule 3 is S-ADENOSYL-L-HOMOCYSTEINE (CCD ID: SAH) (formula:  $C_{14}H_{20}N_6O_5S$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	A	1	Total 26	C 14	N 6	O 5	S 1	0	0


 $\bullet$  Molecule 4 is GLYCEROL (CCD ID: GOL) (formula:  $\mathrm{C_3H_8O_3}).$ 



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0

 $\bullet$  Molecule 5 is S-ADENOSYLMETHIONINE (CCD ID: SAM) (formula:  $C_{15}H_{22}N_6O_5S)$  (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
5	B	1	Total	С	N	О	S	0	0
	D	1	27	15	6	5	1		

#### • Molecule 6 is water.


Mol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
6	A	77	Total O 77 77	0	0
6	В	62	Total O 62 62	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Genome polyprotein





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	53.29Å 95.64Å 110.42Å	Donasiton
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.30 - 2.22	Depositor
Resolution (A)	72.30 - 2.22	EDS
% Data completeness	70.3 (72.30-2.22)	Depositor
(in resolution range)	70.3 (72.30-2.22)	EDS
$R_{merge}$	0.27	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.52 (at 2.22Å)	Xtriage
Refinement program	PHENIX 1.20rc4_4425, PHENIX 1.20rc4_4425	Depositor
D D.	0.266 , 0.276	Depositor
$R, R_{free}$	0.269 , $0.275$	DCC
$R_{free}$ test set	1030 reflections (3.61%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.7	Xtriage
Anisotropy	0.060	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.35, 35.9	EDS
L-test for twinning <sup>2</sup>	$< L >=0.57, < L^2>=0.41$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	4368	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 59.81 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.6706e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SAM, SAH, GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain		nd lengths	Bond angles		
IVIOI		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.73	$4/2116 \ (0.2\%)$	0.70	6/2847 (0.2%)	
1	В	0.39	$2/2132 \ (0.1\%)$	0.46	0/2869	
All	All	0.58	6/4248 (0.1%)	0.59	6/5716 (0.1%)	

#### All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	Observed(A)	$\operatorname{Ideal}( ext{\AA})$
1	A	178	ARG	C-N	-24.62	0.99	1.33
1	A	64	ARG	C-N	12.35	1.50	1.33
1	A	177	PRO	C-N	10.66	1.49	1.33
1	В	46	ARG	C-N	10.27	1.48	1.34
1	A	63	LEU	C-N	8.23	1.44	1.33
1	В	47	ASP	C-N	6.37	1.44	1.33

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\mathbf{Ideal}(^{o})$
1	A	177	PRO	CA-C-N	-13.42	100.92	122.26
1	A	177	PRO	C-N-CA	-13.42	100.92	122.26
1	A	177	PRO	O-C-N	10.67	135.94	123.03
1	A	46	ARG	N-CA-C	-8.92	102.31	113.55
1	A	178	ARG	O-C-N	-8.66	111.70	122.35
1	A	50	LYS	N-CA-C	-5.66	106.22	113.02

There are no chirality outliers.

There are no planarity outliers.



### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2072	0	2061	31	1
1	В	2088	0	2076	43	1
2	A	5	0	0	1	0
2	В	5	0	0	0	0
3	A	26	0	18	0	0
4	A	6	0	8	0	0
5	В	27	0	22	4	0
6	A	77	0	0	8	1
6	В	62	0	0	20	1
All	All	4368	0	4185	75	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:64:ARG:CZ	6:B:401:HOH:O	1.83	1.24
1:B:64:ARG:NH2	6:B:401:HOH:O	1.82	1.07
1:B:42:ARG:NH1	6:B:403:HOH:O	1.89	1.05
1:A:99:GLN:NE2	6:A:403:HOH:O	1.94	1.00
1:B:150:GLU:OE2	6:B:402:HOH:O	1.84	0.96
2:A:301:SO4:O2	6:A:401:HOH:O	1.86	0.93
1:B:259:VAL:CG2	6:B:401:HOH:O	2.15	0.92
1:A:217:ASN:HD22	1:A:217:ASN:C	1.82	0.87
1:A:37:ASP:OD2	6:A:402:HOH:O	1.92	0.87
1:B:179:GLU:OE1	6:B:404:HOH:O	1.98	0.82
1:B:256:GLU:OE1	6:B:405:HOH:O	2.00	0.79
1:B:259:VAL:HG23	6:B:401:HOH:O	1.77	0.77
1:B:133:VAL:O	6:B:406:HOH:O	2.04	0.74
1:A:197:GLU:HG2	6:A:405:HOH:O	1.90	0.72
1:B:229:ASN:HD22	1:B:232:HIS:H	1.38	0.71
1:A:229:ASN:HD22	1:A:232:HIS:H	1.37	0.71
1:B:97:GLY:C	6:B:409:HOH:O	2.33	0.71
1:A:106:LYS:NZ	6:A:404:HOH:O	2.12	0.71

Continued on next page...



 $Continued\ from\ previous\ page...$ 

Continued from pret		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ (\mathring{\rm A})$	overlap (Å)
1:B:64:ARG:NE	6:B:401:HOH:O	2.01	0.70
1:B:159:GLU:OE2	6:B:408:HOH:O	2.10	0.70
1:B:97:GLY:O	6:B:409:HOH:O	2.10	0.69
1:A:14:LYS:O	1:A:18:ASN:ND2	2.27	0.68
1:A:108:GLY:HA3	6:A:417:HOH:O	1.93	0.68
1:B:268:VAL:O	1:B:268:VAL:HG13	1.93	0.68
1:A:197:GLU:CG	6:A:405:HOH:O	2.40	0.67
1:B:196:LEU:HB3	1:B:230:ILE:HD13	1.78	0.66
1:A:249:THR:HB	1:A:251:HIS:CE1	2.36	0.60
1:B:150:GLU:CD	6:B:402:HOH:O	2.37	0.59
1:A:196:LEU:HB3	1:A:230:ILE:HD13	1.84	0.59
1:A:217:ASN:O	1:A:217:ASN:ND2	2.34	0.58
1:A:197:GLU:OE2	6:A:405:HOH:O	2.17	0.57
1:A:217:ASN:C	1:A:217:ASN:ND2	2.55	0.56
1:B:99:GLN:OE1	1:B:268:VAL:HG23	2.06	0.55
1:B:49:ASN:ND2	1:B:51:THR:H	2.04	0.55
1:B:71:PHE:HB3	1:B:223:VAL:HG21	1.89	0.55
1:A:71:PHE:HB3	1:A:223:VAL:HG21	1.90	0.54
1:B:85:ARG:NE	1:B:112[A]:GLU:OE2	2.39	0.54
1:A:13:TRP:CD1	1:A:245:MET:HE3	2.43	0.54
1:B:33:ILE:HA	6:B:418:HOH:O	2.09	0.53
1:B:35:GLU:OE2	1:B:214:ARG:NH1	2.42	0.53
1:A:146:CYS:HB3	1:A:182:ILE:HG23	1.89	0.52
1:A:26:LEU:HD11	1:B:25:PHE:HE2	1.76	0.51
1:A:247:LYS:HB3	1:A:250:TRP:CZ2	2.47	0.50
1:B:31:GLU:HG3	1:B:250:TRP:HE3	1.78	0.48
1:B:133:VAL:HG22	5:B:302:SAM:N1	2.28	0.48
1:A:45:ARG:O	1:A:45:ARG:HG2	2.13	0.48
1:B:259:VAL:HG21	6:B:401:HOH:O	1.97	0.47
1:B:8:THR:HG22	1:B:10:GLY:H	1.79	0.47
1:A:93:ALA:O	1:A:266:ARG:NH1	2.45	0.47
1:B:150:GLU:O	1:B:161:ARG:NH1	2.45	0.46
1:A:79:VAL:HG11	1:A:172:TRP:CZ3	2.51	0.46
1:B:133:VAL:HG13	5:B:302:SAM:H2	1.97	0.46
1:A:44:ALA:HB1	1:A:49:ASN:HB3	1.99	0.45
1:B:158:GLU:HB3	1:B:186:CYS:HB2	1.99	0.45
5:B:302:SAM:H2'	5:B:302:SAM:H8	1.30	0.45
1:A:176:GLY:N	1:A:177:PRO:HD2	2.32	0.44
1:B:97:GLY:HA2	6:B:409:HOH:O	2.17	0.44
1:A:104:TYR:CZ	1:A:136:LYS:HE3	2.54	0.43
1:A:30:LYS:HE2	1:A:214:ARG:HH12	1.83	0.43

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	$\operatorname{Clash}$
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
1:B:28:TYR:CZ	1:B:215:ASN:HB3	2.53	0.43
1:B:43:LYS:O	1:B:46:ARG:HG2	2.19	0.43
1:B:176:GLY:N	1:B:177:PRO:CD	2.82	0.42
1:B:268:VAL:O	1:B:268:VAL:CG1	2.64	0.42
1:A:183:LYS:HB2	1:A:221:TYR:CE2	2.54	0.42
1:B:53:GLY:HA3	1:B:260:ASN:OD1	2.21	0.41
1:B:64:ARG:HH21	1:B:259:VAL:HB	1.86	0.41
1:B:33:ILE:C	6:B:418:HOH:O	2.63	0.41
1:B:34:THR:N	6:B:418:HOH:O	2.52	0.41
1:B:97:GLY:CA	6:B:409:HOH:O	2.66	0.41
1:B:133:VAL:HG13	5:B:302:SAM:C2	2.51	0.41
1:A:168:MET:HE3	1:A:168:MET:HB3	1.94	0.41
1:A:240:VAL:O	1:A:244:ARG:HG3	2.21	0.41
1:B:199:LEU:HB3	1:B:222:TRP:CH2	2.56	0.41
1:A:176:GLY:N	1:A:177:PRO:CD	2.84	0.41
1:A:188:TYR:HA	1:A:234:VAL:HG11	2.04	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:156:GLU:OE1	1:B:51:THR:O[3_645]	2.07	0.13
6:A:461:HOH:O	6:B:407:HOH:O[2_554]	2.10	0.10

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Favoured Allowed		Outliers	Perce	ntiles	
1	A	259/265~(98%)	254 (98%)	5 (2%)	0	100	100
1	В	$261/265 \ (98\%)$	258 (99%)	3 (1%)	0	100	100
All	All	520/530 (98%)	512 (98%)	8 (2%)	0	100	100



There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Rotameric   Outliers	
1	A	$220/221 \ (100\%)$	219 (100%)	1 (0%)	86 93
1	В	222/221 (100%)	219 (99%)	3 (1%)	62 75
All	All	442/442 (100%)	438 (99%)	4 (1%)	75 85

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	217	ASN
1	В	257	GLU
1	В	258	ASP
1	В	260	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	ASN
1	A	49	ASN
1	A	54	HIS
1	A	111	HIS
1	A	217	ASN
1	A	229	ASN
1	A	232	HIS
1	В	18	ASN
1	В	49	ASN
1	В	111	HIS
1	В	200	GLN
1	В	229	ASN



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain Res		Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	В	301	-	4,4,4	0.14	0	6,6,6	0.04	0
4	GOL	A	303	-	5,5,5	0.09	0	5,5,5	0.34	0
2	SO4	A	301	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SAH	A	302	-	24,28,28	4.75	7 (29%)	25,40,40	2.66	6 (24%)
5	SAM	В	302	-	24,29,29	4.84	8 (33%)	23,42,42	4.48	6 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	303	-	-	2/4/4/4	-
3	SAH	A	302	-	-	1/11/31/31	0/3/3/3
5	SAM	В	302	-	-	2/12/33/33	0/3/3/3



All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	Observed(A)	$\operatorname{Ideal}( ext{\AA})$
5	В	302	SAM	C2'-C1'	-17.43	1.27	1.53
3	A	302	SAH	C2'-C1'	-15.43	1.30	1.53
3	A	302	SAH	O4'-C1'	13.95	1.60	1.41
5	В	302	SAM	C3'-C4'	-9.72	1.28	1.53
5	В	302	SAM	O4'-C1'	9.40	1.54	1.41
3	A	302	SAH	O4'-C4'	-7.01	1.29	1.45
5	В	302	SAM	C2'-C3'	4.46	1.65	1.53
5	В	302	SAM	O4'-C4'	3.97	1.53	1.45
3	A	302	SAH	O3'-C3'	-3.90	1.33	1.43
3	A	302	SAH	O2'-C2'	2.93	1.49	1.43
5	В	302	SAM	C5-C4	-2.92	1.33	1.40
5	В	302	SAM	C6-N6	2.82	1.44	1.34
3	A	302	SAH	C5-C4	-2.48	1.34	1.40
3	A	302	SAH	C6-N6	2.40	1.42	1.34
5	В	302	SAM	OXT-C	-2.23	1.23	1.30

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$Ideal(^{o})$
5	В	302	SAM	C1'-N9-C4	14.28	151.73	126.64
5	В	302	SAM	C5-C6-N6	11.53	137.88	120.35
3	A	302	SAH	C5-C6-N6	8.35	133.04	120.35
5	В	302	SAM	N6-C6-N1	-7.78	102.43	118.57
3	A	302	SAH	N6-C6-N1	-5.68	106.78	118.57
5	В	302	SAM	N3-C2-N1	-5.44	120.18	128.68
3	A	302	SAH	N3-C2-N1	-5.05	120.78	128.68
3	A	302	SAH	C1'-N9-C4	-4.34	119.01	126.64
5	В	302	SAM	O4'-C1'-C2'	-4.02	101.05	106.93
3	A	302	SAH	C3'-C2'-C1'	2.99	105.47	100.98
3	A	302	SAH	C5'-C4'-C3'	-2.47	108.88	115.06
5	В	302	SAM	CG-SD-C5'	2.19	108.98	103.40

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	SAH	CA-CB-CG-SD
4	A	303	GOL	O1-C1-C2-C3
4	A	303	GOL	O1-C1-C2-O2
5	В	302	SAM	CA-CB-CG-SD
5	В	302	SAM	C4'-C5'-SD-CG




There are no ring outliers.


2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	SO4	1	0
5	В	302	SAM	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	178:ARG	С	179:GLU	N	0.99



# 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$		$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9		
1	A	$261/265 \ (98\%)$	1.08	48 (18%)	4	4		18, 34, 58, 74	0
1	В	$262/265 \ (98\%)$	0.83	32 (12%)	10	8		11, 33, 54, 70	1 (0%)
All	All	523/530 (98%)	0.95	80 (15%)	6	5		11, 34, 55, 74	1 (0%)

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	47	ASP	7.2
1	A	250	TRP	6.0
1	В	249	THR	5.4
1	A	46	ARG	5.3
1	A	249	THR	5.2
1	A	267	ALA	5.2
1	A	48	GLY	5.1
1	В	268	VAL	4.7
1	A	37	ASP	4.6
1	В	8	THR	4.6
1	A	248	ARG	4.2
1	A	43	LYS	4.0
1	В	45	ARG	3.9
1	A	39	SER	3.9
1	A	251	HIS	3.9
1	В	257	GLU	3.9
1	В	236	MET	3.8
1	В	251	HIS	3.7
1	В	250	TRP	3.7
1	A	53	GLY	3.6
1	В	248	ARG	3.5
1	A	259	VAL	3.4
1	В	58	ARG	3.3
1	A	26	LEU	3.3

Continued on next page...



Continued from previous page...

Mol	nued fron Chain	Res	Type	RSRZ
1	В	259	VAL	3.2
1	В	48	GLY	3.2
1	A	27	LYS	3.1
1	A	253	PRO	3.1
1	A	255	TYR	3.1
1	A	49	ASN	3.1
1	В	267	ALA	3.1
1	A	44	ALA	3.1
1	A	204	GLY	3.0
1	A	246	GLU	3.0
1	В	232	HIS	3.0
1	A	247	LYS	3.0
1	A	20	LEU	3.0
1	В	260	ASN	2.9
1	A	52	GLY	2.8
1	В	20	LEU	2.8
1	A	217	ASN	2.8
1	A	40	ALA	2.8
1	A	257	GLU	2.8
1	В	47	ASP	2.7
1	A	236	MET	2.7
1	A	239	GLN	2.7
1	A	232	HIS	2.7
1	В	255	TYR	2.6
1	В	256	GLU	2.5
1	В	156	GLU	2.5
1	A	19	GLY	2.5
1	В	247	LYS	2.5
1	В	39	SER	2.5
1	A	9	LEU	2.4
1	В	240	VAL	2.4
1	A	153	SER	2.4
1	В	7	ARG	2.4
1	A	17	LEU	2.4
1	A	258	ASP	2.3
1	В	26	LEU	2.3
1	A	41	ALA	2.3
1	A	156	GLU	2.3
1	A	8	THR	2.3
1	В	46	ARG	2.3
1	A	260	ASN	2.2
1	В	51	THR	2.2

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	В	42	ARG	2.2
1	A	240	VAL	2.2
1	A	171	ASP	2.2
1	A	154	SER	2.1
1	В	242	ILE	2.1
1	A	21	SER	2.1
1	В	16	LYS	2.1
1	В	171	ASP	2.1
1	В	43	LYS	2.1
1	A	230	ILE	2.1
1	A	265	THR	2.1
1	A	45	ARG	2.0
1	A	203	TYR	2.0
1	A	197	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
4	GOL	A	303	6/6	0.73	0.17	20,20,20,20	0
2	SO4	A	301	5/5	0.75	0.13	51,61,66,88	0
2	SO4	В	301	5/5	0.79	0.11	49,55,62,77	0
3	SAH	A	302	26/26	0.90	0.10	20,20,20,20	0
5	SAM	В	302	27/27	0.91	0.11	23,23,24,24	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




#### Electron density around SAH A 302:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around SAM B 302:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)









# 6.5 Other polymers (i)

There are no such residues in this entry.

