



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:37 PM UTC

PDB ID : 8BQ8 / pdb_00008bq8
Title : Crystal structure of Trichoplax Dlg PDZ2 domain in complex with Trichoplax Vangl peptide
Authors : Madduamge, J.C.; Kvensakul, M.
Deposited on : 2022-11-20
Resolution : 2.70 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : **NOT EXECUTED**
Xtrriage (Phenix) : 2.0
EDS : **NOT EXECUTED**
Buster-report : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	94	
1	B	94	
1	C	94	
2	D	8	
2	E	8	
2	F	8	

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 4317 atoms, of which 2146 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Disks large-like protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	89	1332	416	673	114	128	1	0	0	0
1	B	90	1333	417	673	115	127	1	0	0	0
1	C	93	1373	431	693	117	131	1	0	0	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	312	GLY	-	expression tag	UNP A0A369SI82
A	313	PRO	-	expression tag	UNP A0A369SI82
A	314	LEU	-	expression tag	UNP A0A369SI82
A	315	GLY	-	expression tag	UNP A0A369SI82
A	316	SER	-	expression tag	UNP A0A369SI82
B	312	GLY	-	expression tag	UNP A0A369SI82
B	313	PRO	-	expression tag	UNP A0A369SI82
B	314	LEU	-	expression tag	UNP A0A369SI82
B	315	GLY	-	expression tag	UNP A0A369SI82
B	316	SER	-	expression tag	UNP A0A369SI82
C	312	GLY	-	expression tag	UNP A0A369SI82
C	313	PRO	-	expression tag	UNP A0A369SI82
C	314	LEU	-	expression tag	UNP A0A369SI82
C	315	GLY	-	expression tag	UNP A0A369SI82
C	316	SER	-	expression tag	UNP A0A369SI82

- Molecule 2 is a protein called Vang-like protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	D	5	71	22	34	5	10	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
			Total	C	H	N				O
2	F	5	Total	C	H	N	O	0	0	0
			71	22	34	5	10			
2	E	6	Total	C	H	N	O	0	0	0
			84	26	39	7	12			

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	17	Total	O	0	0
			17	17		
3	B	18	Total	O	0	0
			18	18		
3	C	18	Total	O	0	0
			18	18		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: Disks large-like protein 1

Chain A:  84% 11% 5%




- Molecule 1: Disks large-like protein 1

Chain B:  83% 13% .



- Molecule 1: Disks large-like protein 1

Chain C:  81% 16% ...




- Molecule 2: Vang-like protein 1

Chain D:  50% 12% 38%




- Molecule 2: Vang-like protein 1

Chain F:  38% 25% 38%



- Molecule 2: Vang-like protein 1



4 Data and refinement statistics

EDS was not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	64.67Å 101.98Å 106.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.34 – 2.70	Depositor
% Data completeness (in resolution range)	97.7 (32.34-2.70)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 2.68Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.222 , 0.272	Depositor
Wilson B-factor (Å ²)	30.4	Xtrriage
Anisotropy	0.551	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	4317	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.14	0/664	0.35	0/897
1	B	0.15	0/665	0.35	0/898
1	C	0.15	0/687	0.38	0/930
2	D	0.16	0/37	0.33	0/48
2	E	0.18	0/45	0.38	0/60
2	F	0.18	0/37	0.25	0/48
All	All	0.15	0/2135	0.36	0/2881

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	659	673	673	8	0
1	B	660	673	674	9	0
1	C	680	693	694	12	0
2	D	37	34	34	1	0
2	E	45	39	39	0	0
2	F	37	34	34	3	0
3	A	17	0	0	0	0
3	B	18	0	0	0	0
3	C	18	0	0	0	0
All	All	2171	2146	2148	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:317:LEU:HD21	1:B:403:VAL:HG22	1.44	0.99
1:B:317:LEU:CD2	1:B:403:VAL:HG22	1.98	0.94
1:A:326:ASP:OD1	1:A:328:GLY:N	2.22	0.71
1:C:391:LEU:HB3	2:F:195:VAL:HG21	1.75	0.67
1:C:342:ILE:HD12	1:C:349:PHE:CZ	2.33	0.63
1:C:391:LEU:CB	2:F:195:VAL:HG21	2.35	0.56
1:A:320:ILE:HG22	1:A:322:LEU:CD1	2.36	0.56
1:C:369:ASP:OD2	1:C:404:LYS:NZ	2.41	0.53
1:B:351:THR:O	1:B:352:LYS:HG3	2.09	0.52
1:C:371:ILE:HD13	1:C:402:VAL:HG12	1.92	0.52
1:A:320:ILE:HG22	1:A:322:LEU:HD11	1.92	0.50
1:A:325:GLU:O	1:A:326:ASP:CB	2.60	0.49
1:C:393:GLU:O	1:C:394:THR:O	2.32	0.47
1:A:325:GLU:O	1:A:326:ASP:HB3	2.17	0.45
1:A:321:VAL:C	1:A:322:LEU:HD12	2.42	0.45
1:C:315:GLY:CA	1:C:404:LYS:O	2.66	0.44
1:C:342:ILE:HD12	1:C:349:PHE:CE2	2.52	0.44
1:B:373:LYS:HB3	1:B:401:VAL:HB	1.99	0.44
1:B:373:LYS:HE2	1:B:375:ASN:O	2.18	0.43
1:A:391:LEU:HB3	2:D:152:VAL:HG21	1.99	0.43
1:B:325:GLU:O	1:B:326:ASP:C	2.62	0.42
1:C:372:THR:HG22	1:C:401:VAL:O	2.20	0.42
1:C:373:LYS:HB3	1:C:401:VAL:HB	2.03	0.41
1:B:366:GLU:HB2	1:B:404:LYS:HZ2	1.84	0.41
1:C:393:GLU:HG2	1:C:394:THR:N	2.36	0.41
2:F:195:VAL:HG22	2:F:195:VAL:OXT	2.21	0.41
1:B:369:ASP:OD1	1:B:404:LYS:HD3	2.21	0.40
1:C:317:LEU:CD2	1:C:403:VAL:HG22	2.52	0.40
1:A:370:ARG:O	1:A:402:VAL:HA	2.21	0.40
1:B:317:LEU:HD23	1:B:403:VAL:HG22	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	85/94 (90%)	82 (96%)	3 (4%)	0	100	100
1	B	16/94 (17%)	16 (100%)	0	0	100	100
1	C	91/94 (97%)	87 (96%)	2 (2%)	2 (2%)	5	14
2	D	3/8 (38%)	3 (100%)	0	0	100	100
2	E	4/8 (50%)	4 (100%)	0	0	100	100
2	F	3/8 (38%)	3 (100%)	0	0	100	100
All	All	202/306 (66%)	195 (96%)	5 (2%)	2 (1%)	12	32

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	394	THR
1	C	393	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	72/76 (95%)	71 (99%)	1 (1%)	59	82
1	B	71/76 (93%)	71 (100%)	0	100	100
1	C	73/76 (96%)	70 (96%)	3 (4%)	27	56
2	D	5/8 (62%)	5 (100%)	0	100	100
2	E	6/8 (75%)	4 (67%)	2 (33%)	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	F	5/8 (62%)	4 (80%)	1 (20%)	1	4
All	All	232/252 (92%)	225 (97%)	7 (3%)	36	66

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	343	ILE
1	C	314	LEU
1	C	376	THR
1	C	394	THR
2	F	194	SER
2	E	256	ASN
2	E	261	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	341	HIS
1	B	341	HIS
1	C	319	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

Mogul was not executed - this section is therefore empty.

5.5 Carbohydrates [i](#)

Mogul was not executed - this section is therefore empty.

5.6 Ligand geometry [i](#)

Mogul was not executed - this section is therefore empty.

5.7 Other polymers [i](#)

Mogul was not executed - this section is therefore empty.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.