



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 18, 2025 – 08:09 PM EST

PDB ID : 8EGZ / pdb\_00008egz  
Title : Engineered tyrosine synthase (TmTyrS1) derived from *T. maritima* TrpB with Ser bound as the amino-acrylate intermediate  
Authors : Porter, N.J.; Almhjell, P.J.; Arnold, F.H.  
Deposited on : 2022-09-13  
Resolution : 1.90 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.46


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	397	 91% 6%
1	B	397	 90% 7%

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 6262 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Engineered tyrosine synthase (TmTyrS1).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	385	2970	1896	510	556	8	0	5	0
1	B	386	2954	1883	514	550	7	0	2	0

There are 52 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	4	ASN	TYR	engineered mutation	UNP P50909
A	12	ASN	TYR	engineered mutation	UNP P50909
A	19	GLY	PRO	engineered mutation	UNP P50909
A	30	GLY	GLU	engineered mutation	UNP P50909
A	41	TYR	PHE	engineered mutation	UNP P50909
A	69	VAL	ILE	engineered mutation	UNP P50909
A	96	LEU	LYS	engineered mutation	UNP P50909
A	103	THR	ILE	engineered mutation	UNP P50909
A	105	GLY	GLU	engineered mutation	UNP P50909
A	140	LEU	PRO	engineered mutation	UNP P50909
A	167	ASP	ASN	engineered mutation	UNP P50909
A	184	PRO	ILE	engineered mutation	UNP P50909
A	213	PRO	LEU	engineered mutation	UNP P50909
A	228	SER	GLY	engineered mutation	UNP P50909
A	291	ALA	VAL	engineered mutation	UNP P50909
A	292	SER	THR	engineered mutation	UNP P50909
A	302	PRO	SER	engineered mutation	UNP P50909
A	389	HIS	ARG	engineered mutation	UNP P50909
A	390	LEU	-	expression tag	UNP P50909
A	391	GLU	-	expression tag	UNP P50909
A	392	HIS	-	expression tag	UNP P50909
A	393	HIS	-	expression tag	UNP P50909
A	394	HIS	-	expression tag	UNP P50909
A	395	HIS	-	expression tag	UNP P50909
A	396	HIS	-	expression tag	UNP P50909

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
A	397	HIS	-	expression tag	UNP P50909
B	4	ASN	TYR	engineered mutation	UNP P50909
B	12	ASN	TYR	engineered mutation	UNP P50909
B	19	GLY	PRO	engineered mutation	UNP P50909
B	30	GLY	GLU	engineered mutation	UNP P50909
B	41	TYR	PHE	engineered mutation	UNP P50909
B	69	VAL	ILE	engineered mutation	UNP P50909
B	96	LEU	LYS	engineered mutation	UNP P50909
B	103	THR	ILE	engineered mutation	UNP P50909
B	105	GLY	GLU	engineered mutation	UNP P50909
B	140	LEU	PRO	engineered mutation	UNP P50909
B	167	ASP	ASN	engineered mutation	UNP P50909
B	184	PRO	ILE	engineered mutation	UNP P50909
B	213	PRO	LEU	engineered mutation	UNP P50909
B	228	SER	GLY	engineered mutation	UNP P50909
B	291	ALA	VAL	engineered mutation	UNP P50909
B	292	SER	THR	engineered mutation	UNP P50909
B	302	PRO	SER	engineered mutation	UNP P50909
B	389	HIS	ARG	engineered mutation	UNP P50909
B	390	LEU	-	expression tag	UNP P50909
B	391	GLU	-	expression tag	UNP P50909
B	392	HIS	-	expression tag	UNP P50909
B	393	HIS	-	expression tag	UNP P50909
B	394	HIS	-	expression tag	UNP P50909
B	395	HIS	-	expression tag	UNP P50909
B	396	HIS	-	expression tag	UNP P50909
B	397	HIS	-	expression tag	UNP P50909

- Molecule 2 is 2-{{(E)-{3-hydroxy-2-methyl-5-[(phosphonoxy)methyl]pyridin-4-yl}methylidene}amino}prop-2-enoic acid (CCD ID: 0JO) (formula: C<sub>11</sub>H<sub>13</sub>N<sub>2</sub>O<sub>7</sub>P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			21	11	2	7	1		
2	B	1	Total	C	N	O	P	0	0
			21	11	2	7	1		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0

- Molecule 4 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total K 1 1	0	0
4	B	1	Total K 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	157	Total O 158 158	0	1
5	B	112	Total O 112 112	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Engineered tyrosine synthase (TmTyrS1)



- Molecule 1: Engineered tyrosine synthase (TmTyrS1)



## 4 Data and refinement statistics i

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	164.57Å 164.57Å 84.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.13 – 1.90 40.13 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (40.13-1.90) 99.9 (40.13-1.90)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.72 (at 1.89Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.170 , 0.192 0.170 , 0.192	Depositor DCC
$R_{free}$ test set	4532 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.8	Xtrriage
Anisotropy	0.075	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 34.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.027 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6262	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K, OJO, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.84	0/3047	0.86	1/4126 (0.0%)
1	B	0.80	0/3019	0.83	0/4089
All	All	0.82	0/6066	0.85	1/8215 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	22	GLU	CA-CB-CG	-5.10	103.90	114.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2970	0	2968	13	0
1	B	2954	0	2941	20	0
2	A	21	0	9	0	0
2	B	21	0	9	1	0
3	A	16	0	24	4	0
3	B	8	0	12	3	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	158	0	0	2	0
5	B	112	0	0	1	0
All	All	6262	0	5963	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:95:LYS:HD3	1:B:124:MET:HE2	1.58	0.84
1:B:113:VAL:O	1:B:117:THR:HG23	1.83	0.78
1:B:8:TYR:HE1	1:B:271:ILE:HD11	1.49	0.76
1:A:50:GLY:O	3:A:402:EDO:H21	1.85	0.76
1:A:28:TYR:CZ	1:A:32:MET:HE3	2.21	0.75
3:A:402:EDO:H12	5:A:528:HOH:O	1.94	0.67
1:B:166:ILE:HD13	3:B:403:EDO:H21	1.80	0.64
3:A:404:EDO:H22	3:A:405:EDO:H12	1.84	0.60
1:A:228:SER:HB2	1:A:299:LEU:HA	1.85	0.58
1:B:8:TYR:CE1	1:B:271:ILE:HD11	2.38	0.54
1:A:271:ILE:HG13	5:A:615:HOH:O	2.09	0.53
1:A:60:ARG:HD3	1:A:337:LEU:O	2.09	0.52
1:B:206:GLU:O	1:B:210:LYS:HG2	2.10	0.52
1:B:158:GLY:HA3	1:B:164:ASP:OD2	2.11	0.50
1:A:32:MET:HE1	1:A:38:TRP:HZ2	1.76	0.49
1:A:372[B]:ASP:OD1	1:A:373:LYS:N	2.44	0.49
3:A:402:EDO:H22	5:B:578:HOH:O	2.11	0.49
1:A:59:ARG:O	1:A:63:GLU:HG3	2.14	0.48
1:B:11:GLN:OE1	1:B:18:MET:HE1	2.15	0.47
1:B:182:TYR:OH	3:B:403:EDO:H22	2.16	0.46
1:B:31:ILE:HD12	1:B:32:MET:N	2.31	0.45
1:A:95:LYS:HE2	1:A:122:PHE:O	2.17	0.45
1:A:210:LYS:HB3	1:A:210:LYS:HE2	1.88	0.45
1:A:28:TYR:CE1	1:A:32:MET:HE3	2.51	0.45
1:B:182:TYR:OH	3:B:403:EDO:H11	2.17	0.44
1:B:212:ILE:CG2	1:B:213:PRO:HD3	2.49	0.43
1:B:344:LEU:HD23	1:B:344:LEU:HA	1.89	0.42
1:A:75:ASP:HB2	1:A:370:ARG:HB3	2.00	0.42
1:B:296:SER:OG	1:B:345:GLU:HG3	2.20	0.42
1:A:212:ILE:HG21	1:A:220:PRO:HD3	2.02	0.41
1:B:212:ILE:HG22	1:B:213:PRO:HD3	2.02	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:60:ARG:HH11	1:B:60:ARG:HG3	1.85	0.41
1:B:189:GLY:HA2	1:B:275:HIS:O	2.20	0.41
2:B:401:OJO:O3	2:B:401:OJO:N	2.54	0.41
1:B:117:THR:HG22	1:B:148:LEU:HD13	2.02	0.40
1:B:168:GLU:OE2	1:B:171:ARG:NH2	2.51	0.40
1:B:31:ILE:HD12	1:B:31:ILE:C	2.47	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/397 (98%)	381 (98%)	7 (2%)	0	100	100
1	B	386/397 (97%)	379 (98%)	7 (2%)	0	100	100
All	All	774/794 (98%)	760 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	306/319 (96%)	305 (100%)	1 (0%)	91	92
1	B	301/319 (94%)	300 (100%)	1 (0%)	91	92

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	607/638 (95%)	605 (100%)	2 (0%)	91	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	356	ILE
1	B	117	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	ASN
1	A	90	GLN
1	B	86	ASN
1	B	90	GLN
1	B	290	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	A	403	-	3,3,3	0.43	0	2,2,2	0.67	0
3	EDO	A	404	-	3,3,3	0.76	0	2,2,2	0.29	0
3	EDO	B	402	-	3,3,3	0.64	0	2,2,2	0.59	0
2	0JO	B	401	-	20,21,21	1.68	6 (30%)	23,30,30	1.77	9 (39%)
3	EDO	A	405	-	3,3,3	0.39	0	2,2,2	0.54	0
3	EDO	B	403	-	3,3,3	0.73	0	2,2,2	0.51	0
3	EDO	A	402	-	3,3,3	0.51	0	2,2,2	0.64	0
2	0JO	A	401	-	20,21,21	2.03	8 (40%)	23,30,30	1.92	5 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	403	-	-	0/1/1/1	-
3	EDO	A	404	-	-	0/1/1/1	-
3	EDO	B	402	-	-	0/1/1/1	-
2	0JO	B	401	-	-	0/11/15/15	0/1/1/1
3	EDO	A	405	-	-	0/1/1/1	-
3	EDO	B	403	-	-	1/1/1/1	-
3	EDO	A	402	-	-	0/1/1/1	-
2	0JO	A	401	-	-	0/11/15/15	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	0JO	C5A-C5	3.60	1.60	1.50
2	A	401	0JO	C2A-C2	3.46	1.55	1.50
2	A	401	0JO	CA-N	3.37	1.51	1.36
2	A	401	0JO	O-C	3.05	1.30	1.22
2	A	401	0JO	C5A-C5	3.03	1.58	1.50
2	A	401	0JO	C4-C3	2.94	1.45	1.41
2	B	401	0JO	C2A-C2	2.77	1.54	1.50
2	B	401	0JO	C4-C3	2.53	1.45	1.41
2	B	401	0JO	C6-C5	2.49	1.42	1.37
2	A	401	0JO	C4A-N	2.34	1.31	1.28
2	A	401	0JO	C4-C4A	2.28	1.51	1.46
2	A	401	0JO	OXT-C	-2.28	1.24	1.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	OJO	CA-N	2.18	1.46	1.36
2	B	401	OJO	P-OP4	2.14	1.67	1.60

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	OJO	C4-C3-C2	-5.31	117.15	120.14
2	A	401	OJO	OP4-P-OP3	-3.90	95.91	106.44
2	B	401	OJO	C2A-C2-C3	-3.63	116.55	120.80
2	B	401	OJO	C4-C3-C2	-2.85	118.54	120.14
2	B	401	OJO	C3-C2-N1	2.79	124.48	120.96
2	B	401	OJO	OP1-P-OP3	2.67	121.24	110.83
2	A	401	OJO	C4-C4A-N	-2.65	116.90	123.01
2	B	401	OJO	OP1-P-OP4	-2.64	99.80	106.67
2	A	401	OJO	C2A-C2-C3	-2.46	117.92	120.80
2	B	401	OJO	C4-C4A-N	-2.41	117.46	123.01
2	A	401	OJO	O3-C3-C2	2.25	122.25	117.58
2	B	401	OJO	OP4-C5A-C5	-2.13	105.35	109.36
2	B	401	OJO	OP4-P-OP3	-2.13	100.69	106.44
2	B	401	OJO	C3-C4-C5	-2.02	116.66	118.28

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	403	EDO	O1-C1-C2-O2


There are no ring outliers.


5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	404	EDO	1	0
2	B	401	OJO	1	0
3	A	405	EDO	1	0
3	B	403	EDO	3	0
3	A	402	EDO	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	385/397 (96%)	-0.18	2 (0%) 87 88	16, 30, 44, 74	5 (1%)
1	B	386/397 (97%)	0.12	7 (1%) 67 70	17, 33, 51, 72	2 (0%)
All	All	771/794 (97%)	-0.03	9 (1%) 76 78	16, 31, 49, 74	7 (0%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	293[A]	HIS	3.7
1	B	2	LYS	3.4
1	B	387	ARG	2.9
1	B	164	ASP	2.6
1	B	133	GLU	2.5
1	A	29	GLU	2.4
1	B	32	MET	2.3
1	A	3	GLY	2.2
1	B	386	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	EDO	A	403	4/4	0.68	0.20	45,48,57,69	0
3	EDO	A	405	4/4	0.82	0.13	53,54,55,60	0
3	EDO	B	403	4/4	0.86	0.23	39,48,50,50	0
3	EDO	A	404	4/4	0.87	0.16	47,56,59,62	0
3	EDO	B	402	4/4	0.90	0.13	38,44,46,52	0
3	EDO	A	402	4/4	0.91	0.23	40,42,46,49	0
2	0JO	B	401	21/21	0.97	0.07	26,30,38,41	0
2	0JO	A	401	21/21	0.98	0.06	25,29,34,37	0
4	K	A	406	1/1	0.99	0.05	32,32,32,32	0
4	K	B	404	1/1	0.99	0.09	36,36,36,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 0JO B 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.