

Full wwPDB X-ray Structure Validation Report (i)

Dec 2, 2025 - 04:02 am GMT

PDB ID : 8H26 / pdb 00008h26

Title: Crystal structure of MnmM from S. aureus complexed with SAH (1.50 A)

Authors: Kim, J.; Cho, G.; Lee, J.

Deposited on : 2022-10-05

Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4\text{-}5\text{-}2 \text{ with } Phenix 2.0$

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 2.0

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.010 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.46

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	164625	3717 (1.50-1.50)
Clashscore	180529	4048 (1.50-1.50)
Ramachandran outliers	177936	3970 (1.50-1.50)
Sidechain outliers	177891	3967 (1.50-1.50)
RSRZ outliers	164620	3718 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
			8%	
1	A	195	78%	12% 10%
			6%	
1	В	195	84%	6% 10%
			5%	
1	С	195	79%	15% • 6%
			9%	
1	D	195	80%	9% • 10%
			9%	
1	Е	195	76%	10% 13%



 $Continued\ from\ previous\ page...$

Mol	Chain	Length	Quality of chain			
	_		7%			
1	F'	195	78%	7%	٠	14%



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 9201 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 16S rRNA (Cytosine(1402)-N(4))-methyltransferase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	175	Total	С	N	О	S	0	2	0
1	A	175	1406	896	241	263	6	0	<u> </u>	U
1	В	175	Total	С	N	О	S	0	2	0
1	Б	175	1402	893	241	262	6	0	<u> </u>	U
1	С	184	Total	С	N	О	S	0	5	0
1		104	1495	951	259	278	7			U
1	D	175	Total	С	N	О	S	0	4	0
1	D	175	1413	897	244	266	6	0	4	U
1	Е	169	Total	С	N	О	S	0	2	0
1	E	109	1353	860	235	253	5	0		U
1	F	168	Total	С	N	О	S	0	2	0
1	I'	100	1350	860	234	251	5		<u> </u>	U

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	188	GLY	-	expression tag	UNP Q2FXG9
A	189	HIS	-	expression tag	UNP Q2FXG9
A	190	HIS	-	expression tag	UNP Q2FXG9
A	191	HIS	-	expression tag	UNP Q2FXG9
A	192	HIS	-	expression tag	UNP Q2FXG9
A	193	HIS	-	expression tag	UNP Q2FXG9
A	194	HIS	-	expression tag	UNP Q2FXG9
A	195	GLY	-	expression tag	UNP Q2FXG9
В	188	GLY	-	expression tag	UNP Q2FXG9
В	189	HIS	-	expression tag	UNP Q2FXG9
В	190	HIS	-	expression tag	UNP Q2FXG9
В	191	HIS	_	expression tag	UNP Q2FXG9
В	192	HIS	-	expression tag	UNP Q2FXG9
В	193	HIS	_	expression tag	UNP Q2FXG9
В	194	HIS	-	expression tag	UNP Q2FXG9
В	195	GLY	-	expression tag	UNP Q2FXG9
С	188	GLY	-	expression tag	UNP Q2FXG9




 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
С	189	HIS	-	expression tag	UNP Q2FXG9
С	190	HIS	-	expression tag	UNP Q2FXG9
С	191	HIS	-	expression tag	UNP Q2FXG9
С	192	HIS	-	expression tag	UNP Q2FXG9
С	193	HIS	-	expression tag	UNP Q2FXG9
С	194	HIS	-	expression tag	UNP Q2FXG9
С	195	GLY	-	expression tag	UNP Q2FXG9
D	188	GLY	-	expression tag	UNP Q2FXG9
D	189	HIS	-	expression tag	UNP Q2FXG9
D	190	HIS	-	expression tag	UNP Q2FXG9
D	191	HIS	-	expression tag	UNP Q2FXG9
D	192	HIS	-	expression tag	UNP Q2FXG9
D	193	HIS	-	expression tag	UNP Q2FXG9
D	194	HIS	-	expression tag	UNP Q2FXG9
D	195	GLY	-	expression tag	UNP Q2FXG9
Е	188	GLY	-	expression tag	UNP Q2FXG9
Е	189	HIS	-	expression tag	UNP Q2FXG9
Е	190	HIS	-	expression tag	UNP Q2FXG9
Е	191	HIS	-	expression tag	UNP Q2FXG9
Е	192	HIS	-	expression tag	UNP Q2FXG9
Е	193	HIS	-	expression tag	UNP Q2FXG9
Е	194	HIS	-	expression tag	UNP Q2FXG9
Е	195	GLY	-	expression tag	UNP Q2FXG9
F	188	GLY	-	expression tag	UNP Q2FXG9
F	189	HIS	-	expression tag	UNP Q2FXG9
F	190	HIS	-	expression tag	UNP Q2FXG9
F	191	HIS	-	expression tag	UNP Q2FXG9
F	192	HIS	-	expression tag	UNP Q2FXG9
F	193	HIS	-	expression tag	UNP Q2FXG9
F	194	HIS	-	expression tag	UNP Q2FXG9
F	195	GLY	-	expression tag	UNP Q2FXG9

 \bullet Molecule 2 is S-ADENOSYL-L-HOMOCYSTEINE (CCD ID: SAH) (formula: $C_{14}H_{20}N_6O_5S)$ (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf				
2	Λ.	٨	٨	Λ	1	Total	С	N	О	S	0	0
2	A	1	26	14	6	5	1	0	0			
2	D	1	Total	С	N	О	S	0	0			
2	В	1	26	14	6	5	1	0				
2	C	1	Total	С	N	О	S	0	0			
2		1	26	14	6	5	1	0	U			
2	D	1	Total	С	N	О	S	0	0			
2	ש	1	26	14	6	5	1		U			


• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	119	Total O 119 119	0	0
3	В	134	Total O 134 134	0	0
3	C	130	Total O 130 130	0	0
3	D	90	Total O 90 90	0	0
3	E	111	Total O 111 111	0	0
3	F	94	Total O 94 94	0	0




3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.








• Molecule 1: 16S rRNA (Cytosine(1402)-N(4))-methyltransferase



• Molecule 1: 16S rRNA (Cytosine(1402)-N(4))-methyltransferase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	62.33Å 69.36Å 74.61Å	Depositor
a, b, c, α , β , γ	91.25° 112.97° 114.05°	Depositor
Resolution (Å)	67.09 - 1.50	Depositor
resolution (11)	67.08 - 1.50	EDS
% Data completeness	63.4 (67.09-1.50)	Depositor
(in resolution range)	63.4 (67.08-1.50)	EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.82 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.183 , 0.219	Depositor
it, it free	0.190 , 0.225	DCC
R_{free} test set	5112 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	21.6	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33 , 37.4	EDS
L-test for twinning ²	$ < L > = 0.51, < L^2> = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9201	wwPDB-VP
Average B, all atoms $(Å^2)$	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.37% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
Wioi Chain		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.90	1/1432 (0.1%)	0.98	0/1937	
1	В	0.88	0/1427	0.95	0/1930	
1	С	0.94	0/1525	0.99	3/2063 (0.1%)	
1	D	0.91	1/1438 (0.1%)	1.00	1/1944 (0.1%)	
1	Е	0.90	1/1379 (0.1%)	0.96	$2/1865 \ (0.1\%)$	
1	F	0.91	1/1376 (0.1%)	1.07	2/1861 (0.1%)	
All	All	0.91	4/8577 (0.0%)	0.99	8/11600 (0.1%)	

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$Ideal(\AA)$
1	F	75	ASP	C-O	-6.23	1.16	1.23
1	Е	179	PHE	CA-C	-5.19	1.46	1.52
1	D	58	GLU	CG-CD	5.19	1.65	1.52
1	A	163	ALA	C-O	-5.11	1.17	1.23

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$\operatorname{Ideal}({}^{o})$
1	F	150	ALA	N-CA-C	9.31	122.37	111.02
1	Е	188	GLY	N-CA-C	-5.87	105.50	113.37
1	F	172	ASN	N-CA-C	5.71	118.55	109.24
1	D	145	GLN	N-CA-C	-5.66	103.47	111.56
1	С	172	ASN	N-CA-C	5.13	117.98	109.46
1	Е	172	ASN	N-CA-C	5.12	117.08	108.73
1	С	29[A]	CYS	N-CA-C	5.01	116.74	111.28
1	С	29[B]	CYS	N-CA-C	5.01	116.74	111.28

There are no chirality outliers.

There are no planarity outliers.



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1406	0	1401	16	0
1	В	1402	0	1403	9	0
1	С	1495	0	1488	27	0
1	D	1413	0	1407	14	0
1	Ε	1353	0	1342	10	0
1	F	1350	0	1339	8	0
2	A	26	0	19	0	0
2	В	26	0	19	1	0
2	С	26	0	19	2	0
2	D	26	0	19	2	0
3	A	119	0	0	1	0
3	В	134	0	0	0	0
3	С	130	0	0	6	1
3	D	90	0	0	3	0
3	Ε	111	0	0	3	1
3	F	94	0	0	2	0
All	All	9201	0	8456	74	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:C:155:LEU:HD13	1:C:181[B]:CYS:SG	1.94	1.06
1:A:155:LEU:HD13	1:A:181[B]:CYS:SG	2.05	0.97
1:F:139:HIS:ND1	3:F:201:HOH:O	2.04	0.80
1:D:21:GLU:OE1	3:D:301:HOH:O	2.01	0.78
1:C:118:GLN:OE1	3:C:301:HOH:O	2.02	0.77
1:C:155:LEU:CD1	1:C:181[B]:CYS:SG	2.75	0.73
1:D:144:GLY:N	3:D:303:HOH:O	2.26	0.69
1:C:123:LEU:O	1:C:127[A]:MET:HG2	1.94	0.68
1:B:123:LEU:O	1:B:127[B]:MET:HG2	1.93	0.67
1:B:153:ASP:HB3	1:D:145:GLN:NE2	2.10	0.67
1:A:123:LEU:O	1:A:127[A]:MET:HG2	1.94	0.67



 $Continued\ from\ previous\ page...$

Atom 1		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	$\text{overlap } (\mathring{\mathbf{A}})$
1:C:147:GLU:OE1	3:C:302:HOH:O	2.12	0.66
1:C:55:LEU:C	1:C:55:LEU:HD12	2.22	0.65
1:C:145:GLN:OE1	3:C:303:HOH:O	2.13	0.65
1:C:7:LEU:HD23	3:C:393:HOH:O	1.97	0.64
1:D:157:THR:OG1	3:D:302:HOH:O	2.02	0.64
1:E:84:ILE:O	1:E:89:LYS:NZ	2.33	0.62
1:D:123:LEU:O	1:D:127[A]:MET:HG2	2.00	0.61
1:A:7:LEU:HD12	1:C:21:GLU:HG2	1.81	0.61
1:A:155:LEU:CD1	1:A:181[B]:CYS:SG	2.86	0.61
1:D:14:ILE:HD12	1:D:94:ALA:HB1	1.82	0.60
1:C:17:HIS:HE1	1:C:184:GLU:OE1	1.85	0.60
1:A:7:LEU:HD12	1:C:21:GLU:CG	2.34	0.58
1:F:174:ARG:HG3	1:F:177:ALA:HB3	1.86	0.57
1:A:17:HIS:HE1	1:A:184:GLU:OE1	1.89	0.55
1:F:152:LEU:HD21	1:F:179:PHE:HZ	1.73	0.53
1:B:171:LEU:HD21	1:C:167:GLN:OE1	2.10	0.52
1:A:5:ARG:HD3	3:A:402:HOH:O	2.10	0.52
1:C:145:GLN:NE2	1:F:157:THR:OG1	2.44	0.50
1:C:7:LEU:HB2	1:C:8:PRO:HD3	1.93	0.50
1:C:98:ASN:HD22	2:C:201:SAH:HB2	1.77	0.50
1:B:127[A]:MET:HE1	1:B:133:ILE:HG13	1.93	0.49
1:F:139:HIS:CE1	1:F:175:ASN:O	2.65	0.49
1:D:98:ASN:HB2	2:D:201:SAH:HG1	1.95	0.48
1:C:65:LYS:HG2	3:C:312:HOH:O	2.14	0.48
1:C:13:LEU:O	1:C:17:HIS:HD2	1.97	0.47
1:B:159:ASP:OD1	1:B:161:LYS:HD3	2.14	0.47
1:E:148:LYS:N	3:E:209:HOH:O	2.47	0.47
1:D:98:ASN:HD22	2:D:201:SAH:HB2	1.80	0.47
1:C:101:TYR:CE1	1:C:110:VAL:HG13	2.50	0.46
1:A:13:LEU:O	1:A:17:HIS:HD2	1.98	0.46
1:C:98:ASN:ND2	2:C:201:SAH:HN1	2.14	0.46
1:B:153:ASP:HB3	1:D:145:GLN:CD	2.41	0.46
1:E:175:ASN:O	1:E:175:ASN:CG	2.59	0.46
1:A:61:ARG:HD2	1:A:72:LEU:HD12	1.97	0.45
1:A:7:LEU:CD1	1:C:21:GLU:CG	2.95	0.45
1:C:127[B]:MET:HE1	1:C:133:ILE:HG13	1.98	0.45
1:A:173:GLN:HG2	3:E:252:HOH:O	2.17	0.45
1:A:148:LYS:HE2	1:A:179:PHE:CE1	2.52	0.44
1:C:55:LEU:HD12	1:C:56:ALA:N	2.32	0.44
1:F:130:GLU:HG2	3:F:269:HOH:O	2.16	0.44
1:B:98:ASN:ND2	2:B:201:SAH:HN1	2.17	0.43



Continued from previous page...

A 4 a sec 1	A4 0	Interatomic	Clash
Atom-1	Atom-2	$\operatorname{distance}\ (ext{\AA})$	overlap (Å)
1:D:1:MET:SD	1:F:164:GLN:HG3	2.59	0.43
1:A:164:GLN:HG3	1:E:1:MET:SD	2.58	0.43
1:C:29[A]:CYS:SG	1:C:57:LEU:HD23	2.58	0.43
1:D:127[B]:MET:CE	1:D:133:ILE:HG13	2.49	0.43
1:E:41:GLN:NE2	3:E:212:HOH:O	2.52	0.43
1:E:162:HIS:HD2	1:E:187:SER:O	2.02	0.43
1:D:127[B]:MET:HE1	1:D:133:ILE:HG13	2.00	0.42
1:A:135:LEU:HD13	1:A:151:LEU:HD21	2.01	0.42
1:D:25:VAL:O	1:D:95:ALA:HA	2.19	0.42
1:D:112:LYS:HD3	1:D:114:ASP:HB2	2.02	0.42
1:C:25:VAL:O	1:C:95:ALA:HA	2.19	0.42
1:A:25:VAL:O	1:A:95:ALA:HA	2.20	0.41
1:A:139:HIS:HE1	1:A:177:ALA:O	2.04	0.41
1:B:25:VAL:O	1:B:95:ALA:HA	2.20	0.41
1:E:139:HIS:HA	1:E:148:LYS:NZ	2.36	0.41
1:E:162:HIS:HB2	1:E:189:HIS:O	2.21	0.41
1:E:37:PHE:O	1:E:41:GLN:HG2	2.21	0.41
1:E:25:VAL:O	1:E:95:ALA:HA	2.22	0.40
1:B:164:GLN:HG3	1:C:1:MET:SD	2.62	0.40
1:C:130:GLU:HG2	3:C:319:HOH:O	2.20	0.40
1:F:25:VAL:O	1:F:95:ALA:HA	2.21	0.40
1:C:135:LEU:HD13	1:C:151:LEU:HD21	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
3:C:406:HOH:O	3:E:280:HOH:O[1_665]	2.17	0.03

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	171/195 (88%)	168 (98%)	3 (2%)	0	100 100
1	В	171/195 (88%)	170 (99%)	1 (1%)	0	100 100
1	С	185/195 (95%)	182 (98%)	3 (2%)	0	100 100
1	D	173/195 (89%)	171 (99%)	2 (1%)	0	100 100
1	E	165/195~(85%)	163 (99%)	2 (1%)	0	100 100
1	F	164/195 (84%)	161 (98%)	1 (1%)	2 (1%)	11 2
All	All	1029/1170 (88%)	1015 (99%)	12 (1%)	2 (0%)	44 22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	140	GLY
1	F	175	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	${f ntiles}$
1	A	157/171 (92%)	156 (99%)	1 (1%)	84	70
1	В	157/171 (92%)	156 (99%)	1 (1%)	84	70
1	С	167/171 (98%)	167 (100%)	0	100	100
1	D	158/171 (92%)	155 (98%)	3 (2%)	52	24
1	E	151/171 (88%)	147 (97%)	4 (3%)	41	14
1	F	150/171 (88%)	147 (98%)	3 (2%)	50	22
All	All	940/1026~(92%)	928 (99%)	12 (1%)	65	41

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	HIS
1	В	68	ASN
1	D	57	LEU



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type
1	D	68	ASN
1	D	77	HIS
1	Е	77	HIS
1	Е	115	THR
1	Е	118	GLN
1	Е	132	ILE
1	F	77	HIS
1	F	139	HIS
1	F	152	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (35) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	GLN
1	A	17	HIS
1	A	68	ASN
1	A A A B	139	HIS
1		33	ASN
1	В	41	GLN
1	В	68	ASN ASN
1	В	79	ASN
1	В	98	ASN
1	В	118	GLN
1	В	145	GLN
1	В	176	HIS
1	С	16	GLN
1	С	17	HIS
1	С	41	GLN
1	С	53	GLN
1	С	98	ASN
1	С	118	GLN
1	С	139	HIS
1	С	145	GLN
1	С	160	GLN
1	C C C C C C D D	33	ASN
1		41	GLN
1	D	82	HIS
1	D	98	ASN
1	D	176	HIS
1	Е	16	GLN
1	Е	41	GLN
1	Е	82	HIS



Continued from previous page...

Mol	Chain	Res	Type
1	Е	91	HIS
1	Е	162	HIS
1	Е	176	HIS
1	F	41	GLN
1	F	82	HIS
1	F	141	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trme	Chain Re	Dec	Res Link	Bond lengths			Bond angles		
MIOI	Mol Type		nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SAH	В	201	-	24,28,28	0.79	2 (8%)	25,40,40	0.97	2 (8%)
2	SAH	D	201	-	24,28,28	0.74	0	25,40,40	0.98	1 (4%)
2	SAH	A	201	-	24,28,28	0.74	1 (4%)	25,40,40	1.18	4 (16%)
2	SAH	С	201	-	24,28,28	0.68	0	25,40,40	0.98	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

\mathbf{Mol}	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAH	В	201	-	-	0/11/31/31	0/3/3/3
2	SAH	D	201	-	-	0/11/31/31	0/3/3/3
2	SAH	A	201	-	-	0/11/31/31	0/3/3/3
2	SAH	С	201	-	-	0/11/31/31	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(ext{\AA})$
2	A	201	SAH	OXT-C	-2.21	1.23	1.30
2	В	201	SAH	C8-N7	-2.17	1.30	1.34
2	В	201	SAH	OXT-C	-2.09	1.23	1.30

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	201	SAH	OXT-C-O	-3.38	116.41	124.09
2	С	201	SAH	O4'-C1'-C2'	-2.78	102.86	106.93
2	A	201	SAH	OXT-C-CA	2.67	122.47	113.38
2	С	201	SAH	C5-C6-N6	2.52	124.19	120.35
2	В	201	SAH	C5-C6-N6	2.47	124.11	120.35
2	A	201	SAH	O4'-C1'-C2'	-2.45	103.34	106.93
2	D	201	SAH	C5-C6-N6	2.08	123.51	120.35
2	В	201	SAH	O4'-C1'-C2'	-2.04	103.94	106.93
2	A	201	SAH	C5-C6-N6	2.02	123.42	120.35

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.


3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	201	SAH	1	0
2	D	201	SAH	2	0
2	С	201	SAH	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is



within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	#RSRZ>2	$OWAB(Å^2)$	Q < 0.9
1	A	175/195 (89%)	0.61	16 (9%) 16 16	9, 28, 58, 71	2 (1%)
1	В	175/195 (89%)	0.43	12 (6%) 24 25	10, 25, 51, 89	2 (1%)
1	С	184/195 (94%)	0.33	9 (4%) 36 38	8, 24, 43, 53	5 (2%)
1	D	175/195 (89%)	0.54	18 (10%) 13 13	10, 27, 52, 78	4 (2%)
1	Е	169/195 (86%)	0.55	18 (10%) 12 12	11, 27, 58, 88	2 (1%)
1	F	168/195 (86%)	0.62	13 (7%) 21 21	11, 28, 52, 82	2 (1%)
All	All	1046/1170 (89%)	0.51	86 (8%) 19 19	8, 27, 53, 89	17 (1%)

All (86) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	101	TYR	6.0
1	Е	139	HIS	5.6
1	Е	113	PRO	5.5
1	D	110	VAL	5.4
1	Е	189	HIS	5.1
1	A	110	VAL	4.7
1	F	141	HIS	4.4
1	F	148	LYS	4.3
1	A	176	HIS	4.3
1	D	144	GLY	4.3
1	В	144	GLY	4.2
1	Е	149	HIS	4.1
1	Е	176	HIS	3.8
1	В	139	HIS	3.7
1	С	188	GLY	3.7
1	В	109	ILE	3.6
1	D	188	GLY	3.6
1	D	55	LEU	3.5
1	С	145	GLN	3.4



Continued from previous page...

Mol	nued fron Chain	Res	Type	RSRZ
1	С	187	SER	3.3
1	С	101	TYR	3.3
1	В	146	ILE	3.1
1	D	176	HIS	3.1
1	Е	186	ILE	3.0
1	A	112	LYS	3.0
1	В	67	PHE	2.9
1	A	55	LEU	2.9
1	В	176	HIS	2.8
1	D	139	HIS	2.8
1	A	144	GLY	2.7
1	В	100	GLY	2.7
1	A	57	LEU	2.7
1	A	52	ILE	2.7
1	A	101	TYR	2.7
1	Е	148	LYS	2.7
1	A	139	HIS	2.6
1	С	171	LEU	2.6
1	A	145	GLN	2.6
1	Е	100	GLY	2.6
1	С	69[A]	HIS	2.6
1	Е	173	GLN	2.6
1	D	113	PRO	2.5
1	F	52	ILE	2.5
1	Е	175	ASN	2.5
1	D	146	ILE	2.4
1	С	139	HIS	2.4
1	В	175	ASN	2.4
1	F	125	SER	2.4
1	Е	115	THR	2.4
1	С	162[A]	HIS	2.4
1	A	67	PHE	2.4
1	D	114	ASP	2.4
1	F	118	GLN	2.3
1	Е	80	ILE	2.3
1	A	162	HIS	2.3
1	D	174	ARG	2.3
1	A	154	TYR	2.3
1	Е	177	ALA	2.3
1	F	173	GLN	2.3
1	A	85	ASN	2.3
1	F	31	ASN	2.3



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	В	145	GLN	2.3
1	Е	138	TYR	2.3
1	D	67	PHE	2.2
1	D	154	TYR	2.2
1	В	110	VAL	2.2
1	С	157	THR	2.2
1	В	177	ALA	2.2
1	A	82	HIS	2.2
1	Е	174	ARG	2.2
1	F	139	HIS	2.2
1	F	187	SER	2.2
1	Е	82	HIS	2.2
1	F	66	ASP	2.1
1	Е	118	GLN	2.1
1	D	56	ALA	2.1
1	A	146	ILE	2.1
1	D	173	GLN	2.1
1	D	115	THR	2.1
1	D	175	ASN	2.1
1	В	1	MET	2.0
1	F	152	LEU	2.0
1	D	100	GLY	2.0
1	D	112	LYS	2.0
1	Е	87	ALA	2.0
1	F	162	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	SAH	A	201	26/26	0.93	0.08	25,29,43,52	0
2	SAH	D	201	26/26	0.93	0.08	22,27,32,39	0
2	SAH	В	201	26/26	0.95	0.07	16,22,37,46	0
2	SAH	С	201	26/26	0.96	0.06	16,18,22,25	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around SAH D 201:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around SAH B 201:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

